

2008

# The Distribution of DMSP in Green Macroalgae from Northern New Zealand, Eastern Australia and Southern Tasmania

Kathryn L. Van Alstyne Dr.

Western Washington University, [kathy.vanalstyne@wwu.edu](mailto:kathy.vanalstyne@wwu.edu)

Follow this and additional works at: [https://cedar.wwu.edu/shannonpoint\\_facpubs](https://cedar.wwu.edu/shannonpoint_facpubs)

 Part of the [Marine Biology Commons](#)

---

## Recommended Citation

Van Alstyne, Kathryn L. Dr., "The Distribution of DMSP in Green Macroalgae from Northern New Zealand, Eastern Australia and Southern Tasmania" (2008). *Shannon Point Marine Center Faculty Publications*. 3.  
[https://cedar.wwu.edu/shannonpoint\\_facpubs/3](https://cedar.wwu.edu/shannonpoint_facpubs/3)

This Article is brought to you for free and open access by the Shannon Point Marine Center at Western CEDAR. It has been accepted for inclusion in Shannon Point Marine Center Faculty Publications by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).

# The distribution of DMSP in green macroalgae from northern New Zealand, eastern Australia and southern Tasmania

KATHRYN L. VAN ALSTYNE

Shannon Point Marine Center, Western Washington University, 1900 Shannon Point Road, Anacortes, WA 98221, USA

*The sulphonium compound dimethylsulphoniopropionate (DMSP) is commonly found in temperate green macroalgae. To examine taxonomic and regional and local geographical patterns of DMSP production in Australasian algae, I collected 30 species of green algae from 14 sites in three regions, eastern Australia, Tasmania, Australia, and the North Island of New Zealand. The distribution of DMSP content was similar to that seen from other areas of the world. DMSP was found in high concentrations in Ulva and Codium spp. It tended to be undetectable or in lower concentrations in other members of the orders Bryopsidales and Cladophorales. There was no evidence for differences in concentrations among the three regions in the genera Codium and Ulva; however, the invasive subspecies of Codium fragile, C. fragile ssp. tomentosoides, had significantly higher concentrations of DMSP than the non-invasive subspecies. The herbivorous sea slug Elysia maoria had whole body concentrations that were not significantly different from those of its host alga C. fragile ssp. tomentosoides. The distribution patterns of DMSP in Codium spp. do not support the hypothesis that DMSP is used as an anti-oxidant in this genus. Based on the data collected here and previous reports from the literature, I speculate that one function of DMSP in these algae may be to deter herbivores.*

**Keywords:** DMSP distribution, green macroalgae, *Codium* spp., northern New Zealand, eastern Australia, southern Tasmania

Submitted 26 April 2006; accepted 12 February 2008

## INTRODUCTION

The tertiary sulphonium compound dimethylsulphoniopropionate (DMSP) is commonly found in temperate green macroalgae (phylum Chlorophyta) and in the red algal (phylum Rhodophyta) genus *Polysiphonia* (Karsten *et al.*, 1990, 1992; Kirst *et al.*, 1991; Van Alstyne *et al.*, 2001b; Van Alstyne, 2008). It is also found in numerous phytoplankton (Kiene *et al.*, 1996) and zooxanthellate invertebrates, such as scleractinian corals, soft corals, anemones, corallimorphs (phylum Cnidaria) and in *Tridacna* spp. (phylum Mollusca) (Van Alstyne *et al.*, 2006 and references therein).

The reported functions of DMSP in macroalgae are diverse and vary by geographical region. For example, DMSP has been reported to be a cryoprotectant in Antarctic macroalgae (Kirst *et al.*, 1991; Karsten *et al.*, 1992), a compatible solute in north-eastern Atlantic macroalgae (Reed, 1983a, b; Edwards *et al.*, 1987, 1988), an antioxidant in south-western Atlantic algae (Ross & Van Alstyne, 2007), and as a precursor to an antiherbivore activated defence in north-eastern Pacific species (Van Alstyne *et al.*, 2001b; Van Alstyne & Houser, 2003). Although these functions are reported for different regions, it is likely that DMSP serves multiple functions in some, if not all, areas. It is also possible that the current geographical patterns are a result of there only being a handful of studies conducted

by researchers with different interests or expertise. The reported patterns may also be confounded by differences in the production of DMSP among algal taxa and by geographical variation in its production within taxa.

Patterns of spatial variation in algal natural products at scales of cms to 1000s of km have been described, both within species and across taxonomic groups (reviewed by Steinberg, 1992; Van Alstyne *et al.*, 2001a; Van Alstyne & Puglisi, 2007). When natural product concentrations are phenotypically plastic, these patterns can be generated by a variety of factors including responses to local environmental conditions (Karban & Baldwin, 1997; Tollrian & Harvell, 1999; Pigliucci, 2001). When they are produced at constitutive levels and dispersal is limited, local conditions can select for specific but different natural product levels at different sites. Alternatively, if dispersal is limited, natural product concentrations may become fixed at different levels at different sites by genetic drift.

Understanding the geographical and taxonomic patterns in the distribution of macroalgal DMSP concentrations at a range of scales could provide additional insights into the range of functions of the compound. For example, if DMSP plays an important role in protecting algae from reactive oxygen species (ROS) that are generated in response to physiological stresses (Sunda *et al.*, 2002; Ross & Van Alstyne, 2007), then algae in relatively stressful environments would be expected to produce higher concentrations of DMSP. Species growing higher in the intertidal zone would be expected to have higher concentrations of DMSP than algae growing in the lower intertidal or subtidal zone. Likewise, algae growing at lower latitudes would be expected to

### Corresponding author:

K.L. Van Alstyne

Email: kathy.vanalstyne@wwu.edu

produce higher concentrations of DMSP than species in more temperate areas because of the higher levels of ultraviolet (UV) radiation at lower latitudes.

The purpose of this study was to survey DMSP concentrations in green macroalgae in temperate Australasian waters to assess both taxonomic and local and regional-scale geographical patterns in the compound's distribution. Algae were collected from 14 sites in Australia and New Zealand and DMSP was measured in them. It was also measured in the herbivorous ascoglossan sea slug *Elysia maoria* Powell, 1937, which lives on and consumes siphonaceous green algae.

## MATERIALS AND METHODS

Marine green algae were collected by hand from intertidal or shallow subtidal habitats at five sites in northern New Zealand (Figure 1). These included: (1) the Goat Island Marine Reserve ( $36^{\circ} 16' S$ ,  $174^{\circ} 47' E$ ) near the University of Auckland's Leigh Marine Laboratory (LML) in Leigh, NZ; (2) Leigh Harbor ( $36^{\circ} 17' S$   $174^{\circ} 48' E$ ), located approximately 2 km from the LML site; (3) Army Bay ( $36^{\circ} 36' S$   $174^{\circ} 48' E$ ) on the south side of Whangaparoa Bay, approximately 35 km south of the LML site; and (4) Waitangi ( $35^{\circ} 15' S$   $174^{\circ} 04' E$ ), located approximately 75 km north of the LML site. A collection of *Ulva lactuca* L. was also made from the Mokohinau Islands ( $36^{\circ} 16' S$   $174^{\circ} 47' E$ ). The herbivorous sea slug *Elysia maoria* was noted to be occurring abundantly on *Codium fragile* ssp. *tomentosoides* (Van Goor) Silva at

Army Bay so several individuals were collected in order to compare DMSP concentrations in the slugs and their host alga.

In Australia, algae were collected by hand or with SCUBA from the high intertidal zone to 15 m depth from four sites in eastern Australia and five sites in Tasmania (Figure 1). Three of the eastern Australia sites were located within 16 km of one another in Sydney, New South Wales (NSW). The northernmost site was at Vaucluse Point ( $33^{\circ} 51' S$   $151^{\circ} 16' E$ ) in Jackson Harbor, the middle site was at Shark Point ( $33^{\circ} 55' S$   $151^{\circ} 16' E$ ), and the southernmost site was at Bare Island ( $34^{\circ} 00' S$   $151^{\circ} 14' E$ ) in Botany Bay. A fourth site was located at Jervis Bay, NSW ( $35^{\circ} 08' S$   $150^{\circ} 44' E$ ), approximately 125 km south of the Sydney sites. Sites in Tasmania included Bicheno Harbor ( $41^{\circ} 55' S$   $148^{\circ} 19' E$ ) and Rice Cove ( $41^{\circ} 55' S$   $148^{\circ} 19' E$ ) on the east side of the island, and on the south-eastern side, the blowhole at Blackman's Bay ( $43^{\circ} 01' S$   $147^{\circ} 20' E$ ), Adventure Bay ( $43^{\circ} 16' S$   $147^{\circ} 21' E$ ) on Bruny Island, and a location 1 km east of the Nine Pin Point marine reserve ( $43^{\circ} 17' S$   $147^{\circ} 11' E$ ) near Verona Sands. The three southernmost sites were located within 35 km of one another and approximately 150 to 180 km from the two southern sites, which were approximately 2 km apart.

The algae were transported to either the Leigh Marine Laboratory in Leigh, New Zealand or the University of New South Wales in Sydney, Australia and processed within about 2 days of collection. They were identified to species or subspecies whenever possible using Womersley (1984), Adams (1994) and Huisman (2000). Approximately 0.1 g pieces of each alga were dried in a drying oven at  $60^{\circ} C$




Fig. 1. Collection sites in Australia and New Zealand. (A) North Island of New Zealand (NZ), LML, Leigh Marine Laboratory, Goat Island Reserve, Leigh; (B) eastern Australia; (C) Tasmania, Australia.

overnight. The dried algae were then shipped to the Shannon Point Marine Center (SPMC) in Anacortes, WA, USA for measurement of DMSP. Previous studies have shown that drying green algae prior to measuring tissue DMSP concentrations is preferable to measuring DMSP in fresh tissues (Karsten *et al.*, 1994; Van Alstyne *et al.*, 2003).

At SPMC, the dried algae and opisthobranchs were weighed to the nearest 0.1 mg and placed in 4 ml of 4 N NaOH in 30 ml gas-tight containers. Known amounts of commercially-obtained DMSP (Center for Analysis, Spectroscopy and Synthesis, University of Groningen; purity > 98%) were added to 4 N NaOH in gas-tight containers to generate standard curves. All containers were incubated in the dark at 4°C overnight to allow for the conversion of DMSP to DMS. DMSP was measured as DMS by directly injecting 10–20 µl of headspace gas into an SRI gas chromatograph (Chromasil 330 column, flame photometric detector; oven temperature 90°C; lower detection limit: 5 µg DMS).

Analyses at the ordinal level for the green algae were based on the classification of these organisms given on the AlgaeBase website (<http://www.algaebase.org/>) on 1 November 2007. A two-way analysis of variance was conducted to examine the effects of region (New Zealand, eastern Australia and Tasmania) and genus on DMSP concentrations in *Ulva* ssp. and *Codium* ssp. that contained detectable amounts of DMSP after ascertaining that the data met the assumption of normality. A Student's *t*-test was used to compare DMSP concentrations in *E. maoria* and its host alga *C. fragile* ssp. *tomentosoides*.

## RESULTS

Concentrations of DMSP ranged from being undetectable in many algae in the orders Bryopsidales and Cladophorales to 3.7% of the alga's dry mass (DM) or 278 µmol g<sup>-1</sup> DM (Tables 1–3). The presence of DMSP tended to follow

taxonomic affinities rather than varying by geographical location. A two-way analysis of variance with region (eastern Australia, Tasmania and New Zealand) and genus (*Codium* and *Ulva*) as factors indicated that the effects of region were not significant ( $F = 1.05$ ,  $P = 0.405$ ), but the effects of genus were ( $F = 8.11$ ,  $P = 0.029$ ). There was no significant interaction effect ( $F = 0.42$ ,  $P = 0.672$ ).

In the Bryopsidales, DMSP was absent in all species of *Caulerpa*, but was present in detectable quantities in two-thirds of the species of *Codium* examined. It was uniformly absent in *Codium convolutum* (Dell) Silva, which was collected in the intertidal zone (Table 1). In *Codium fragile* (Suringar) Hariot, DMSP was present in significantly higher amounts in the invasive subspecies *C. fragile* ssp. *tomentosoides*, which was collected in low intertidal to shallow subtidal waters in New Zealand, than the non-invasive *C. fragile*, which was collected in Tasmania and eastern Australia (Student's  $t = 2.98$ ,  $P = 0.041$ ,  $N = 5$  species).

In the order Cladophorales, concentrations of DMSP varied among species, even within genera (Tables 1–3). DMSP occurred in high concentrations in an unidentified species of *Chaetomorpha* from Leigh Harbor, New Zealand, in low concentrations in *Chaetomorpha aerea* (Dillwyn) Kuetzing from the seawater outflow at the Leigh Marine Laboratory, and was not detected in a population of *C. aerea* from Army Bay (Table 1). It was also not detected in *Chaetomorpha coliformis* (Montagne) Kuetzing from three sites on Tasmania (Table 3). In *Cladophora*, DMSP was not detected in four out of the eight species collected. It occurred in low concentrations in three of the remaining species and in high concentrations in a single species collected from the seawater outflow at the Leigh Marine Laboratory (Table 1). DMSP occurred in relatively high concentrations in all members of the order Ulvales.

Whole animal concentrations of DMSP in *Elysia maoria* were  $95 \pm 22$  µmol g<sup>-1</sup> DM or  $1.3 \pm 0.3\%$  DM ( $N = 10$ ). These

**Table 1.** DMSP concentrations in New Zealand macroalgae. Values are given in µmol g<sup>-1</sup> tissue dry mass (DM) and as the per cent of the tissue DM to facilitate comparisons with the literature. All collections were made in 2002. LML, Leigh Marine Laboratory; nd, not detected.

Species	Location	Collection date	[DMSP] (µmol g <sup>-1</sup> DM)	[DMSP] (% DM)	N
<b>PHYLUM CHLOROPHYTA</b>					
Order Bryopsidales					
<i>Caulerpa geminata</i>	Goat Island Reserve	13 February	nd	nd	5
<i>Codium convolutum</i> (intertidal)	Army Bay	8 February	nd	nd	10
<i>Codium convolutum</i> (intertidal)	Leigh Harbor	6 February	nd	nd	10
<i>Codium convolutum</i> (intertidal)	Waitangi Park, Bay of Islands	10 February	nd	nd	10
<i>Codium fragile</i> ssp. <i>tomentosoides</i>	Army Bay	8 February	110 ± 28	1.5 ± 0.4	20
<i>Codium fragile</i> ssp. <i>tomentosoides</i>	Bay of Islands	10 February	118 ± 25	1.6 ± 0.3	20
<i>Codium fragile</i> ssp. <i>tomentosoides</i>	Leigh Harbor	7 February	125 ± 29	1.7 ± 0.4	20
<i>Codium spongiosum</i>	Leigh Harbor	6 February	35 ± 20	0.5 ± 0.3	20
Order Cladophorales					
<i>Chaetomorpha aerea</i>	Army Bay	8 February	nd	nd	10
<i>Chaetomorpha aerea</i>	Seawater Outflow, LML	11 February	19 ± 8	0.3 ± 0.1	15
<i>Chaetomorpha</i> sp.	Leigh Harbor	7 February	142 ± 41	1.9 ± 0.6	20
<i>Cladophora</i> sp. 1	Leigh Harbor	6 February	5 ± 5	0.1 ± 0.1	20
<i>Cladophora</i> sp. 2	Seawater Outflow, LML	11 February	61 ± 22	0.9 ± 0.3	20
Order Ulvales					
<i>Ulva intestinalis</i>	Seawater Outflow, LML	6 February	120 ± 40	1.6 ± 0.6	20
<i>Ulva intestinalis</i>	Tide pool below LML	6 February	168 ± 32	2.3 ± 0.4	20
<i>Ulva lactuca</i>	Goat Island Reserve	13 February	192 ± 61	2.6 ± 0.8	20
<i>Ulva lactuca</i>	Mokohinau Islands		103 ± 19	1.4 ± 0.3	5
<i>Ulva lactuca</i>	Seawater Outflow, LML	7 February	187 ± 35	2.5 ± 0.5	20

**Table 2.** DMSP concentrations of algae collected from eastern Australia. Values are given in  $\mu\text{mol g}^{-1}$  tissue dry mass (DM) and as the per cent of the tissue DM to facilitate comparisons with the literature. All collections were made in 2002. nd, not detected.

Species	Location	Collection date	[DMSP] ( $\mu\text{mol g}^{-1}$ DM)	[DMSP] (% DM)	N
<b>PHYLUM CHLOROPHYTA</b>					
Order Bryopsidales					
<i>Caulerpa filiformis</i>	Vaucluse Point, Sydney	19 February	nd	nd	5
<i>Caulerpa flexilis</i>	Jervis Bay	4 March	nd	nd	5
<i>Caulerpa taxifolia</i>	Sydney	27 February	nd	nd	5
<i>Codium fragile</i>	Jervis Bay	4 March	$88 \pm 15$	$1.2 \pm 0.2$	10
<i>Codium fragile</i>	Vaucluse Point, Sydney	19 February	$47 \pm 24$	$0.6 \pm 0.3$	20
Order Cladophorales					
<i>Cladophora coelothrix</i>	Jervis Bay	4 March	nd	nd	20
<i>Cladophora feredayi</i>	Bare Island, Sydney	22 February	nd	nd	20
<i>Cladophora</i> sp. 1	Bare Island, Sydney	22 February	$8 \pm 12$	$0.1 \pm 0.2$	10
<i>Cladophora</i> sp. 2	Shark Point, Sydney	21 February	$12 \pm 7$	$0.2 \pm 0.1$	20
Order Ulvales					
<i>Ulva intestinalis</i>	Bare Island, Sydney	22 February	$150 \pm 5$	$2.0 \pm 0.1$	20
<i>Ulva intestinalis</i>	Jervis Bay	4 March	$127 \pm 29$	$1.7 \pm 0.4$	20
<i>Ulva intestinalis</i>	Shark Point, Sydney	21 February	$193 \pm 25$	$2.6 \pm 0.3$	20
<i>Ulva intestinalis</i>	Vaucluse Point, Sydney	19 February	$78 \pm 37$	$1.1 \pm 0.5$	10
<i>Ulva lactuca</i>	Shark Point, Sydney	21 February	$278 \pm 51$	$3.7 \pm 0.7$	20

values were not significantly different from the DMSP concentrations of  $110 \pm 28 \mu\text{mol g}^{-1}$  DM or  $1.5 \pm 0.4\%$  DM in the *Codium fragile* ssp. *tomentosoides* from which they were collected (Student's *t*-test:  $t = 1.72$ ,  $P = 0.10$ ).

## DISCUSSION

The patterns of DMSP concentrations of Australian and New Zealand marine macroalgae are similar to those seen in other

**Table 3.** DMSP concentrations of algae collected from Tasmania, Australia. Values are given in  $\mu\text{mol g}^{-1}$  tissue dry mass (DM) and as the per cent of the tissue DM to facilitate comparisons with the literature. All collections were made in 2002. nd, not detected.

Species	Location	Collection date	[DMSP] ( $\mu\text{mol g}^{-1}$ DM)	[DMSP] (% DM)	N
<b>PHYLUM CHLOROPHYTA</b>					
Order Bryopsidales					
<i>Caulerpa brownii</i>	Bruny Island, Tasmania	28 February	nd	nd	5
<i>Caulerpa brownii</i>	Rice Cove, Tasmania	27 February	nd	nd	5
<i>Caulerpa geminata</i>	Nine Pin Point, Tasmania	28 February	nd	nd	5
<i>Caulerpa geminata</i>	Rice Cove, Tasmania	27 February	nd	nd	5
<i>Caulerpa longifolia</i>	Rice Cove, Tasmania	27 February	nd	nd	5
<i>Caulerpa scapelliformis</i>	Nine Pin Point, Tasmania	28 February	nd	nd	5
<i>Caulerpa simpliciusaeta</i>	Nine Pin Point, Tasmania	28 February	nd	nd	5
<i>Codium dimorphum</i>	Bruny Island, Tasmania	28 February	$65 \pm 20$	$0.9 \pm 0.3$	10
<i>Codium fragile</i>	Bicheno Harbor, Tasmania	27 February	$8 \pm 5$	$0.1 \pm 0.1$	10
<i>Codium fragile</i> ssp. <i>tasmanicum</i>	Blackman's Bay, Tasmania	28 February	$10 \pm 16$	$0.1 \pm 0.2$	10
<i>Codium harveyi</i>	Bicheno Harbor, Tasmania	27 February	$66 \pm 41$	$0.9 \pm 0.6$	10
<i>Codium</i> sp.	Rice Cove, Tasmania	27 February	nd	nd	20
Order Cladophorales					
<i>Chaetomorpha coliformis</i>	Bicheno Harbor, Tasmania	27 February	nd	nd	20
<i>Chaetomorpha coliformis</i>	Bruny Island, Tasmania	28 February	nd	nd	10
<i>Chaetomorpha coliformis</i>	Nine Pin Point, Tasmania	28 February	nd	nd	20
<i>Cladophora feredayi</i>	Blackman's Bay, Tasmania	28 February	nd	nd	20
<i>Cladophora subsimplex</i>	Bicheno Harbor, Tasmania	27 February	nd	nd	10
<i>Rhizoclonium riparium</i>	Bruny Island, Tasmania	28 February	nd	nd	10
Order Siphonocladales					
<i>Apjohnia laetevirens</i>	Blackman's Bay, Tasmania	28 February	nd	nd	10
Order Ulvales					
<i>Ulva intestinalis</i>	Bicheno Harbor, Tasmania	27 February	$99 \pm 19$	$1.3 \pm 0.3$	10
<i>Ulva intestinalis</i>	Blackman's Bay, Tasmania	28 February	$178 \pm 31$	$2.4 \pm 0.4$	20
<i>Ulva intestinalis</i>	Bruny Island, Tasmania	28 February	$155 \pm 19$	$2.1 \pm 0.3$	20
<i>Ulva australis</i>	Bicheno Harbor, Tasmania	27 February	$82 \pm 20$	$1.1 \pm 0.3$	20
<i>Ulva taeniata</i>	Bicheno Harbor, Tasmania	27 February	$149 \pm 23$	$2 \pm 0.3$	20
<i>Ulva taeniata</i>	Blackman's Bay, Tasmania	28 February	$124 \pm 22$	$1.7 \pm 0.3$	20
<i>Ulva taeniata</i>	Bruny Island, Tasmania	28 February	$146 \pm 16$	$2.0 \pm 0.2$	20
<i>Ulva taeniata</i>	Nine Pin Point, Tasmania	28 February	$102 \pm 26$	$1.4 \pm 0.4$	20

regional geographical comparisons. The general pattern is that DMSP is present in high concentrations in members of the order Ulvales, in some members of the Bryopsidales and Ulotrichales, and in the red algal genus *Polysiphonia*. It is absent or in very low concentration in red algae except *Polysiphonia* spp., brown algae, siphonaceous green algae except most species of *Codium* and *Acrosiphonia* and cyanobacteria (Van Alstyne & Puglisi, 2007). A few of the algae within these groups have low concentrations of DMSP but it is not known whether the source of the DMSP is the algae themselves or epiphytic microalgae growing on them.

The concentrations of DMSP in *Ulva* reported in this survey were comparable to values reported in *Ulva* from the northern hemisphere. In this study, DMSP in *Ulva* spp. ranged from 78 to 278  $\mu\text{mol g}^{-1}$  DM, which is similar to concentrations of DMSP in *Ulva* spp. from northern hemisphere warm and cold temperate sites (Van Alstyne & Puglisi, 2007). In *Ulva intestinalis*, local differences in DMSP concentrations among sites were as great as the differences in concentrations between sites from New Zealand, eastern Australia and Tasmania. This suggests that DMSP concentrations in these algae may be responding to localized environmental conditions, such as differences in salinity or grazer densities, rather than large scale environmental gradients.

Although DMSP concentrations in marine macroalgae tend to increase with increasing latitude in the northern hemisphere over a range of geographical and taxonomic scales (Van Alstyne *et al.*, 2007; Van Alstyne & Puglisi, 2007), I saw no evidence that similar patterns of DMSP concentrations occurred in the algae examined here. In the northern hemisphere, DMSP concentrations in *Ulva* spp. increase from the tropics to polar regions (Van Alstyne & Puglisi, 2007) and within *Ulva lactuca* over a 70 km change in latitude (Van Alstyne *et al.*, 2007). Here, we found no tendency for DMSP to differ among the three regions examined, even though the sampling sites from Tasmania were about 1000 km south of the eastern Australia and New Zealand sites. At this point, there are insufficient data available to determine whether geographical clines in macroalgal DMSP concentration that are seen in the northern hemisphere are absent in the southern hemisphere.

Members of the genus *Codium* spanned a range of tidal heights making it possible to test the hypothesis that DMSP is used to counter environmental stresses. If this hypothesis were true, then DMSP should be higher in species growing higher in the intertidal zone. In this study, I found that *Codium convolutum*, which I collected from the intertidal zone, did not contain detectable amounts of DMSP, whereas *Codium fragile*, which I collected from lower in the intertidal zone and in the subtidal zone, consistently contained DMSP, usually in relatively high quantities. With the exception of an unidentified species of *Codium* from Tasmania (Table 3), the several other species of *Codium* that were collected from the low intertidal zone to the subtidal zone also contained measurable amounts of DMSP; however, for these species, only single collections were made so it is not known how much inter-population variability exists in their DMSP concentrations. These data suggest that DMSP, at least in *Codium* spp., is not functioning to protect the algae from stresses such as desiccation and extreme temperatures that are found in the upper intertidal zone.

The presence of higher DMSP concentrations in the invasive subspecies of *Codium fragile*, *C. fragile* ssp. *tomentosoides*,

may in part be responsible for the success of the invasive alga. DMSP has been shown to be the precursor to an activated anti-herbivore defence that can deter feeding by some sea urchins (Van Alstyne *et al.*, 2001b; Van Alstyne & Houser, 2003; Lyons *et al.*, 2007). In Nova Scotia, Canada, DMSP in *C. fragile* ssp. *tomentosoides* occurs in high enough concentrations to deter feeding by green sea urchins and is thought to contribute to the alga's ability to become established and persist in former kelp beds that are heavily grazed (Lyons *et al.*, 2007).

Indirect evidence suggests that DMSP may also protect *C. fragile* ssp. *tomentosoides* from grazing in New Zealand. Although many species of herbivorous invertebrates inhabit the shores of the North Island of New Zealand, they are generally uncommon on *C. fragile* ssp. *tomentosoides* (Trowbridge, 1995). When local grazers were offered choices between fresh tissues of *C. fragile* ssp. *tomentosoides* and other sympatric species, they usually exhibited distinct preferences, although whether they preferred or avoided *C. fragile* ssp. *tomentosoides* was herbivore-specific (Trowbridge, 1995). However, when the algae were freeze-dried, ground, and incorporated into an agar-based food, almost all the preferences disappeared. Trowbridge (1995) attributed the loss of preference to a loss of morphology or volatile chemicals in the artificial foods and noted that the *C. fragile* ssp. *tomentosoides* used in these assays had a strong pungent odour, which is consistent with the smell of DMS, one of the cleavage products of DMSP. Because DMSP is lost during freeze-drying, it is very likely that the artificial foods in Trowbridge's (1995) experiments contained little or none of the DMSP that was present in the fresh algae. If DMSP or its cleavage products were causing the preferences exhibited for the fresh algae, then a loss of DMSP due to freeze-drying in the ground *C. fragile* ssp. *tomentosoides* would explain the loss of preferences when herbivores were offered the agar-based diets.

The relatively high concentrations of DMSP found in the herbivorous sea slug *Elysia maoria* suggest that the animal is taking up DMSP from its host alga *C. fragile* ssp. *tomentosoides*. Many fish and shellfish species are reported to have high tissue concentrations of DMSP that are suspected to be derived from their diets (e.g. Ackman & Hingley, 1968; Brooke *et al.*, 1968; Ackman *et al.*, 1972; Iida & Tokunaga, 1986; White *et al.*, 1995). Ascoglossan opisthobranchs, including other species in the family Elysiidae, are known to take up algal components from their foods, including chloroplasts and secondary metabolites (e.g. Paul & Van Alstyne, 1988; Williams & Walker, 1999; Trowbridge & Todd, 2001). Therefore, it is not surprising that *E. maoria* contains DMSP in concentrations that are comparable to those in its host alga. An ascoglossan in the family Elysiidae from Guam has been shown to take up compounds from the host algae, which are then modified, stored, and emitted when the animal is perturbed (Paul & Van Alstyne, 1988). The compound produced by the sea slug is a deterrent towards both herbivorous and carnivorous fish and is presumably being used as an anti-predator defence. Whether *E. maoria* uses sequestered DMSP as a defence is not currently known.

In summary, DMSP is found in many species of macroalgae and in an herbivorous sea slug in Australia and New Zealand. The taxonomic distribution of DMSP is comparable to that seen in other areas of the world; however, there was no evidence that the latitudinal clines that have been described in the northern hemisphere are present in this region. This may

reflect the fact that there are currently fewer data available on the geographical distribution of DMSP in the southern hemisphere than the northern hemisphere. There is some support for DMSP functioning as an anti-herbivore defence in Australasia; however, further experimental work is needed to conclusively demonstrate this.

## ACKNOWLEDGEMENTS

I would like to thank P. Steinberg and R. Taylor for hosting my visits to Australia and New Zealand, respectively. I thank the members of the Steinberg Laboratory and the faculty and staff at the Leigh Marine Laboratory for their invaluable assistance with many of the logistical aspects of this project. I would particularly like to acknowledge N. Paul for his assistance with fieldwork in Tasmania. This work was funded by a grant from the National Science Foundation (IOB-0090825).

## REFERENCES

- Ackman R. and Hingley H.** (1968) The occurrence and retention of dimethyl- $\beta$ -propiothetin in some filter-feeding animals. *Journal of the Fisheries Research Board of Canada* 25, 267–284.
- Ackman R., Hingley J. and MacKay K.** (1972) Dimethyl sulfide as an odor component in Nova Scotia fall mackerel. *Journal of the Fisheries Research Board of Canada* 29, 1085–1088.
- Adams N.M.** (1994) *Seaweeds of New Zealand*. Christchurch: Canterbury University Press.
- Brooke R., Mendelsohn J. and King F.** (1968) Significance of dimethyl sulfide to the odor of soft-shell clams. *Journal of the Fisheries Research Board of Canada* 25, 2453–2460.
- Edwards D.M., Reed R.H., Chudek J.A., Foster R. and Stewart W.D.P.** (1987) Organic solute accumulation in osmotically-stressed *Enteromorpha intestinalis*. *Marine Biology* 95, 583–592.
- Edwards D.M., Reed R.H. and Stewart W.D.P.** (1988) Osmoacclimation in *Enteromorpha intestinalis*: long-term effects of osmotic stress on organic solute accumulation. *Marine Biology* 88, 457–476.
- Huisman J.** (2000) *Marine plants of Australia*. Nedlands: University of Western Australia Press.
- Iida H. and Tokunaga T.** (1986) Dimethyl sulfide and dimethyl- $\beta$ -propiothetin in shellfish. *Bulletin of the Japanese Society of Scientific Fisheries* 52, 557–563.
- Karban R. and Baldwin I.** (1997) *Induced responses to herbivory*. Chicago: University of Chicago Press.
- Karsten U., Kuck K., Daniel C., Wiencke C. and Kirst G.O.** (1994) A method for complete determination of dimethylsulfoniopropionate (DMSP) in marine macroalgae for different geographical regions. *Phycologia* 33, 171–176.
- Karsten U., Wiencke C. and Kirst G.O.** (1990) A beta-dimethylsulphoniopropionate (DMSP) content of macroalgae from Antarctica and Southern Chile. *Botanica Marina* 33, 143–146.
- Karsten U., Wiencke C. and Kirst G.O.** (1992) Dimethyl sulphoniopropionate (DMSP) accumulation in green macroalgae from polar to temperate regions: interactive effects of light versus salinity and light versus temperature. *Polar Biology* 12, 603–607.
- Kiene R.P., Visscher P., Keller M. and Kirst G.O.** (1996) *Biological and environmental chemistry of DMSP and related sulfonium compounds*. New York: Plenum Press.
- Kirst G.O., Thiel C., Wolff H., Nothnagel J., Wanzek M. and Ulmke R.** (1991) Dimethylsulphoniopropionate (DMSP) in ice-algae and its possible biological role. *Marine Chemistry* 35, 381–388.
- Lyons D., Van Alstyne K.L. and Scheibling R.** (2007) Anti-grazing activity and seasonal variation of dimethylsulphoniopropionate (DMSP)-associated compounds in the invasive alga *Codium fragile* ssp. *tomentosoides*. *Marine Biology*, 153, 179–188.
- Paul V.J. and Van Alstyne K.L.** (1988) Use of ingested algal diterpenoids by *Elysia halimeda* Macnae (Opisthobranchia: Ascoglossa) as antipredator defenses. *Journal of Experimental Marine Biology and Ecology* 119, 15–29.
- Pigliucci M.** (2001) *Phenotypic plasticity: beyond nature and nurture*. Baltimore: Johns Hopkins University Press.
- Reed R.H.** (1983a) Measurement and osmotic significance of beta-dimethylsulphoniopropionate in marine macroalgae. *Marine Biology Letters* 4, 173–181.
- Reed R.H.** (1983b) The osmotic responses of *Polysiphonia lanosa* (L.) Tandy from marine and estuarine sites: evidence for incomplete recovery of turgor. *Journal of Experimental Marine Biology and Ecology* 68, 169–193.
- Ross C. and Van Alstyne K.L.** (2007) Intraspecific variation in stress-induced hydrogen peroxide scavenging by the ulvoid macroalga *Ulva lactuca*. *Journal of Phycology* 43, 466–474.
- Steinberg P.D.** (1992) Geographical variation in the interaction between marine herbivores and brown algal secondary metabolites. In Paul V. (ed.) *Ecological roles of marine natural products*. Ithaca: Cornell University Press, pp. 51–92.
- Sunda W.G., Kieber D., Kiene R. and Huntsman S.** (2002) An antioxidant function for DMSP and DMS in marine algae. *Nature* 418, 317–320.
- Tollrian R. and Harvell C.D.** (1999) *The ecology and evolution of inducible defenses*. Princeton: Princeton University Press.
- Trowbridge C.D.** (1995) Establishment of the green alga *Codium fragile* ssp. *tomentosoides* on New Zealand rocky shores: current distribution and invertebrate grazers. *Journal of Ecology* 83, 949–965.
- Trowbridge C.D. and Todd C.** (2001) Host-plant change in marine specialist herbivores: ascoglossan sea slugs on introduced macroalgae. *Ecological Monographs* 71, 219–243.
- Van Alstyne K.L.** (2008) Ecological and physiological roles of dimethylsulphoniopropionate (DMSP) and its products in marine macroalgae. In Amsler C.D. (ed.) *Algal chemical ecology*. Heidelberg: Springer, pp. 173–194.
- Van Alstyne K.L., Dethier M.N. and Duggins D.O.** (2001a) Spatial patterns in macroalgal chemical defenses. In McClintock J. and Baker W. (eds) *Marine chemical ecology*. Boca Raton, FL: CRC Press, pp. 301–324.
- Van Alstyne K.L. and Houser L.T.** (2003) Dimethylsulfide release during macroinvertebrate grazing and its role as an activated chemical defense. *Marine Ecology Progress Series* 250, 175–181.
- Van Alstyne K.L., Koellermaier L. and Nelson T.** (2007) Spatial variation in dimethylsulphoniopropionate (DMSP) production in *Ulva lactuca* (Chlorophyta) from the northeast Pacific. *Marine Biology* 150, 1127–1135.
- Van Alstyne K.L., Pelletreau K.N. and Rosario K.** (2003) The effects of salinity on dimethylsulphoniopropionate production in the green alga *Ulva fenestrata* Postels et Ruprecht (Chlorophyta). *Botanica Marina* 46, 350–356.
- Van Alstyne K.L. and Puglisi M.P.** (2007) DMSP in marine macroalgae and macroinvertebrates: distribution, function, and ecological impacts. *Aquatic Science* 69, 394–402.

- Van Alstyne K.L., Schupp P. and Slattery M.** (2006) The distribution of dimethylsulfoniopropionate (DMSP) in tropical Pacific coral reef invertebrates. *Coral Reefs* 25, 321–327.
- Van Alstyne K.L., Wolfe G.V., Freidenburg T.L., Neill A. and Hicken C.** (2001b) Activated defense systems in marine macroalgae: evidence for an ecological role for DMSP cleavage. *Marine Ecology Progress Series* 213, 53–65.
- White B., Hill R.W. and Dacey J.W.H.** (1995) Accumulation of DMSP in *Geukensia demissa* depends on trophic interactions. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 189, 235–236.
- Williams S. and Walker D.I.** (1999) Mesoherbivore–macroalgal interactions: feeding ecology of sacoglossan sea slugs (Mollusca, Opisthobranchia) and their effects on their food algae. *Oceanography and Marine Biology: an Annual Review* 37, 87–128.
- and
- Womersley H.B.S.** (1984) *The marine benthic flora of Southern Australia. Part I.* Adelaide: South Australia Printing Division.

**Correspondence should be addressed to:**

Kathryn L. Van Alstyne  
Shannon Point Marine Center  
Western Washington University  
1900 Shannon Point Road, Anacortes, WA 98221, USA  
email: kathy.vanalstyne@wwu.edu