

Assessing the Effects of Chemical Mixtures using a Bayesian Network- Relative Risk Model (BN-RRM) Integrating Adverse Outcome Pathways (AOPs)

Chu, V.R.*¹, Harris, M.J.¹, Graham, S. E.¹, Wallis, L.K.¹, Mitchell, C.J.², Stark, J.D.², von Stackleberg, K.E.³ and Landis, W.G.¹

¹Western Washington University, Bellingham, WA 98225, ²Washington State University- Puyallup Research and Extension Center, Puyallup, WA 98371, ³Harvard Center for Risk Analysis, Boston, MA 02115; Supported by USEPA STAR Grant RD-83569501; Email:chuv@wwu.edu


Introduction

There are long-standing uncertainties about toxicity of chemical mixtures to populations. Laboratory toxicity tests have confirmed synergistic and antagonistic effects to individuals, but not to populations. We will conduct a regional scale ecological risk assessment by evaluating the effects chemical mixtures to populations with a new Bayesian Network- Relative Risk Model (BN-RRM) incorporating an Adverse Outcome Pathway (AOP) (Figure 7). We started applying this new BN-RRM framework in a case study with organophosphate pesticide (OP) mixtures (diazinon, chlorpyrifos, and malathion). Acetylcholinesterase inhibition (AChE) was chosen the molecular initiating event and the Puget Sound Chinook salmon (*Oncorhynchus tshawytscha*) and Coho salmon (*Oncorhynchus kisutch*) Evolutionary Significant Units (ESU) were chosen as population endpoints.


Background

Malathion and chlorpyrifos (Figure 1 and 2) make up of 50% of OP pesticides detected in urban streams in the Pacific Northwest [9]. Urban streams are where Coho and Chinook salmon inhabit and spawn and the streams typically contain two or more OPs [1] [9]. Because mixtures are normally encountered in urban streams, toxicity testing of OP mixtures (Figure 3) is comparable to environmentally realistic exposures [4] [5] [9].


Organophosphate pesticides such as chlorpyrifos and malathion act synergistically to Chinook salmon. The presence of an organophosphate compound such as chlorpyrifos inhibits the enzyme carboxylesterase (CaE) which results in the potentiation of malaoxon, a metabolite of malathion, leading to synergism [3] [6].



Figures 1, 2 and 3. Dose response curves for the individual chemicals and mixtures will be combined to fit one dose-response surface.




Data from Laetz et al. (2009)



The conversion from chemical concentrations to moles will help facilitate the fit.

Figure 4. Isoboles can account for mixtures that fail the assumption of additivity.



This isobole indicates synergistic toxicity. The point is the EC50 of this mixture and the error bars are the confidence intervals.

Isobole Model Design

Isoboles are contours that can connect two chemicals of equal effect into one response surface (Figure 4). If the isoboles are linear, then the mixture indicates concentration addition. Isoboles which curve towards the origin indicate synergistic mixtures; while ones that curve away from the origin are antagonistic [11]. The next steps will be to create an isobologram and fit mixture rays to a model.

NOTE: Probabilities of population estimations from the organophosphate pesticide mixtures have **NOT** been compiled yet. However, results from single pesticide exposures have been compiled.




Figure 5 and 6. Sensitivity analysis for toxicological effects and water temperature by watershed per simulation year

Sensitivity Analysis for Single Organophosphate Pesticides

In our model, toxicological effects did not have much of an effect on salmon populations (Figure 5). This is not surprising because our model only accounted for single chemicals. Toxicological effects were more sensitive depending on the watershed. Total toxicological effects are more important in the Cedar and Lower Skagit watersheds for all years. Water temperature (Figure 6) is more important in the Nooksack and Lower Yakima watersheds.

Bayesian Network Depicting the Probability of Chinook Population Size





Figure 8. Four study sites will be examined, three in the Puget Sound basin and one in the Columbia river watershed.



Uncertainty Analysis for Single Organophosphate Pesticides

We attempted to make migration nodes to connect olfaction to adult reduction in survival because these OP pesticides have been found to inhibit olfaction [7] [10] [12]. Olfactory inhibition can adversely affect the ability of salmon to return to spawn. However, there is not a clearly defined dose-response relationship between what the role olfaction inhibition plays in homing ability. Therefore, our confidence in the available dataset is low.

We also examined code wire tag (CWT) data to calculate rates of homing and straying during migration. Estimates of smolt-adult return (SAR) percent rates and hatchery escapement numbers were calculated. Our analysis indicates that the CWT data are not sufficient to estimate SAR and escapement rates for our study areas because no percent error rates were given.

What is Next?

- We expect to see a **greater** contribution of toxicological effects with the mixtures
- We will add suitable habitat nodes from Geographic Information Systems (GIS) and other data sets. Adults pick suitable habitats to spawn based on discharge, substrate, streambank condition, riparian vegetation as well as cover & refuge [2]
- Climate change will also be added as a stressor

See also:

Landis et al. (2017) Integrating Adverse Outcome Pathways into the Bayesian Network Relative Risk Model for Landscape Scale Ecological Risk Assessment

Mitchell et al. (2017) Incorporating Spatially Explicit Metapopulation Models as the Adverse Outcome Pathway Endpoint of a Bayesian Network-Relative Risk Model.

References

[1] Anderson PD, Dugger D. 2008. Surface Water Monitoring Program for Pesticides in Salmonid-Bearing Streams, 2007 Data Summary. Washington State Department of Ecology, Olympia Washington.

[2] Burnett KM, Reeves GH, Miller DJ, Clarke S, Vance-Borland K, Christiansen K. 2007. Distribution of salmon-habitat potential relative to landscape characteristics and implications for conservation. *Ecological Applications*, 17(1), 66-80.

[3] Casida JE, Baron RL, Eto M, Engel JL. 1963. Potentiation and neurotoxicity induced by certain organophosphates. *Biochemical pharmacology*, 12(1), 73-83.

[4] Gilliom, R.J. 2007. Pesticides in U.S. streams and groundwater. *Environmental Science and Technology*, 41:3408-3414.

[5] Harris KA, Dangerfield N, Woudneh M, Brown TG, Verrin S, Ross PS. 2008. Partitioning of current-use and legacy pesticides in salmon habitat in British Columbia, Canada. *Environmental Toxicology and Chemistry* 27:2253-2262.

[6] Jansen KL, Cole TB, Park SS, Furlong CE, Costa LG. 2009. Paraoxonase 1 (PON1) modulates the toxicity of mixed organophosphorus compounds. *Toxicology and Applied Pharmacology* 236:142-153.

[7] Jarrard HE, Delaney KR, Kennedy CJ. 2004. Impacts of carbamate pesticides on olfactory neurophysiology and cholinesterase activity in coho salmon (*Oncorhynchus tshawytscha*). *Aquatic Toxicology* 69:133-148.

[8] Laetz CA, Baldwin DH, Collier TK, Hebert V, Stark JD, Scholz NL. 2009. The synergistic toxicity of pesticide mixtures: implications for risk assessment and the conservation of endangered pacific salmon. *Environmental Health Perspectives*. 117: 348-353.

[9] Sargeant D, Newell E, Anderson P, Cook A. 2013. Surface Water Monitoring Program for Pesticides in Salmon-Bearing Streams, 2009-2011 Triennial Report. A Cooperative Study by the Washington State Departments of Ecology and Agriculture. Washington State Department of Ecology Publication 13-03-002 Department of Agriculture Publication AGR PUB 102-377 Washington State Department of Ecology, Olympia, WA

[10] Scholz NL, Truelove NK, French BL, Berjikian B, Quinn T, Casillas E, Collier TK. 2000. Diazinon disrupts antipredator and homing behaviors in chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries Aquatic Science* 57:1911-1918.

[11] Sorensen H, Cuedegreen N, Skovgaard IM, Streibig JC. 2007. An isobole-based statistical model and test for synergism/antagonism in binary mixture toxicity experiments. *Environmental Ecology Statistics* 14:383-397

[12] Tierney KB, Sampson JL, Ross PS, Sekela MA, Kennedy CJ. 2008. Salmon olfaction is impaired by an environmentally realistic pesticide mixture. *Environmental Science & Technology* 42:4996-5001

Individual contributors
Abigail Nickelson, WA Department of Agriculture
Cathy Laetz, NOAA Fisheries
Julann Stromberg, NOAA Fisheries/Ocean Associates

Contributing Agencies-Data on Sites and Exposure-Response
Northwest Fisheries Science Center, NOAA
USGS
WA Department of Agriculture
WA Department of Ecology
City of Bellingham
Nooksack Salmon Enhancement Association
King County DNR
King Conservation District
Skagit Conservation district
Yakima Tribal DNR
Samish Tribal DNR