



2-28-2022

## Lake Whatcom Monitoring Project 2020/2021 Report

Angela Strecker

*Western Washington University*

Michael Hilles

*Western Washington University*

Joan Pickens

*Western Washington University*

Robert Mitchell

*Western Washington University*

Robin Matthews

*Western Washington University*

*See next page for additional authors*

Follow this and additional works at: [https://cedar.wvu.edu/lakewhat\\_annualreps](https://cedar.wvu.edu/lakewhat_annualreps)



Part of the [Environmental Monitoring Commons](#)

---

### Recommended Citation

Strecker, Angela; Hilles, Michael; Pickens, Joan; Mitchell, Robert; Matthews, Robin; and Matthews, Geoffrey, "Lake Whatcom Monitoring Project 2020/2021 Report" (2022). *Lake Whatcom Annual Reports*. 30.

[https://cedar.wvu.edu/lakewhat\\_annualreps/30](https://cedar.wvu.edu/lakewhat_annualreps/30)

This Report is brought to you for free and open access by the Lake Whatcom at Western CEDAR. It has been accepted for inclusion in Lake Whatcom Annual Reports by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).

---

**Authors**

Angela Strecker, Michael Hilles, Joan Pickens, Robert Mitchell, Robin Matthews, and Geoffrey Matthews

# Lake Whatcom Monitoring Project 2020/2021 Report

Dr. Angela Strecker  
Michael Hilles  
Joan Pickens  
*Institute for Watershed Studies,  
College of the Environment*

Dr. Robert Mitchell  
*Geology Department,  
College of Science and Engineering*

Dr. Robin Matthews, Emeritus  
*Environmental Sciences Department  
College of the Environment*

Dr. Geoffrey Matthews, Emeritus  
*Computer Science Department  
College of Science and Engineering*

Western Washington University  
Bellingham, Washington 98225

February 28, 2022

---

Funding for this project was provided by the City of Bellingham, as part of their long-term commitment to environmental education and their concern for maintaining the water quality of Lake Whatcom. We thank Eion Atkins, Tessa Beaver, Ian Browning, Bella Colbert, Pilar Deniston, Ariana Dionisio, Emily Flarry, Ian Penner, Kathryn Queen, Emily Smoot, Russ Thompson, and Emma Yenney for assistance with the project.

(This page blank)

# Contents

<b>1</b>	<b>Background</b>	<b>1</b>
1.1	Objectives . . . . .	2
<b>2</b>	<b>Lake Whatcom Monitoring</b>	<b>2</b>
2.1	Site Descriptions . . . . .	2
2.2	Field Sampling and Analytical Methods . . . . .	3
2.3	Results and Discussion . . . . .	3
2.3.1	Water temperature . . . . .	4
2.3.2	Dissolved oxygen . . . . .	5
2.3.3	Conductivity and pH . . . . .	8
2.3.4	Alkalinity and turbidity . . . . .	9
2.3.5	Nitrogen and phosphorus . . . . .	9
2.3.6	Chlorophyll, plankton, and Secchi depth . . . . .	13
2.3.7	Coliform bacteria . . . . .	15
2.3.8	Total organic carbon and disinfection by-products . . . . .	16
<b>3</b>	<b>Tributary Monitoring</b>	<b>56</b>
3.1	Site Descriptions . . . . .	56
3.2	Field Sampling and Analytical Methods . . . . .	56
3.3	Results and Discussion . . . . .	57
<b>4</b>	<b>Storm Water Monitoring</b>	<b>77</b>
4.1	Hydrograph Monitoring . . . . .	77
4.2	Field Sampling and Analytical Methods . . . . .	77

4.3	Site Descriptions . . . . .	77
4.4	Field Sampling and Analytical Methods . . . . .	77
<b>5</b>	<b>References and Related Reports</b>	<b>86</b>
5.1	Cited References . . . . .	86
5.2	Related Reports . . . . .	90
<b>A</b>	<b>Site Descriptions</b>	<b>94</b>
A.1	Lake Whatcom Monitoring Sites . . . . .	94
A.2	Tributary Monitoring Sites . . . . .	94
A.3	Storm Water Monitoring Sites . . . . .	96
<b>B</b>	<b>Long-Term Water Quality Figures</b>	<b>100</b>
B.1	Monthly YSI Profiles . . . . .	101
B.2	Long-term YSI/Hydrolab Data (1988-present) . . . . .	167
B.3	Long-term Water Quality Data (1988-present) . . . . .	188
B.4	Lake Whatcom Tributary Data (2004-present) . . . . .	249
<b>C</b>	<b>Quality Control</b>	<b>290</b>
C.1	Performance Evaluation Reports . . . . .	290
C.2	Laboratory Duplicates, Spikes, and Check Standards . . . . .	290
C.3	Field Duplicates . . . . .	290
<b>D</b>	<b>Lake Whatcom Online Data</b>	<b>342</b>

## List of Figures

2.1	October 2020 temperature and dissolved oxygen compared to historic ranges . . . . .	28
2.2	November 2020 temperature and dissolved oxygen compared to historic ranges . . . . .	29
2.3	December 2020 temperature and dissolved oxygen compared to historic ranges . . . . .	30
2.4	February 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	31
2.5	April historic temperature and dissolved oxygen compared to historic ranges . . . . .	32
2.6	May 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	33
2.7	June 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	34
2.8	July 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	35
2.9	August 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	36
2.10	September 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	37
2.11	October 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	38
2.12	November 2021 temperature and dissolved oxygen compared to historic ranges . . . . .	39
2.13	December 2021 temperature and dissolved oxygen compared to historic ranges (preliminary data) . . . . .	40
2.14	Relationship between dissolved oxygen and time at Site 1, 12 m .	41
2.15	Relationship between dissolved oxygen and time at Site 1, 14 m .	42

2.16	Relationship between dissolved oxygen and time at Site 1, 16 m .	43
2.17	Relationship between dissolved oxygen and time at Site 1, 18 m .	44
2.18	Minimum summer, near-surface DIN concentrations . . . . .	45
2.19	Median spring vs. summer near-surface DIN concentrations . . . .	46
2.20	Spring/summer near-surface DIN concentration differences . . . .	47
2.21	Median summer, near-surface total phosphorus concentrations . .	48
2.22	Median summer near-surface chlorophyll concentrations . . . . .	49
2.23	Log <sub>10</sub> plots of median summer, near-surface algae counts . . . . .	50
2.24	Log <sub>10</sub> plots of median summer, near-surface Cyanobacteria counts	51
2.25	THMs in the Bellingham water distribution system . . . . .	52
2.26	Quarterly THMs in the Bellingham water distribution system . .	53
2.27	HAAs in the Bellingham water distribution system . . . . .	54
2.28	Quarterly HAAs in the Bellingham water distribution system . .	55
4.1	Austin Creek hydrograph for WY2021 . . . . .	82
4.2	Smith Creek hydrograph for WY2021 . . . . .	83
4.3	Aquarius rating curve for Austin Creek . . . . .	84
4.4	Aquarius rating curve for Smith Creek . . . . .	85
A1	Lake Whatcom lake sampling sites . . . . .	98
A2	Lake Whatcom tributary and storm water sampling sites . . . . .	99
B1	Water column profiles for Site 1, Oct. 7, 2020 . . . . .	102
B2	Water column profiles for Site 2, Oct. 7, 2020 . . . . .	103
B3	Water column profiles for the Intake, Oct. 7, 2020 . . . . .	104
B4	Water column profiles for Site 3, Oct. 5, 2020 . . . . .	105
B5	Water column profiles for Site 4, Oct. 5, 2020 . . . . .	106
B6	Water column profiles for Site 1, Nov. 9, 2020 . . . . .	107



B7	Water column profiles for Site 2, Nov. 9, 2020 . . . . .	108
B8	Water column profiles for the Intake, Nov. 9, 2020 . . . . .	109
B9	Water column profiles for Site 3, Nov. 2, 2020 . . . . .	110
B10	Water column profiles for Site 4, Nov. 2, 2020 . . . . .	111
B11	Water column profiles for Site 1, Dec. 7, 2020 . . . . .	112
B12	Water column profiles for Site 2, Dec. 7, 2020 . . . . .	113
B13	Water column profiles for the Intake, Dec. 7, 2020 . . . . .	114
B14	Water column profiles for Site 3, Dec. 2, 2020 . . . . .	115
B15	Water column profiles for Site 4, Dec. 2, 2020 . . . . .	116
B16	Water column profiles for Site 1, Feb. 27, 2021 . . . . .	117
B17	Water column profiles for Site 2, Feb. 24, 2021 . . . . .	118
B18	Water column profiles for the Intake, Feb. 27, 2021 . . . . .	119
B19	Water column profiles for Site 3, Feb. 17, 2021 . . . . .	120
B20	Water column profiles for Site 4, Feb. 17, 2021 . . . . .	121
B21	Water column profiles for Site 1, April 8, 2021 . . . . .	122
B22	Water column profiles for Site 2, April 8, 2021 . . . . .	123
B23	Water column profiles for the Intake, April 8, 2021 . . . . .	124
B24	Water column profiles for Site 3, April 6, 2021 . . . . .	125
B25	Water column profiles for Site 4, April 6, 2021 . . . . .	126
B26	Water column profiles for Site 1, May 10, 2021 . . . . .	127
B27	Water column profiles for Site 2, May 6, 2021 . . . . .	128
B28	Water column profiles for the Intake, May 10, 2021 . . . . .	129
B29	Water column profiles for Site 3, May 4, 2021 . . . . .	130
B30	Water column profiles for Site 4, May 4, 2021 . . . . .	131
B31	Water column profiles for Site 1, Jun. 3, 2021 . . . . .	132

B32	Water column profiles for Site 2, Jun. 3, 2021 . . . . .	133
B33	Water column profiles for the Intake, Jun. 3, 2021 . . . . .	134
B34	Water column profiles for Site 3, Jun. 1, 2021 . . . . .	135
B35	Water column profiles for Site 4, Jun. 1, 2021 . . . . .	136
B36	Water column profiles for Site 1, Jul. 6, 2021 . . . . .	137
B37	Water column profiles for Site 2, Jul. 8, 2021 . . . . .	138
B38	Water column profiles for the Intake, Jul. 8, 2021 . . . . .	139
B39	Water column profiles for Site 3, Jul. 8, 2021 . . . . .	140
B40	Water column profiles for Site 4, Jul. 6, 2021 . . . . .	141
B41	Water column profiles for Site 1, Aug. 9, 2021 . . . . .	142
B42	Water column profiles for Site 2, Aug. 9, 2021 . . . . .	143
B43	Water column profiles for the Intake, Aug. 9, 2021 . . . . .	144
B44	Water column profiles for Site 3, Aug. 11, 2021 . . . . .	145
B45	Water column profiles for Site 4, Aug. 11, 2021 . . . . .	146
B46	Water column profiles for Site 1, Sept. 9, 2021 . . . . .	147
B47	Water column profiles for Site 2, Sept. 9, 2021 . . . . .	148
B48	Water column profiles for the Intake, Sept. 9, 2021 . . . . .	149
B49	Water column profiles for Site 3, Sept. 7, 2021 . . . . .	150
B50	Water column profiles for Site 4, Sept. 7, 2021 . . . . .	151
B51	Water column profiles for Site 1, Oct. 7, 2021 . . . . .	152
B52	Water column profiles for Site 2, Oct. 7, 2021 . . . . .	153
B53	Water column profiles for the Intake, Oct. 7, 2021 . . . . .	154
B54	Water column profiles for Site 3, Oct. 19, 2021 . . . . .	155
B55	Water column profiles for Site 4, Oct. 12, 2021 . . . . .	156
B56	Water column profiles for Site 1, Nov. 9, 2021 . . . . .	157

B57	Water column profiles for Site 2, Nov. 9, 2021 . . . . .	158
B58	Water column profiles for the Intake, Nov. 9, 2021 . . . . .	159
B59	Water column profiles for Site 3, Nov. 2, 2021 . . . . .	160
B60	Water column profiles for Site 4, Nov. 2, 2021 . . . . .	161
B61	Water column profiles for Site 1, Dec. 7, 2021 . . . . .	162
B62	Water column profiles for Site 2, Dec. 7, 2021 . . . . .	163
B63	Water column profiles for the Intake, Dec. 7, 2021 . . . . .	164
B64	Water column profiles for Site 3, Dec. 2, 2021 . . . . .	165
B65	Water column profiles for Site 4, Dec. 2, 2021 . . . . .	166
B66	Historic temperature data for Site 1 . . . . .	168
B67	Historic temperature data for Site 2 . . . . .	169
B68	Historic temperature data for the Intake . . . . .	170
B69	Historic temperature data for Site 3 . . . . .	171
B70	Historic temperature data for Site 4 . . . . .	172
B71	Historic dissolved oxygen data for Site 1 . . . . .	173
B72	Historic dissolved oxygen data for Site 2 . . . . .	174
B73	Historic dissolved oxygen data for the Intake . . . . .	175
B74	Historic dissolved oxygen data for Site 3 . . . . .	176
B75	Historic dissolved oxygen data for Site 4 . . . . .	177
B76	Historic pH data for Site 1 . . . . .	178
B77	Historic pH data for Site 2 . . . . .	179
B78	Historic pH data for the Intake . . . . .	180
B79	Historic pH data for Site 3 . . . . .	181
B80	Historic pH data for Site 4 . . . . .	182
B81	Historic conductivity data for Site 1 . . . . .	183

B82	Historic conductivity data for Site 2 . . . . .	184
B83	Historic conductivity data for the Intake . . . . .	185
B84	Historic conductivity data for Site 3 . . . . .	186
B85	Historic conductivity data for Site 4 . . . . .	187
B86	Alkalinity data for Site 1 . . . . .	189
B87	Alkalinity data for Site 2 . . . . .	190
B88	Alkalinity data for the Intake site . . . . .	191
B89	Alkalinity data for Site 3 . . . . .	192
B90	Alkalinity data for Site 4 . . . . .	193
B91	Turbidity data for Site 1 . . . . .	194
B92	Turbidity data for Site 2 . . . . .	195
B93	Turbidity data for the Intake site . . . . .	196
B94	Turbidity data for Site 3 . . . . .	197
B95	Turbidity data for Site 4 . . . . .	198
B96	Ammonium data for Site 1 . . . . .	199
B97	Ammonium data for Site 2 . . . . .	200
B98	Ammonium data for the Intake site . . . . .	201
B99	Ammonium data for Site 3 . . . . .	202
B100	Ammonium data for Site 4 . . . . .	203
B101	Nitrate/nitrite data for Site 1 . . . . .	204
B102	Nitrate/nitrite data for Site 2 . . . . .	205
B103	Nitrate/nitrite data for the Intake site . . . . .	206
B104	Nitrate/nitrite data for Site 3 . . . . .	207
B105	Nitrate/nitrite data for Site 4 . . . . .	208
B106	Total nitrogen data for Site 1 . . . . .	209

B107	Total nitrogen data for Site 2 . . . . .	210
B108	Total nitrogen data for the Intake site . . . . .	211
B109	Total nitrogen data for Site 3 . . . . .	212
B110	Total nitrogen data for Site 4 . . . . .	213
B111	Soluble phosphate data for Site 1 . . . . .	214
B112	Soluble phosphate data for Site 2 . . . . .	215
B113	Soluble phosphate data for the Intake site . . . . .	216
B114	Soluble phosphate data for Site 3 . . . . .	217
B115	Soluble phosphate data for Site 4 . . . . .	218
B116	Total phosphorus data for Site 1 . . . . .	219
B117	Total phosphorus data for Site 2 . . . . .	220
B118	Total phosphorus data for the Intake site . . . . .	221
B119	Total phosphorus data for Site 3 . . . . .	222
B120	Total phosphorus data for Site 4 . . . . .	223
B121	Chlorophyll data for Site 1 . . . . .	224
B122	Chlorophyll data for Site 2 . . . . .	225
B123	Chlorophyll data for the Intake site . . . . .	226
B124	Chlorophyll data for Site 3 . . . . .	227
B125	Chlorophyll data for Site 4 . . . . .	228
B126	Secchi depths for Site 1 . . . . .	229
B127	Secchi depths for Site 2 . . . . .	230
B128	Secchi depths for the Intake site . . . . .	231
B129	Secchi depths for Site 3 . . . . .	232
B130	Secchi depths for Site 4 . . . . .	233
B131	Fecal coliform data for Site 1 . . . . .	234

B132	Fecal coliform data for Site 2 . . . . .	235
B133	Fecal coliform data for the Intake site . . . . .	236
B134	Fecal coliform data for Site 3 . . . . .	237
B135	Fecal coliform data for Site 4 . . . . .	238
B136	Plankton data for Site 1 . . . . .	239
B137	Plankton data for Site 2 . . . . .	240
B138	Plankton data for the Intake Site . . . . .	241
B139	Plankton data for Site 3 . . . . .	242
B140	Plankton data for Site 4 . . . . .	243
B141	Plankton data for Site 1 (omit Chrysophyta) . . . . .	244
B142	Plankton data for Site 2 (omit Chrysophyta) . . . . .	245
B143	Plankton data for the Intake Site (omit Chrysophyta) . . . . .	246
B144	Plankton data for Site 3 (omit Chrysophyta) . . . . .	247
B145	Plankton data for Site 4 (omit Chrysophyta) . . . . .	248
B146	Temperature data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	251
B147	Temperature data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	252
B148	Temperature data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	253
B149	Dissolved oxygen data for Anderson, Austin, Smith, and What- com Creeks . . . . .	254
B150	Dissolved oxygen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	255
B151	Dissolved oxygen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	256
B152	pH data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	257

B153	pH data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks	258
B154	pH data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain	259
B155	Conductivity data for Anderson, Austin, Smith, and Whatcom Creeks	260
B156	Conductivity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks	261
B157	Conductivity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain	262
B158	Alkalinity data for Anderson, Austin, Smith, and Whatcom Creeks	263
B159	Alkalinity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks	264
B160	Alkalinity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain	265
B161	Total suspended solids data for Anderson, Austin, Smith, and Whatcom Creeks	266
B162	Total suspended solids data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks	267
B163	Total suspended solids data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain	268
B164	Turbidity data for Anderson, Austin, Smith, and Whatcom Creeks	269
B165	Turbidity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks	270
B166	Turbidity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain	271
B167	Ammonium data for Anderson, Austin, Smith, and Whatcom Creeks	272
B168	Ammonium data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks	273

B169	Ammonium data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	274
B170	Nitrate/nitrite data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	275
B171	Nitrate/nitrite data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	276
B172	Nitrate/nitrite data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	277
B173	Total nitrogen data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	278
B174	Total nitrogen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	279
B175	Total nitrogen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	280
B176	Soluble phosphate data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	281
B177	Soluble phosphate data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	282
B178	Soluble phosphate data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	283
B179	Total phosphorus data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	284
B180	Total phosphorus data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	285
B181	Total phosphorus data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	286
B182	Fecal coliform data for Anderson, Austin, Smith, and Whatcom Creeks . . . . .	287
B183	Fecal coliform data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks . . . . .	288



B184	Fecal coliform data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain . . . . .	289
C1	Alkalinity laboratory duplicates . . . . .	293
C2	Alkalinity high-range check standards . . . . .	294
C3	Alkalinity low-range check standards . . . . .	295
C4	Chlorophyll laboratory duplicates . . . . .	296
C5	Conductivity laboratory duplicates . . . . .	297
C6	Dissolved oxygen laboratory duplicates . . . . .	298
C7	Nitrogen (ammonium) laboratory duplicates . . . . .	299
C8	Nitrogen (ammonium) spike recoveries . . . . .	300
C9	Nitrogen (ammonium) high-range check standards . . . . .	301
C10	Nitrogen (ammonium) low-range check standards . . . . .	302
C11	Nitrogen (nitrate/nitrite) laboratory duplicates . . . . .	303
C12	Nitrogen (nitrate/nitrite) spike recoveries . . . . .	304
C13	Nitrogen (nitrate/nitrite) high-range check standards . . . . .	305
C14	Nitrogen (nitrate/nitrite) low-range check standards . . . . .	306
C15	Nitrogen (total) laboratory duplicates . . . . .	307
C16	Nitrogen (total) spike recoveries . . . . .	308
C17	Nitrogen (total) high-range check standards . . . . .	309
C18	Nitrogen (total) low-range check standards . . . . .	310
C19	Laboratory pH duplicates . . . . .	311
C20	Phosphorus (soluble reactive phosphate) laboratory duplicates . . . . .	312
C21	Phosphorus (soluble reactive phosphate) spike recoveries . . . . .	313
C22	Phosphorus (soluble reactive phosphate) high-range check standards . . . . .	314

C23	Phosphate (soluble reactive phosphate) low-range check standards	315
C24	Phosphorus (total) laboratory duplicates . . . . .	316
C25	Phosphorus (total) spike recoveries . . . . .	317
C26	Phosphorus (total) high-range check standards . . . . .	318
C27	Phosphorus (total) low-range check standards . . . . .	319
C28	Total suspended solids laboratory duplicates . . . . .	320
C29	Total suspended solids check standards . . . . .	321
C30	Turbidity laboratory duplicates . . . . .	322
C31	Alkalinity field duplicates (lake samples) . . . . .	323
C32	Alkalinity field duplicates (tributary samples) . . . . .	324
C33	Chlorophyll field duplicates (lake samples) . . . . .	325
C34	Conductivity field duplicates (lake samples) . . . . .	326
C35	Dissolved oxygen field duplicates (lake samples) . . . . .	327
C36	Dissolved oxygen field duplicates (tributary samples) . . . . .	328
C37	Nitrogen (ammonium) field duplicates (lake samples) . . . . .	329
C38	Nitrogen (ammonium) field duplicates (tributary samples) . . . . .	330
C39	Nitrogen (nitrate/nitrite) field duplicates (lake samples) . . . . .	331
C40	Nitrogen (nitrate/nitrite) field duplicates (tributary samples) . . . . .	332
C41	Nitrogen (total) field duplicates (lake samples) . . . . .	333
C42	Nitrogen (total) field duplicates (tributary samples) . . . . .	334
C43	Field duplicates for pH (lake samples) . . . . .	335
C44	Phosphorus (soluble reactive phosphate) field duplicates (tributary samples) . . . . .	336
C45	Phosphorus (total) field duplicates (lake samples) . . . . .	337
C46	Phosphorus (total) field duplicates (tributary samples) . . . . .	338

C47	Total suspended solids field duplicates (tributary samples) . . . .	339
C48	Turbidity field duplicates (lake samples) . . . . .	340
C49	Turbidity field duplicates (tributary samples) . . . . .	341

## List of Tables

2.1	Analytical methods and parameter abbreviations . . . . .	18
2.2	Lake Whatcom proposed lake monitoring schedule . . . . .	19
2.3	Summary of missing lake data . . . . .	20
2.4	Summary of Site 1 water quality data, Oct. 2020 – Sept. 2021. . .	21
2.5	Summary of Intake water quality data, Oct. 2020– Sept. 2021 . . .	22
2.6	Summary of Site 2 water quality data, Oct. 2020– Sept. 2021 . . .	23
2.7	Summary of Site 3 water quality data, Oct. 2020– Sept. 2021 . . .	24
2.8	Summary of Site 4 water quality data, Oct. 2020– Sept. 2021 . . .	25
2.9	October hypolimnetic hydrogen sulfide concentrations . . . . .	26
2.10	Lake Whatcom 2020/2021 total organic carbon data . . . . .	27
3.1	Lake Whatcom proposed tributary monitoring schedule . . . . .	61
3.2	Summary of missing tributary data . . . . .	62
3.3	Comparison of 2020–2021 water quality in Lake Whatcom tribu- taries . . . . .	63
3.4	Anderson Creek water quality data, October 2020-September 2021	64
3.5	Austin Creek water quality data, October 2020-September 2021 .	65
3.6	Blue Canyon Creek water quality data, October 2020-September 2021 . . . . .	66
3.7	Brannian Creek water quality data, October 2020-September 2021	67
3.8	Carpenter Creek water quality data, October 2020-September 2021	68
3.9	Euclid Creek water quality data, October 2020-September 2021 .	69
3.10	Millwheel Creek water quality data, October 2020-September 2021	70
3.11	Olsen Creek water quality data, October 2020-September 2021 . .	71
3.12	Park Place outlet water quality data, October 2020-September 2021	72

3.13	Silver Beach Creek water quality data, October 2020-September 2021 . . . . .	73
3.14	Smith Creek water quality data, October 2020-September 2021 . .	74
3.15	Whatcom Creek water quality data, October 2020-September 2021	75
3.16	Lake Whatcom 2021 tributary total organic carbon data . . . . .	76
4.1	Austin Creek rating curves for WY2021 . . . . .	79
4.2	Smith Creek rating curves for WY2021 . . . . .	80
4.3	Summary of storm event sampling in Austin, Euclid, and Silver Beach Creeks . . . . .	81
A1	Approximate GPS coordinates for Lake Whatcom sampling sites.	97
B1	List of outliers omitted from Figures B146–B184 . . . . .	250
C1	Single-blind quality control results . . . . .	291
C2	Quality control values outside plotting range . . . . .	292

# Executive Summary

## Background for the Lake Whatcom Annual Reports

- This report describes the results from the 2020/2021 Lake Whatcom monitoring program conducted by the Institute for Watershed Studies at Western Washington University ([www.wwu.edu/iws](http://www.wwu.edu/iws)).
- The major objectives in 2020/2021 were to continue long-term baseline water quality monitoring in Lake Whatcom and its major tributaries; collect storm runoff water quality data from representative streams in the watershed; and continue collection of hydrologic data from Austin and Smith Creeks.
- Each section in this report contains a brief discussion of the water quality parameters that are measured as part of the monitoring effort. For additional help with understanding the relationship between water quality data and lake, stream, or watershed ecology, we recommend the online resource “Water on the Web” (WOW, 2004; [www.waterontheweb.org](http://www.waterontheweb.org)).
- The online pdf copy of this report contains **red hyperlinks** that will open online citations, and **blue hyperlinks** that will jump to referenced tables and figures or to the section that contains additional information about a specific topic. These hyperlinks are active if the report is opened using Adobe Reader, which can be downloaded free from [www.adobe.com/products/reader.html](http://www.adobe.com/products/reader.html).
- This report is part of an on-going series of annual reports and special project reports that provide a complete documentation of the monitoring program over time. A summary of the Institute for Watershed Studies Lake Whatcom reports, including special project reports, is included in Section 5.2, beginning on page 90, and many of the reports are available online through Western CEDAR, the WWU repository for open access scholarship, under the Institute for Watershed Studies Lake Whatcom collection ([http://cedar.wwu.edu/iws\\_lakewhatcom](http://cedar.wwu.edu/iws_lakewhatcom)).

## Summary of 2020/2021 Monitoring Project

- During the summer the lake's water column was [thermally stratified](#) into a warm surface layer (the epilimnion) and a cool bottom layer (the hypolimnion). Most of the 2020/2021 temperature profiles fell within historical ranges, with stable stratification present at [Sites 1–4](#) by mid-May (Section [2.3.1](#), page [4](#)).<sup>1</sup>
- The [hypolimnetic oxygen](#) concentrations have declined over time at Site 1 (Section [2.3.2](#), page [5](#)), causing the lake to be listed by the Department of Ecology on the 1998 303d list of impaired waterbodies in the state of Washington. The hypolimnetic oxygen loss was apparent at Site 1 after the lake became fully stratified in June, and by August the oxygen concentrations were <2 mg/L from 11 meters to the bottom.
- [Nitrate depletion](#) was evident at all sites in the photosynthetic zone during the summer due to algal uptake of this essential nutrient (Section [2.3.5](#), page [10](#)). Unlike the other [indicators of phytoplankton productivity](#), the dissolved inorganic nitrogen (DIN = nitrate + nitrite + ammonium) trend has not stabilized in recent years. A month-by-month analysis of near-surface DIN showed that water column concentrations have declined in general, not just in the summer. Nitrate depletion also occurred in the hypolimnion at Sites 1 and 2 due to nitrate reduction by bacteria. Anaerobic conditions in the hypolimnion at Sites 1 and 2 resulted in elevated concentrations of ammonium by the end of the summer.
- The summer near-surface [total phosphorus](#) concentrations continued to follow erratic patterns, with no significant correlations with year (Section [2.3.5](#), page [12](#)), reflecting the complicated nature of phosphorus movement in the water column.
- The summer near-surface [chlorophyll concentrations](#) have increased significantly over time at all sites (Section [2.3.6](#), page [13](#)). Despite being quite variable, the concentrations appear to have stabilized since 2004, ranging from 3.7–6.7  $\mu\text{g/L}$  at Site 1 and 2.6–4.6  $\mu\text{g/L}$  at Sites 2–4.

---

<sup>1</sup>These links direct the reader to sections with additional information on the summary topic.

- All of the mid-basin [coliform counts](#) were less than 10 cfu/100 mL (Section [2.3.7](#), page [15](#)). The coliform counts at the Bloedel-Donovan recreational area (collected offshore from the swimming area) were slightly higher than mid-basin counts, but passed the freshwater *Primary Contact Recreational* bacteria standard for Washington in place prior to 2021 (see Section [2.3.7](#), page [15](#) for discussion of changes in standards).
- The concentrations of [trihalomethanes and haloacetic acids](#) (THMs and HAAs) in Bellingham’s treated drinking water have been increasing over time, but the concentrations of both types of disinfection by-products remained below the maximum contaminant levels of 0.080 mg/L and 0.060 mg/L, respectively (Section [2.3.8](#), page [16](#)), and appear to be declining in recent years.
- [Monthly tributary samples](#) were collected at 12 locations in the Lake Whatcom watershed (Section [3](#), page [56](#)). Most of the tributaries had low concentrations of total suspended solids, low alkalinities and conductivities, and low levels of nutrients (phosphorus and nitrogen). The residential streams had higher concentrations of total suspended solids, higher alkalinities and conductivities, higher coliform counts, and higher nutrient concentrations.
- [Hydrograph data](#) were collected at Austin and Smith Creeks using rating curves developed with Aquarius software to calculate discharge (Section [4.1](#), page [77](#)).
- [Storm runoff samples](#) were collected in Austin Creek (two storm events), Euclid Creek (three storm events), and Silver Beach Creek (four storm events) using time-paced automated samplers (Section [4](#), page [77](#)). The water quality data are sent to the City of Bellingham for use in watershed modeling.



# 1 Background

This report is part of an on-going series of annual reports and special project reports that document the Lake Whatcom monitoring program over time. Many of the reports are available online through Western CEDAR, the WWU repository for open access scholarship, under the Institute for Watershed Studies Lake Whatcom collection ([http://cedar.wvu.edu/iws\\_lakewhatcom](http://cedar.wvu.edu/iws_lakewhatcom)). Reports that are not available on CEDAR may be available in the Institute for Watershed Studies (IWS) library or through the City of Bellingham Public Works Department. A summary of the Lake Whatcom annual and special project reports is included in Section 5.2, beginning on page 90.

Each section in this report contains a brief discussion of the water quality parameters that are measured as part of the monitoring effort. For additional help with understanding the relationship between water quality data and lake, stream, or watershed ecology, we recommend the online resource “Water on the Web” ([www.waterontheweb.org](http://www.waterontheweb.org); WOW, 2004).

Lake Whatcom is the primary drinking water source for the City of Bellingham and parts of Whatcom County, including Sudden Valley. It also serves as a primary or supplemental water source to various water systems to the City of Bellingham.

The lake and its watershed provide recreational opportunities, as well as important habitats for fish and wildlife. The lake is used as a storage reservoir to buffer peak storm water flows in Whatcom Creek. Because of its aesthetic appeal, the watershed is highly valued for residential development. Historically, most of the nonresidential portion of the watershed was zoned for forestry and was managed by state or private timber companies.

In January 22, 2014, approximately 7,800 acres of forest lands formerly managed by the Department of Natural Resources was reconveyed to Whatcom County to be managed as low impact park lands. The Lake Whatcom reconveyance planning process is summarized online at [www.whatcomcounty.us/625/Lake-Whatcom-Reconveyance](http://www.whatcomcounty.us/625/Lake-Whatcom-Reconveyance).

## 1.1 Objectives

The City of Bellingham and Western Washington University have collaborated on water quality studies in Lake Whatcom since the early 1960s. Beginning in 1988, a monitoring program was initiated by the City and WWU that was designed to provide long-term lake data for temperature, pH, dissolved oxygen, conductivity, turbidity, nutrients (nitrogen and phosphorus), and other representative water quality measurements. The major goal of the long-term monitoring effort is to provide a record of Lake Whatcom's water quality over time.

The major objectives of the 2020/2021 Lake Whatcom monitoring program were to continue long-term baseline water quality monitoring in Lake Whatcom and its major tributaries; collect storm runoff water quality data from representative streams in the watershed; and continue collection of hydrologic data from Austin and Smith Creeks.

Detailed site descriptions can be found in Appendix A. The historic lake data are plotted in Appendix B. The current quality control results are in Appendix C. The monitoring data are available online at [www.wwu.edu/iws](http://www.wwu.edu/iws) as described in Appendix D (page 342). Table 2.1 (page 18) lists abbreviations and units used to describe water quality analyses; Tables 2.2 & 3.1 (pages 19 & 61) list the locations, depths, and frequency for lake and tributary sampling.

## 2 Lake Whatcom Monitoring

### 2.1 Site Descriptions

Water quality samples were collected at five long-term monitoring sites in Lake Whatcom (Figure A1, page 98 in Appendix A.1). Sites 1–2 are located at the deepest points in their respective basins. The Intake site is located adjacent to the underwater intake point where the City of Bellingham withdraws lake water from basin 2. Site 3 is located at the deepest point in the northern sub-basin of basin 3 and Site 4 is located at the deepest point in the southern sub-basin of basin 3. Water samples were also collected at the City of Bellingham Lake Whatcom Gatehouse, which is located onshore and west of the Intake site.

## 2.2 Field Sampling and Analytical Methods

The lake was sampled on October 5 & 7, November 2 & 9, and December 2 & 7, 2020; and February 17, 24 & 27, April 6 & 8, May 4, 6 & 10, June 1 & 3, July 6 & 8, August 9 & 11, and September 7 & 9, 2021. Each sampling event is a multi-day task; all samples were collected during daylight hours, typically between 10:00 am and 3:00 pm. The analytical and sampling procedures are summarized in Tables 2.1 & 2.2 (pages 18 & 19). Table 2.3 (page 20) summarizes missing data from the 2020/2021 sampling season.

A YSI EXO1 multiparameter field meter was used to measure temperature, pH, dissolved oxygen, and conductivity in the field. Raw water and bacteriological samples were collected using a VanDorn sampler; plankton samples were collected using a 30-L Schindler trap equipped with a 20  $\mu\text{m}$  mesh plankton net. The water and bacteriological samples were stored on ice and in the dark until they reached the laboratory. Plankton samples were placed in a cooler and returned to the laboratory unpreserved. The plankton sample volumes were measured in the laboratory and the samples were preserved with Lugol's solution. Total organic carbon analyses were done by AmTest<sup>2</sup> and by IWS. The bacteria samples were analyzed by the City of Bellingham.

## 2.3 Results and Discussion

The lake monitoring data include monthly field measurements (conductivity, dissolved oxygen, pH, Secchi depth, and water temperature); laboratory analyses for ambient water quality parameters (ammonium,<sup>3</sup> nitrate/nitrite,<sup>4</sup> total nitrogen, soluble phosphate, total phosphorus, alkalinity, turbidity, chlorophyll); plankton and bacteria counts; and total organic carbon measurements.

The 2020/2021 temperature, dissolved oxygen, pH, and conductivity profiles are shown in Figures B1–B50 (Appendix B, pages 102–151). Tables 2.4–2.8 (pages 21–25) summarize the current field measurements, ambient water quality, and

<sup>2</sup>AmTest, 13600 Northeast 126th Place, Suite C, Kirkland, WA, 98034–8720.

<sup>3</sup>Nearly all ammonia ( $\text{NH}_4^+$ ) is ionized to ammonium ( $\text{NH}_3$ ) in surface water. Earlier IWS reports used “ammonia” and “ammonium” interchangeably; we now use “ammonium” to indicate that the data represent the concentration of ionized ammonia.

<sup>4</sup>Nitrate and nitrite were analyzed together because nitrite concentrations are very low in surface water. For simplicity, nitrate/nitrite will be referred to as “nitrate” in this document.

coliform data, and all of the current data are plotted in comparison with historic data in Figures B66–B145 (Appendix B, pages 168–248). These figures are scaled to plot the full range of Lake Whatcom water quality data including minimum, maximum, and outlier values, and do not provide the best illustration of trends that occur in the lake. Separate tables and figures are provided to show trends and illustrate specific patterns in the data. The raw data are available online at [www.wvu.edu/iws](http://www.wvu.edu/iws) as described in Appendix D (page 342).

### 2.3.1 Water temperature

The 2020/2021 monthly temperature profiles for Sites 1–4 were plotted as overlay points on shaded polygons that summarize the 1988–2021 historic temperature ranges (Figures 2.1–2.13, pages 28–40). The monthly YSI profiles for temperature, dissolved oxygen, pH, and conductivity at Sites 1–4 and the Intake were included in Appendix B (Figures B1–B50, pages 102–151).

The summer temperature profiles (e.g., Figure 2.7, page 34) show how the lake stratifies into a warm surface layer (*epilimnion*), and cool bottom layer (*hypolimnion*). The transition zone between the epilimnion and hypolimnion (*metalimnion*) is a region of rapidly changing water temperature. When stratified, the temperature profiles show distinct differences between the surface and bottom of the water column. Stratification develops gradually, and once stable, persists until fall or winter, depending on location in the lake. Seasonal weather differences alter the timing of lake stratification; if the spring is cool, cloudy, and windy, the lake may stratify later than when it has been hot and sunny.

In Lake Whatcom, all sites except the Intake<sup>5</sup> are usually stratified by late spring or early summer. Stratification may begin as early as April, but is often not stable until May or June. The stability of stratification is determined in part by the temperature differences in the water column, but also by water circulation and local weather patterns. Once the water column temperature differs by at least 5°C ( $\Delta T \geq 5^\circ\text{C}$ ), it is unlikely that the lake will destratify.<sup>6</sup>

---

<sup>5</sup>The Intake is too shallow to develop stable stratification (see Appendix B, Figures B1–B46).

<sup>6</sup>The  $\Delta T$  is the difference between the epilimnion and hypolimnion temperatures.

As the weather becomes colder and days shorten, the lake cools and the surface and bottom water temperatures become more similar. Eventually the water column will start to mix from the surface to the bottom and the lake will destratify. Basins 1 and 2 (Sites 1–2) usually destratify by the end of October or early November, but basin 3 (Sites 3–4) is usually still stratified in November (Figure 2.2, page 29). Complete destratification of basin 3 occurs in December or early January, so by February the temperatures are uniform throughout the water column at all sites (Figure 2.4, page 31).

Although destratification is relatively abrupt, the process of mixing the entire water column is not instantaneous. When the lake begins to destratify, water temperatures may be uniform from the surface to the bottom, but the rate of water circulation may not be sufficient to replenish hypolimnetic oxygen concentrations. This phenomenon, where temperature is uniform, but dissolved compounds (e.g., dissolved oxygen) remain partially stratified, is common in the early stages of destratification, when the basin is starting to mix (see November 2013 temperature and oxygen profiles from Site 2; Figure B.7 in Matthews, et al., 2015).

The lake was still stratified at all sites in October 2020 (Figure 2.1, page 28). Sites 1–2 were completely destratified by November (Figure 2.2, page 29). Sites 3–4 were close to destratification in December 2020 ( $\Delta T \leq 3^\circ \text{C}$ ); the entire lake was destratified in February 2021 (Figures 2.3–2.4, pages 30–31).

The water column was starting to stratify in April 2021 ( $\Delta T = 0.7\text{--}1.7^\circ \text{C}$ ). Sites 1–4 developed stable stratification ( $\Delta T > 5^\circ \text{C}$ ) by early May (Figure 2.6, pages 33). All sites remained stratified through September, with temperatures falling within typical historic ranges (Figures 2.7, 2.9, and 2.10, pages 34–37), with the exception of July 2021 (Figure 2.8, page 35). Sampling in July followed a period of exceptionally high air temperatures in the Pacific Northwest. Surface water temperature at Site 1 ( $24.19^\circ \text{C}$ ) was the highest value that has been recorded anywhere in the lake by the Institute for Watershed Studies since data collection began in 1988.

### 2.3.2 Dissolved oxygen

The 2020/2021 monthly oxygen profiles for Sites 1–4 were plotted as overlay points on shaded polygons that summarize the 1988–2021 historic temperature

ranges (Figures 2.1–2.13, pages 28–40).<sup>7</sup> The monthly YSI profiles for temperature, dissolved oxygen, pH, and conductivity at Sites 1–4 and the Intake were included in Appendix B (Figures B1–B50, pages 102–151).

As in past years, Sites 1–2 developed severe hypolimnetic oxygen deficits during the summer (Figures 2.8–2.10, pages 35–37). Hypolimnetic oxygen depletion only becomes apparent after stratification, when the lower waters of the basin are isolated from the lake’s surface and biological respiration consumes the oxygen dissolved in the water. Biological respiration usually increases when there is an abundant supply of organic matter (e.g., decomposing algae). In basin 3, which has a very large, well-oxygenated hypolimnion, respiration has relatively little influence on hypolimnetic oxygen concentrations. In contrast, there is rapid depletion of the hypolimnetic oxygen concentrations at Sites 1–2. These two sites are in shallow basins that have small hypolimnions compared to their photic zones<sup>8</sup> so decomposition of algae and other organic matter causes a significant drop in hypolimnetic oxygen over the summer. This oxygen depletion may be apparent in May if the lake stratifies early in the spring, but is more commonly observed beginning in June (Figure 2.7, page 34).

Low oxygen conditions are associated with a number of unappealing water quality problems in lakes, including loss of aquatic habitat; release of phosphorus from the sediments; increased rates of algal production due to release of phosphorus; unpleasant odors during lake destratification; fish kills, particularly during lake destratification; release of metals and organics from the sediments; increased mercury methylation; increased drinking water treatment costs; increased taste and odor problems in drinking water; and increased risks associated with disinfection by-products created during the drinking water treatment process.

The levels of hypolimnetic oxygen have declined over time at Site 1, causing the lake to be listed by the Department of Ecology as an “impaired” waterbody (Pelletier, 1998).<sup>9</sup> The increasing rate of oxygen loss is most apparent during July and August, after the lake develops stable stratification but before oxygen levels

<sup>7</sup>October–December 2021 are not part of the 2020/2021 sampling period, but the temperature/oxygen profiles were included to provide information on the timing of destratification.

<sup>8</sup>The photic zone is the region with enough light to support algal photosynthesis, which extends to about 10 m below the surface in Lake Whatcom. Assuming a photic zone of 0–10 m, the photic zones for basins 1, 2, and 3 would be 75%, 70%, and 17% of the basin’s volume, respectively (Mitchell, et al., 2010).

<sup>9</sup>[www.ecy.wa.gov/programs/wq/303d](http://www.ecy.wa.gov/programs/wq/303d).

drop near zero. To illustrate this trend we fitted the July and August data using an exponential function (see discussion by Matthews, et al., 2004). As indicated in Figures 2.14–2.17 (pages 41–44), there were significant negative correlations<sup>10</sup> between dissolved oxygen and time for all hypolimnetic samples collected during July and August. Although Site 1 didn't develop stable stratification until May, the July hypolimnetic oxygen concentrations were already <4 mg/L and by August the oxygen concentrations were <1 mg/L (Figures 2.14–2.17).

A region of supersaturated oxygen was evident in the metalimnion at Site 1 in May, June, and July (Figures 2.6–2.8, pages 33–35; Figures B26–B36, pages 127–137). This was caused by the accumulation of phytoplankton along the density gradient between the epilimnion and hypolimnion where light and nutrients are sufficient to support very high levels of photosynthesis. Chlorophyll concentrations within the metalimnetic oxygen peak may be 4–5 times higher than those measured near the surface of the lake (Matthews and DeLuna, 2008). Metalimnetic oxygen peaks are common at Site 1 during the summer, and may occur at Sites 2–4, but will usually be at different depths because the metalimnions are at different depths. When present, the metalimnions form at approximately 5–10 m at Site 1, 10–15 m at Site 2, and 15–20 m at Sites 3–4.

Hypolimnetic oxygen loss is much less obvious in basin 3, in part due to the much larger hypolimnetic volume. Sites 3 and 4 often develop small oxygen sags near the thermocline during late summer. These are caused by respiration of heterotrophic bacteria that accumulate along the density gradient between the epilimnion and hypolimnion (e.g., Figure 2.1, page 28; Figure B4, page 105; Matthews and DeLuna, 2008). From October through December, which is usually the last month of stratification in basin 3, the hypolimnetic oxygen concentrations at Sites 3–4 are often lower than in the epilimnion, which likely reflects continued biological respiration in the isolated hypolimnion (e.g., Figures 2.3 & B14, pages 30 & 115). But the hypolimnion in basin three rarely drops below 5–6 mg/L of dissolved oxygen.<sup>11</sup>

---

<sup>10</sup> Correlation analyses examine the relationships between two variables. The test statistic ranges from -1 to +1; the closer to  $\pm 1$ , the stronger the correlation. The significance is measured using the p-value; significant correlations have p-values <0.05.

<sup>11</sup> From 1998–2019, the deep sample from Site 3 was taken at 80 m; however, this depth is very close to the lake bottom and was frequently contaminated by the bottom sediments. Starting in the 2019–2020 report, we no longer sample at 80 m and instead use 75 m as the deepest measurement.

**Hypolimnetic hydrogen sulfide:** Bacteria require an energy source (e.g., organic carbon) and an electron acceptor (e.g., oxygen) for basic growth and metabolism. Under anaerobic conditions, when oxygen is not available, there is a predictable sequence whereby different types of anaerobic bacteria use alternate electron acceptors.<sup>12</sup> First, bacteria will use nitrate as an alternate to oxygen, converting nitrate to nitrite and nitrogen gas. Next, bacteria use manganese and ferrous ions. When these compounds are exhausted, bacteria use sulfate, converting it to hydrogen sulfide, a colorless gas with a strong, rotten-egg smell. If the electron acceptors listed above are unavailable, bacteria can use carbon dioxide, converting it to methane.

Hydrogen sulfide is commonly present in anaerobic lake sediments, but if the overlying water contains oxygen the sulfide will be converted into sulfates or other compounds. If the overlying water is anaerobic, hydrogen sulfide can build up to detectable levels during stratification. Hydrogen sulfide is an indicator of the degree of anoxia in the hypolimnion because it will not persist in oxygenated waters and is formed after the nitrate, manganese, and ferrous ions are exhausted.

The hypolimnion at Sites 1–2 usually contain detectable concentrations of hydrogen sulfide by October (Table 2.9, page 26). Hydrogen sulfide concentrations are measured in October because that is the latest month that is consistently stratified at Sites 1–2, so the hydrogen sulfide concentrations should be near their highest levels. The values of hydrogen sulfide obtained from Site 1 were consistent with previous years; however, the values measured from Site 2 were relatively high. The hydrogen sulfide values in Site 2 correspond with elevated concentrations of ammonium in the hypolimnion – see Section 2.3.5 for a discussion (page 10).

### 2.3.3 Conductivity and pH

The pH and conductivity data followed trends that were fairly typical for Lake Whatcom (Figures B1–B50 and B76–B85, pages 102–151 and 178–187). Epilimnetic pH values increased during the summer due to photosynthetic activity and hypolimnetic pH values decreased due to decomposition and the release of dissolved compounds from the sediments (Figures B31–B45, pages 132–B45). Given the high water temperatures observed in July, it is unsurprising that in July 2021 we also observed the highest pH value (9.5) recorded anywhere in the lake

---

<sup>12</sup>For a more complete discussion of anaerobic decomposition in lakes, see Wetzel, 2001.



by the Institute for Watershed Studies since data collection began in 1988 (Figure B36, page 137).

The conductivity concentrations were elevated in hypolimnetic samples at Sites 1–2, and periodically at Site 3, coinciding with periods of low oxygen near the bottom (e.g., Figures B41 & B47, pages 142 & 148). The historic data show what appears to be a decreasing trend in the conductivity values from 1988–2002, but this was caused by using increasingly sensitive equipment during the past three decades and does not indicate any actual change in the conductivity in the lake (Matthews, et al., 2004). Occasional spikes in conductivity at Site 3 are associated with low oxygen in samples collected very close to the bottom sediments.

#### **2.3.4 Alkalinity and turbidity**

Lake Whatcom is a soft water lake so most alkalinity values were low ( $\leq 25$  mg/L; Figures B86–B90, pages 189–193). During the summer the alkalinity values at the bottom of Sites 1–2 increased due to decomposition and the release of dissolved compounds from the sediments into the lower portion of the water column.

Turbidity values in the lake were usually low (1–3 NTU) except during late summer in samples from near the bottom of the lake. The high turbidity levels during this time are an indication of increasing turbulence in the lower hypolimnion as the lake begins to destratify. The highest turbidity peaks were measured at Sites 1–2, followed by Site 3 and 4 (Figures B91–B95, pages 194–198).

Suspended sediments from storm events can also cause elevated turbidity levels in the lake. Major storm events usually occur during winter or early spring when the lake is destratified, so the turbidity levels will be high throughout the water column. Storm-related turbidity peaks are easier to see in samples from the Intake and Site 4 where there are fewer distracting late summer hypolimnetic turbidity peaks (e.g., Feb. 2009 turbidity peaks, Figures B93 and B95; pages 196 and 198).

#### **2.3.5 Nitrogen and phosphorus**

The nitrogen and phosphorus data are illustrated in Figures B96–B120 (pages 199–223). Nitrogen and phosphorus are important nutrients that influence the amount and type of microbiota (e.g., algae) that grow in the lake. We measured

inorganic forms of nitrogen and phosphorus (nitrite, nitrate, ammonium, and soluble phosphate) as well as total nitrogen and total phosphorus, which includes inorganic and organic compounds.<sup>13</sup>

**Nitrogen:** Most algae use dissolved inorganic nitrogen (DIN)<sup>14</sup> for growth. Nitrate depletion was evident at all sites in the photosynthetic zone during the summer (Figures B101–B105, pages 204–208), particularly at Site 1, where the epilimnetic nitrate concentrations usually drop below 20  $\mu\text{g-N/L}$  by the end of the summer. Because nitrogen is required for algal growth, depletion of epilimnetic DIN concentrations is an indirect way to measure phytoplankton productivity. And, because algal densities have been increasing throughout the lake, it was not surprising to find that the DIN concentrations were declining over time (Figure 2.18, page 45). But, unlike the other indicators of phytoplankton productivity (see **Indications of eutrophication**, beginning on page 14), the DIN trend has not stabilized in recent years.

A month-by-month analysis of near-surface DIN showed that water column concentrations have declined in general, not just in the summer (Figure 2.19, page 46). Summer DIN concentrations are most likely declining because of higher lake productivity, with phytoplankton depleting DIN through uptake into their cells. When the summer DIN concentrations were adjusted by subtracting them from the median spring DIN values ( $\Delta\text{DIN} = \text{DIN}_{\text{spring}} - \text{DIN}_{\text{summer}}$ ), the trend with year was only marginally significant or not statistically significant, depending on site (Figure 2.20, page 47). Because phytoplankton uptake of DIN would be lower in the spring, this weak trend observed when comparing spring and summer DIN suggests that the overall decline in DIN is not wholly the result of phytoplankton uptake.

The reason for the lake-wide drop in DIN is not known, but similar trends have been reported for lakes in the midwestern and northeastern region of the USA (Oliver, et al., 2017), lakes in the Sierra Nevada (Sickman, et al., 2003), lakes in

---

<sup>13</sup>Organic nitrogen and phosphorus comes from living or decomposing plants and animals, and may include bacteria, algae, leaf fragments, and other organic particles.

<sup>14</sup>Dissolved inorganic nitrogen includes ammonium, nitrate, and nitrite. Usually, epilimnetic concentrations of ammonium and nitrite are low, so DIN is nearly equivalent to nitrate. When DIN is not available, some algae can use organic nitrogen and some Cyanobacteria, and a few uncommon species of diatoms, can convert dissolved nitrogen gas to ammonia (not ammonium) via nitrogen fixation.

the Adirondacks (Waller, et al., 2012), as well as lakes and rivers in northern Italy (Rogora, et al., 2012). Most of these studies attribute the declining DIN concentrations to decreasing amounts of nitrogen entering lakes from atmospheric deposition, but without a detailed nitrogen budget analysis for Lake Whatcom, it would be premature to attribute the declining DIN to a specific cause. The implication, however, is that Lake Whatcom water quality conditions may become increasingly favorable for the growth of nitrogen-fixing Cyanobacteria, many of which are capable of releasing toxins. Recent summer algal counts from Lake Whatcom revealed that the lake contained many species of Cyanobacteria (Matthews, et al., 2012), but the nitrogen-fixing species were not abundant. It will be important to continue tracking the densities of Cyanobacteria in the lake and to watch for increases in the densities of nitrogen-fixing species.

Hypolimnetic nitrate concentrations dropped below  $20 \mu\text{g-N/L}$  at Sites 1–2 (Figures B101–B102, pages 204–205). In anaerobic environments, bacteria reduce nitrate ( $\text{NO}_3^-$ ) to nitrite ( $\text{NO}_2^-$ ) and nitrogen gas ( $\text{N}_2$ ). The historic data indicate that nitrate reduction has been common in the hypolimnion at Site 1, but was not common at Site 2 until the summer of 1999 (Figure B102, page 205). Since then, the only year that Site 2 hypolimnetic nitrate concentrations did not drop below  $20 \mu\text{g-N/L}$  was 2007. Matthews, et al. (2008) hypothesized that the 2007 results were caused by a combination of late spring stratification and early fall destratification, which shortened the period of anoxia in the hypolimnion.

Ammonium, along with hydrogen sulfide, is often an indicator of hypolimnetic anoxia.<sup>15</sup> Ammonium is readily taken up by plants as a growth nutrient. In oxygenated environments, ammonium is rarely present in high concentrations because it is rapidly converted to nitrate through biological and chemical processes. In low oxygen environments, like the hypolimnion at Sites 1–2, ammonium concentrations increase during late summer, reaching maximum concentrations just prior to destratification (Figures B96 & B97, pages 199 & 200). Elevated hypolimnetic ammonium concentrations have been common at both sites throughout the monitoring period, but beginning in 1999 the concentrations increased noticeably at Site 2 (Figure B97, page 200). The hypolimnion in Site 2 tends to be smaller than Site 1, potentially concentrating ammonium in a smaller volume of water. The October 2020 ammonium concentrations near the bottom of Site 1 ( $307 \mu\text{g-N/L}$

---

<sup>15</sup>Ammonium is produced during decomposition of organic matter; hydrogen sulfide is produced by bacteria that use sulfate ( $\text{SO}_4^{2-}$ ) instead of oxygen, creating sulfide ( $\text{S}^{2-}$ ) that reacts with hydrogen ions to form hydrogen sulfide ( $\text{H}_2\text{S}$ ). See hydrogen sulfide discussion on page 8.

at 20 meters) were consistent with previous years, but were much higher at the bottom of Site 2 (620  $\mu\text{g-N/L}$  at 20 meters). This is consistent with the relatively higher hydrogen sulfide concentrations at the bottom of Site 2 in 2020. Both Site 1 and Site 2 had relatively high values of ammonium at 20 meters in October 2021 (425 and 577  $\mu\text{g-N/L}$  at 20 meters for Sites 1–2, respectively). Both sites are usually destratified by November, which causes the ammonium concentrations to drop through winter and spring (see annual patterns in Figures B96 & B97, pages 199 & 200).

Sites 3–4 often have slightly elevated ammonium concentrations in the metalimnion at 20 m, or near the bottom at 80–90 m (Figures B99–B100, pages 202–203). This is caused by bacterial decomposition of organic matter, but the concentrations never approach the levels found in the hypolimnion at Sites 1–2.

**Phosphorus:** Although the Lake Whatcom microbiota require nitrogen, phosphorus is usually what limits microbial growth (Bittner, 1993; Liang, 1994; Matthews, et al., 2002a; McDonald, 1994). The total phosphorus concentration in the water column is a complex mixture of soluble and insoluble phosphorus compounds, only some of which can be used by algae to sustain growth. Soluble forms of phosphorus (e.g., orthophosphate) are easily taken up by algae and other microbiota, and, as a result, are rarely found in high concentrations in the water column. Insoluble phosphorus can be present in the water column bound to the surface of tiny particles or as suspended organic matter (e.g., live or dead algae). Some microbiota produce enzymes that release phosphorus from the surface of suspended soil particles. Liang (1994) and Groce (2011) demonstrated that  $\geq 50\%$  of the total phosphorus associated with soils in the Lake Whatcom watershed was potentially “bioavailable” through enzyme action. Algal growth tests revealed that 37–92% (median=78%) of the total phosphorus in storm runoff from Anderson, Austin, and Smith Creeks was bioavailable (Deacon, 2015).

When hypolimnetic oxygen concentrations are low, sediment-bound phosphorus becomes soluble and leaches into the overlying water. Prior to destratification, hypolimnetic phosphorus may be taken up by microbiota in the hypolimnion or metalimnion (see Section 2.3.2 and Matthews and DeLuna, 2008). When the lake mixes in the fall, the hypolimnetic phosphorus will be distributed throughout the water column. As oxygen concentrations increase during mixing, any soluble phosphorus that has not been taken up by biota will usually be converted back into insoluble phosphorus. Because phosphorus moves back and forth between

soluble and insoluble forms and between organic and inorganic compounds, it can be difficult to interpret total phosphorus trends. For example, when algal densities increase, their growth usually results in the reduction of soluble and bioavailable fractions of phosphorus in the epilimnion. This uptake moves the phosphorus into the “live-algae” fraction of organic phosphorus, which should show up in total phosphorus measurements. But algae are not distributed homogeneously in the water column (Matthews and Deluna, 2008), making it difficult to estimate the amount of phosphorus that is incorporated into algal biomass.

In Lake Whatcom, total phosphorus and soluble phosphate concentrations were usually low except in the hypolimnion at Sites 1–2 just prior to destratification (Figures B111–B115, pages 214–218 and B116–B120, pages 219–223). Epilimnetic total phosphorus concentrations are usually lower than late-summer hypolimnetic peaks. Prior to 2000, the median epilimnetic phosphorus concentrations were  $<5 \mu\text{g-P/L}$  at Sites 2–4 and approximately  $5\text{--}8 \mu\text{g-P/L}$  at Site 1 (Figure 2.21, page 48). Since 2000, the median epilimnetic phosphorus concentrations have often been in the detectable range ( $\geq 5 \mu\text{g-P/L}$ ), but the pattern is quite erratic, reflecting the complicated nature of phosphorus movement in the water column (Figure 2.21, page 48).

### 2.3.6 Chlorophyll, plankton, and Secchi depth

Site 1 continued to have the highest chlorophyll concentrations of all the sites (Figures B121–B125, pages 224–228). Peak chlorophyll concentrations were usually collected at 0–15 m, while samples from 20 m had relatively low chlorophyll concentrations because light levels are not optimal for algal growth at this depth.

The plankton counts (Figures B136–B145, pages 239–248) were usually dominated by golden algae (Chrysophyta). Substantial numbers of green algae (Chlorophyta) and bluegreen bacteria (Cyanobacteria) were also measured at all sites during summer and late fall. Previous analyses of algal biomass in Lake Whatcom indicated that although Chrysophyta dominate the numerical plankton counts, Chlorophyta and Cyanobacteria may dominate the plankton biomass, particularly in late summer and early fall (Ashurst, 2003; Matthews, et al., 2002b).

Secchi depths (Figures B126–B130, pages 229–233) showed no clear seasonal pattern because transparency in Lake Whatcom is affected by particulates from storm events as well as algal blooms.

**Indications of eutrophication:** Eutrophication is the term used to describe a lake that is becoming more biologically productive. It can apply to an unproductive lake that is becoming slightly more eutrophic, or a productive lake that is becoming extremely eutrophic (see Wetzel, 2001, for more about eutrophication and Matthews, et al., 2005, for a description of the chemical and biological indicators of eutrophication in Lake Whatcom).

Chlorophyll is a direct measure of algal biomass and generally provides a better indication of changes in the lake's biological productivity than phosphorus. Similarly, although algal counts are useful for looking at trends within the same type of algae (e.g., are the numbers of Cyanobacteria increasing?), cell counts are not as good as chlorophyll for estimating algal biomass. The actual relationship between chlorophyll and algae cell counts is complex. The amount of chlorophyll in a cell is influenced by the physiological age and condition of the cell, light intensity, nutrient availability, and many other factors. In addition, while most types of algae are counted by individual cells, a few types must be counted by colonies because the cells are too difficult to see.

The median near-surface summer chlorophyll concentrations have increased significantly at all sites since 1994 (Figure 2.22, page 49). Site 1 has shown the most year-to-year variability, which is reflected by a slightly lower correlation statistic compared to Sites 2–4 (Site 1 Kendall's  $\tau = 0.451$ ; Sites 2–4 Kendalls  $\tau = 0.517, 0.587, 0.536$ , respectively).<sup>16</sup> Although the annual chlorophyll concentrations are quite variable, the median near-surface summer concentrations seem to have stabilized since 2004, ranging from 3.7–6.7  $\mu\text{g/L}$  at Site 1 and 2.6–4.6  $\mu\text{g/L}$  at Sites 2–4 (Appendix B, page 100).

Under certain conditions and in certain lakes, a thin layer of algae can form deep in the water column (i.e., not at the surface) – this is known as a deep chlorophyll maximum. Deep chlorophyll maxima are thought to be a product of lake depth, stratification, and light, with deeper, relatively clear stratified lakes frequently observing this pattern (Fee, 1976). These deep chlorophyll maxima occur frequently in Lake Whatcom, with the highest values of chlorophyll often observed at 10 or 15 m (Figures B121–B125, pages 224–228). For example, the very high chlorophyll value at Site 1 in July 2010 was at 10 m. These layers can be thin (from a few centimeters to a few meters) and may not be observed with discrete sampling. Another way to detect them is to examine dissolved oxygen profiles, which will

---

<sup>16</sup>See discussion of correlation in footnote on page 7

spike near the deep chlorophyll layer because of increased algal photosynthesis (Figures 2.7–2.10, pages 34–37). For further discussion, see page 7.

Except for the dinoflagellates,<sup>17</sup> the algae counts have increased significantly since 1994 (Figure 2.23, page 50). Cyanobacteria, which are often used as bioindicators of eutrophication, have increased at all sites (Figure 2.24, page 51). The Cyanobacteria counts are dominated by *Aphanothece*, *Aphanocapsa*, *Cyanodictyon*, and *Snowella*, genera that are not usually associated with toxic blooms, but some of which have led to filter clogging incidences at the water treatment plant.<sup>18</sup> As with chlorophyll, the algae counts appear to have stabilized since 2004.

### 2.3.7 Coliform bacteria

This sampling period (2020–2021) spanned a change in the surface water standards based on “designated use” categories, which for Lake Whatcom is “Primary Contact Recreation,” described in Chapter 173–201A–200 of the Washington Administrative Code, Water Quality Standards for Surface Waters of the state of Washington. The standard for bacteria prior to December 31, 2020 was:

*Fecal coliform organism levels within an averaging period must not exceed a geometric mean value of 100 CFU or MPN per 100 mL, with not more than 10 percent of all samples (or any single sample when less than ten sample points exist) obtained within an averaging period exceeding 200 CFU or MPN per 100 mL.*

The standard for bacteria after December 31, 2020 is:

*E. coli organism levels within an averaging period must not exceed a geometric mean value of 100 CFU or MPN per 100 mL, with not more than 10 percent of all samples (or any single sample when less than ten sample points exist) obtained within the averaging period exceeding 320 CFU or MPN per 100 mL.*

<sup>17</sup>Dinoflagellates are small single-cell algae that are common in Lake Whatcom, but rarely have high densities in the plankton counts.

<sup>18</sup>P. Wendling, pers. comm., City of Bellingham Public Works Dept.

The city is currently in the process of examining whether fecal coliform data can transfer to the new *E. coli* standard, in collaboration with the Washington Department of Ecology.<sup>19</sup> During this transition phase, fecal coliform data will continue to be included in this annual report and the standard in place prior to 2021 will be used.

All of the mid-basin (Sites 1–4) and Intake values for fecal coliforms<sup>20</sup> were less than 10 cfu<sup>21</sup>/100 mL (Figures B131–B135, pages 234–238) and passed the freshwater *Primary Contact Recreation* bacteria standard in place prior to 2021.

Coliform samples collected offshore from the Bloedel-Donovan swimming area had slightly higher counts than at Site 1 (mid-basin). None of the Bloedel-Donovan counts exceeded 100 cfu/100 mL and the geometric mean was 3 cfu/100 mL, so this site passed both parts of the freshwater *Primary Contact Recreation* bacteria standard in place prior to 2021.

### 2.3.8 Total organic carbon and disinfection by-products

Total organic carbon concentrations, along with plankton and chlorophyll data, are used to help assess the likelihood of developing potentially harmful disinfection by-products through the reaction of chlorine with organic compounds during the drinking water treatment process. Algae excrete dissolved organic carbon into water, which can react with chlorine to form disinfection by-products, predominately chloroform and other trihalomethanes (THMs). When algal densities (or total organic carbon concentrations) increase, we expect to see an increase in THMs. To minimize risk, limits are set on the levels of disinfection by-products allowed in treated drinking water through the Safe Drinking Water Act’s Disinfection Byproduct Rule. This Rule was adopted in 1979 and has undergone two major revisions (Phase I in 1998; Phase II in 2005). The sampling requirement doubled under Phase II; currently the City samples eight locations in the water distribution system.<sup>22</sup>

The 2020/2021 total organic carbon concentrations ranged from 1.7–8.2 mg/L (AmTest) and 1.7–2.2 mg/L (IWS; Table 2.10, page 27). The samples were split

<sup>19</sup>P. Wendling, pers. comm., City of Bellingham Public Works Dept.

<sup>20</sup>Fecal coliforms are currently called “thermotolerant” coliforms (APHA, 2017).

<sup>21</sup>Colony forming unit/100 mL; cfu/100 mL is sometimes labeled “colonies/100 mL.”

<sup>22</sup>P. Wendling, pers. comm., City of Bellingham Public Works Dept.



and analyzed by AmTest and the IWS laboratory to compare results. The median difference between AmTest and IWS concentrations was  $\pm 0.5$  mg/L. Larger differences could have been caused by small particulates that were unevenly distributed in the split samples or differences in the analytical methodologies.

The 2020/2021 THMs and HAAs remained below the maximum contaminant levels of 0.080 mg/L and 0.060 mg/L, respectively, described in Chapter 246–290–310 of Washington Administrative Code, Water Quality Standards for Public Water Supplies. The THMs concentrations (1991–2021) have showed a significant increase over time, particularly during the spring and summer (Quarters 2–3; Figures 2.25–2.26, pages 52–53), when algal densities are higher. However, in recent years THMs concentrations appear to be declining, likely due to operational changes by the City of Bellingham.<sup>23</sup>

Haloacetic acids (HAAs), another type of disinfection by-product, also increased significantly over time, with more stable concentrations during the past 5–6 years (Figure 2.27, page 54). Only the winter and spring HAAs data followed this trend (Quarters 1–2); the summer and fall data were not significantly correlated with year (Figure 2.28, page 55). According to Sung, et al. (2000), HAAs are not as closely linked to algal concentrations and chlorine dose as THMs. In addition, HAAs can be degraded by the microbial biofilm that grows on the surface of water treatment filtration media (Baribeau, et al., 2005). Although microbial biofilm on filtration media can be a major site of HAA degradation (Grigorescu and Hozalski, 2010), bioremediation is thought to be occurring in the City of Bellingham’s distribution system by pipe and reservoir biofilm.<sup>24</sup>

---

<sup>23</sup>P. Wendling, pers. comm., City of Bellingham Public Works Dept.

<sup>24</sup>P. Wendling, pers. comm., City of Bellingham Public Works Dept.

Abbrev.	Parameter	Method	Historic DL <sup>†</sup>	2020/2021 MDL <sup>†</sup>	Sensitivity or Confidence limit
<b>IWS field measurements:</b>					
cond	Conductivity	YSI (2017)	–	–	± 2 µS/cm
do	Dissolved oxygen	YSI (2017)	–	–	± 0.1 mg/L
ph	pH	YSI (2017)	–	–	± 0.1 pH unit
temp	Temperature	YSI (2017)	–	–	± 0.1° C
disch	Discharge	Rantz et al. (1982); SOP-IWS-6	–	–	–
secchi	Secchi depth	Lind (1985)	–	–	± 0.1 m
<b>IWS laboratory analyses:</b>					
alk	Alkalinity	APHA (2017) #2320; SOP-IWS-8	–	–	± 0.4 mg/L
cond	Conductivity	APHA (2017) #2510; SOP-IWS-8	–	–	± 2.1 µS/cm
do	Dissolved oxygen	APHA (2017) #4500-O.C.; SOP-IWS-8	–	–	± 0.1 mg/L
ph	pH-lab	APHA (2017) #4500-H <sup>+</sup> ; SOP-IWS-8	–	–	± 0.1 pH unit
tss	T. suspended solids	APHA (2017) #2540 D; SOP-IWS-13	2 mg/L	2.1 mg/L	± 3.8 mg/L
turb	Turbidity	APHA (2017) #2130; SOP-IWS-8	–	–	± 0.2 NTU
nh4	Ammonium (auto)	APHA (2017) #4500-NH <sub>3</sub> H; SOP-IWS-19	10 µg-N/L	6.4 µg-N/L	± 9.7 µg-N/L
no3	Nitrite/nitrate (auto)	APHA (2017) #4500-NO <sub>3</sub> I; SOP-IWS-22	20 µg-N/L	24.7 µg-N/L	± 26.9 µg-N/L
tn	T. nitrogen (auto)	APHA (2017) #4500-N C; SOP-IWS-22	100 µg-N/L	48.6 µg-N/L	± 45.2 µg-N/L
srp	Sol. phosphate (auto)	APHA (2017) #4500-P G; SOP-IWS-22	5 µg-P/L	2.8 µg-P/L	± 1.1 µg-P/L
tp	T. phosphorus (auto)	APHA (2017) #4500-P J; SOP-IWS-22	5 µg-P/L	2.0 µg-P/L	± 2.1 µg-P/L
toc <sup>‡</sup>	T. organic carbon	EPA #414.3; Potter and Wimsatt (2009)	1.0 mg/L	0.09 mg/L	± 0.11 mg/L
<b>IWS plankton analyses:</b>					
chl	Chlorophyll	APHA (2017) #10200 H; SOP-LW-16	–	–	± 0.1 µg/L
chlo	Chlorophyta	Lind (1985), Schindler trap	–	–	–
cyan	Cyanobacteria	Lind (1985), Schindler trap	–	–	–
chry	Chrysophyta	Lind (1985), Schindler trap	–	–	–
pyrr	Pyrrophyta	Lind (1985), Schindler trap	–	–	–
<b>City coliform analyses:</b>					
fc	Fecal coliform <sup>§</sup>	APHA (2017) #9222 D	1 cfu/100 mL	1 cfu/100 mL	–
<b>Edge Analytical analyses:</b>					
H <sub>2</sub> S	Hydrogen sulfide	APHA (2017) #4500-S2 F	–	0.044 mg/L	–
<b>AmTest analyses:</b>					
toc <sup>‡</sup>	T. organic carbon	APHA (2017) #5310 B	1.0 mg/L	0.5 mg/L	–

<sup>†</sup>Historic detection limits (DL) are usually higher than current method detection limits (MDL).

<sup>‡</sup>Total organic carbon analyses are run in duplicate by IWS and AmTest to evaluate analytical equivalence.

<sup>§</sup>Fecal coliforms are now called thermotolerant coliforms (APHA 2017).

Table 2.1: Summary of IWS, AmTest, Edge Analytical, and City of Bellingham analytical methods and parameter abbreviations.

Parameter	Feb	Apr	May	Jun	Jul	Aug	Sep	Oct <sup>†</sup>	Nov <sup>†</sup>	Dec <sup>†</sup>	Locations <sup>‡</sup>
DO - field	•	•	•	•	•	•	•	•	•	•	Sites 1, 2, Intake - every 1 m;
pH - field	•	•	•	•	•	•	•	•	•	•	Sites 3, 4 - every 1 m to 10 m
Temp - field	•	•	•	•	•	•	•	•	•	•	then every 5 m; Gatehouse
Cond - field	•	•	•	•	•	•	•	•	•	•	
Secchi depth	•	•	•	•	•	•	•	•	•	•	Sites 1, 2, 3, 4, Intake
Alkalinity	•	•	•	•	•	•	•	•	•	•	Sites 1, 2 - 0.3, 5, 10, 15, 20 m;
Ammonium	•	•	•	•	•	•	•	•	•	•	Intake - 0.3, 5, 10 m;
Nitrate/nitrite	•	•	•	•	•	•	•	•	•	•	Site 3 - 0.3, 5, 10, 20, 40, 60,
T. nitrogen	•	•	•	•	•	•	•	•	•	•	75 m; Site 4 - 0.3, 5, 10, 20, 40
Sol. phosphate	•	•	•	•	•	•	•	•	•	•	60, 80, 90 m; Gatehouse
T. phosphorus	•	•	•	•	•	•	•	•	•	•	
Turbidity	•	•	•	•	•	•	•	•	•	•	
T. organic carbon	•				•						Sites 1, 2, 3, 4, Intake - 0.3 m and bottom only
Chlorophyll	•	•	•	•	•	•	•	•	•	•	Sites 1, 2, 3, 4 - 0.3, 5, 10, 15, 20 m; Intake - 0.3, 5, 10 m
Plankton	•	•	•	•	•	•	•	•	•	•	Sites 1, 2, 3, 4, Intake; 5 m
Bacteria (City)	•	•	•	•	•	•	•	•	•	•	Sites 1, 2, 3, 4, Intake, Bloedel-Donovan; 0.3 m
H <sub>2</sub> S - opt								•			Sites 1, 2 - 10, 15, 20 m

<sup>†</sup>Samples will be collected Feb-Dec in 2019 and 2020; field work will end in September 2021 to allow time to complete all analyses unless the monitoring contract is extended past December 2021.

<sup>‡</sup>Samples within each parameter subgroup are collected at all locations listed in this column.

Table 2.2: Lake Whatcom proposed lake monitoring schedule. All field and laboratory methods are summarized in Table 2.1; missing data resulting from the sampling and laboratory issues are summarized in Table 2.3.

Month	Missing Sample Summary	Comments
October 2020	No missing data	
November 2020	No missing data	
December 2020	No missing data	
February 2021	No missing data	
April 2021	No missing data	
May 2021	Temperature, DO, pH, and conductivity data missing from Site 2 at 11–14 m and 16–19 m, Site 3 at 3 m Temperature, DO, pH, and conductivity data missing from Site 2 at 20 m	missed collecting sample data rejected because YSI probe was in the lake sediment
June 2021	No missing data	
July 2021	Total nitrogen data missing for all sites and depths Chlorophyll missing from Site 3 at 15 m	unacceptable laboratory variability missed collecting sample
August 2021	Alkalinity data missing for Site 3 at surface	missed collecting sample
September 2021	No missing data	

Table 2.3: Summary of missing lake data due to sampling or laboratory issues.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	18.6	21.1	21.2	27.9
Conductivity (μS/cm)	57.8	59.4	60.9	77.1
Dissolved oxygen (mg/L)	0.0	9.6	8.3	13.0
pH	6.4	7.3	7.3	9.5
Temperature (°C)	5.5	10.1	11.6	24.2
Turbidity (NTU)	0.5	0.8	1.0	4.9
Nitrogen, ammonium (μg-N/L)	<10	<10	36.0	307.2
Nitrogen, nitrate/nitrite (μg-N/L)	<20	148.3	137.8	322.8
Nitrogen, total (μg-N/L)	<100	311.2	284.9	486.9
Phosphorus, soluble (μg-P/L)	<5	1.1	1.5	37.9
Phosphorus, total (μg-P/L)	5.3	7.5	10.8	57.9
Chlorophyll (μg/L)	0.5	3.2	3.4	14.3
Secchi depth (m)	2.7	4.2	4.1	5.0
Coliforms, fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	1

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 2.4: Summary of Site 1 water quality data, Oct. 2020 – Sept. 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	18.2	20.2	20.0	21.1
Conductivity (μS/cm)	56.8	57.7	57.8	59.0
Dissolved oxygen (mg/L)	8.1	10.5	10.3	12.0
pH	6.6	7.6	7.7	8.6
Temperature (°C)	6.2	14.1	14.5	22.8
Turbidity (NTU)	0.4	0.5	0.5	0.7
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	19.8
Nitrogen, nitrate/nitrite (μg-N/L)	<20	159.2	166.4	341.9
Nitrogen, total (μg-N/L)	<100	308.5	258.4	420.2
Phosphorus, soluble (μg-P/L)	<5	<5	<5	<5
Phosphorus, total (μg-P/L)	<5	<5	<5	8.3
Chlorophyll (μg/L)	1.8	2.7	2.9	4.5
Secchi depth (m)	4.5	5.4	5.2	6.2
Coliforms, fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	1

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 2.5: Summary of Intake water quality data, Oct. 2020– Sept. 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.4	20.4	20.5	28.3
Conductivity (μS/cm)	56.7	57.7	58.9	90.9
Dissolved oxygen (mg/L)	0.0	9.8	9.1	12.0
pH	6.5	7.3	7.4	8.7
Temperature (°C)	6.2	11.8	12.8	22.7
Turbidity (NTU)	0.3	0.6	0.8	6.1
Nitrogen, ammonium (μg-N/L)	<10	<10	43.7	619.9
Nitrogen, nitrate/nitrite (μg-N/L)	<20	199.5	177.5	453.8
Nitrogen, total (μg-N/L)	<100	336.5	324.9	791.8
Phosphorus, soluble (μg-P/L)	<5	<5	<5	31.8
Phosphorus, total (μg-P/L)	<5	<5	9.5	79.9
Chlorophyll (μg/L)	0.6	2.6	2.7	5.8
Secchi depth (m)	4.2	5.3	5.5	6.8
Coliforms, fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	1

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 2.6: Summary of Site 2 water quality data, Oct. 2020– Sept. 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.5	18.8	19.3	21.2
Conductivity ( $\mu$ S/cm)	56.8	57.3	57.9	62.4
Dissolved oxygen (mg/L)	6.8	10.0	9.9	12.0
pH	6.4	7.1	7.3	8.4
Temperature (°C)	6.4	7.6	10.6	22.0
Turbidity (NTU)	0.1	0.4	0.4	0.8
Nitrogen, ammonium ( $\mu$ g-N/L)	<10	<10	<10	21.7
Nitrogen, nitrate/nitrite ( $\mu$ g-N/L)	<20	305.2	261.6	401.8
Nitrogen, total ( $\mu$ g-N/L)	117.6	378.4	346	477.5
Phosphorus, soluble ( $\mu$ g-P/L)	<5	<5	<5	<5
Phosphorus, total ( $\mu$ g-P/L)	<5	<5	<5	17.8
Chlorophyll ( $\mu$ g/L)	0.8	2.6	2.7	5.5
Secchi depth (m)	5.0	5.7	6.0	7.8
Coliforms, fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	1

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1  $\Rightarrow$  1).

Table 2.7: Summary of Site 3 water quality data, Oct. 2020– Sept. 2021.



Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	15.0	18.7	19.1	21.1
Conductivity (μS/cm)	56.8	57.3	57.8	61.8
Dissolved oxygen (mg/L)	7.8	9.9	9.9	12.1
pH	6.3	7.0	7.2	8.4
Temperature (°C)	6.5	7.1	10.2	21.9
Turbidity (NTU)	0.2	0.4	0.4	0.6
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	20.0
Nitrogen, nitrate/nitrite (μg-N/L)	<20	339.2	289.9	410.3
Nitrogen, total (μg-N/L)	123.2	403.2	378.5	574.5
Phosphorus, soluble (μg-P/L)	<5	<5	<5	<5
Phosphorus, total (μg-P/L)	<5	<5	<5	9.0
Chlorophyll (μg/L)	0.7	2.5	2.7	5.8
Secchi depth (m)	5.0	6.2	6.5	8.5
Coliforms, fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	1

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 2.8: Summary of Site 4 water quality data, Oct. 2020– Sept. 2021.

Year	H <sub>2</sub> S (mg/L)		Year	H <sub>2</sub> S (mg/L)	
	Site 1	Site 2		Site 1	Site 2
1999 <sup>†</sup>	0.03–0.04	0.40	2011	0.12	0.16
2000 <sup>†</sup>	0.27	0.53	2012	na	na
2001 <sup>†</sup>	0.42	0.76	2013	0.20 <sup>§</sup>	0.16
2002 <sup>†</sup>	0.09	0.32	2014	0.28	0.66
2003 <sup>†</sup>	0.05	0.05	2015	0.51	0.41
2004 <sup>†</sup>	0.25	0.25	2016	0.64	0.51
2005 <sup>‡</sup>	0.13, 0.12	0.25, 0.42	2017	0.68*	<0.05
2006	0.20	0.42	2018	0.32	0.39
2007	0.40	0.20	2019	0.10	0.22
2008	0.28	0.38	2020	0.29	0.51
2009	0.15	0.47	2021	0.25	0.51
2010	0.38	0.40			

<sup>†</sup>H<sub>2</sub>S samples analyzed by HACH test kit.

<sup>‡</sup>HACH (first value) vs. Edge Analytical (second value)

<sup>§</sup>Corrected value (1.20 in Matthews, et al., 2015)

\*Sample collected at 15 meters; sample from 20 m contained sediment.

Table 2.9: October hypolimnetic hydrogen sulfide concentrations at Sites 1 and 2 (20 m). The H<sub>2</sub>S samples have been analyzed by Edge Analytical since 2005; earlier samples were analyzed using a HACH field test kit.

Site	Depth (m)	Date	AmTest TOC (mg/L)	IWS TOC (mg/L)	Date	AmTest TOC (mg/L)	IWS TOC (mg/L)
Site 1	0	Feb 27, 2021	2.2	1.8	July 6, 2021	2.7	2.2
	20	Feb 27, 2021	2.2	1.9	July 6, 2021	8.2	1.9
Intake	0	Feb 27, 2021	2.0	1.8	July 8, 2021	2.7	2.2
	10	Feb 27, 2021	2.4	1.8	July 8, 2021	2.9	2.1
Site 2	0	Feb 24, 2021	2.1	1.7	July 8, 2021	2.5	2.1
	20	Feb 24, 2021	2.1	1.8	July 8, 2021	2.1	1.9
Site 3	0	Feb 17, 2021	4.0	1.8	July 8, 2021	2.7	2.2
	75	Feb 17, 2021	3.6	1.7	July 8, 2021	2.0	1.7
Site 4	0	Feb 17, 2021	1.7	1.7	July 6, 2021	2.4	2.0
	90	Feb 17, 2021	4.3	1.7	July 6, 2021	1.9	1.7

Table 2.10: Lake Whatcom 2020/2021 total organic carbon data. February and July samples were split and analyzed by AmTest (TOC-AM) and IWS (TOC-IWS). Differences can be expected when concentrations are low and if there are particles present in one sample but not the other.




Figure 2.1: October 2020 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2020 data.




Figure 2.2: November 2020 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2020 data.




Figure 2.3: December 2020 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2020 data.




Figure 2.4: February 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.




Figure 2.5: April 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.






Figure 2.6: May 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.




Figure 2.7: June 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.




Figure 2.8: July 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.




Figure 2.9: August 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.




Figure 2.10: September 2021 temperature (---●---) and dissolved oxygen (---●---) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data.




Figure 2.11: October 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data. October 2021 is not part of the 2020/2021 sampling period, but were included to to provide information on the timing of destratification.




Figure 2.12: November 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data. November 2021 is not part of the 2020/2021 sampling period, but were included to to provide information on the timing of destratification.




Figure 2.13: December 2021 temperature (-●-) and dissolved oxygen (-●-) profiles compared to historic ranges. The gradation in shading shows the approximate frequency of the 1988–2021 data. December 2021 is not part of the 2020/2021 sampling period, but were included to to provide information on the timing of destratification.






Figure 2.14: Relationship between dissolved oxygen and time at Site 1, 12 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.15: Relationship between dissolved oxygen and time at Site 1, 14 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.16: Relationship between dissolved oxygen and time at Site 1, 16 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.

Site 1 Dissolved Oxygen by Year at Depth 18




Figure 2.17: Relationship between dissolved oxygen and time at Site 1, 18 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.18: Minimum summer, near-surface dissolved inorganic nitrogen (DIN) concentrations (1994–2021 June–Oct, depths  $\leq 5$  m). Uncensored (raw) data were used to illustrate that minimum values are dropping below analytical detection limits (dashed red line); negative values represent regression results for concentrations below the detection limit. Note differences in y-axis scale between Site 1 and Sites 2–4. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.19: Comparison of median spring (Feb-May) vs. summer (June-Oct) near-surface dissolved inorganic nitrogen (DIN) concentrations (1994–2021, depths  $\leq 5$  m). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.20: Differences between median spring (Feb-May) and summer (June-Oct) near-surface dissolved inorganic nitrogen (DIN) concentrations (1994–2021, depths  $\leq 5$  m; DIN difference =  $DIN_{spring} - DIN_{summer}$ ). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; correlations were marginally significant (p-value  $< 0.05$ ) or not significant (p-value  $> 0.05$ ).




Figure 2.21: Median summer, near-surface total phosphorus concentrations (1994–2021, June–Oct, depths  $\leq 5$  m). Uncensored (raw) data were used to illustrate when median values are below analytical detection limits (dashed red line). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; none of the correlations were significant.






Figure 2.22: Median summer near-surface chlorophyll concentrations (1994–2021, June–October, depths  $\leq 5$  m). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.23:  $\text{Log}_{10}$  plots of median summer, near-surface algae counts (1994-2021, June-October, all sites and depths). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations except Dinoflagellates were significant. Note difference in vertical axis scales.




Figure 2.24: Log<sub>10</sub> plots of median summer, near-surface Cyanobacteria counts (1994–2021, June–October, depths  $\leq 5$  m). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 2.25: Total trihalomethanes (THMs) quarterly average concentrations in the Bellingham water distribution system (data provided by the City of Bellingham Public Works Department). The recommended maximum contaminant level for total THMs is 0.080 mg/l; all samples were below the level. The number of sites used to calculate the quarterly averages increased from four to eight in the fourth quarter of 2012 (vertical red line). Kendall's  $\tau$  correlation was used because the data were not monotonic-linear; the correlation was significant.




Figure 2.26: Total trihalomethanes (THMs) quarterly average concentrations in the Bellingham water distribution system plotted by quarter (data provided by the City of Bellingham Public Works Department). The recommended maximum contaminant level for total THMs is 0.080 mg/l; all samples were below the level. The number of sites used to calculate the quarterly averages increased from four to eight in the fourth quarter of 2012 (vertical red line). Kendall’s  $\tau$  correlations were used because the data were not monotonic-linear; all correlations of total THMs with time were significant.




Figure 2.27: Haloacetic acids (HAAs) quarterly average concentrations in the Bellingham water distribution system (data provided by the City of Bellingham Public Works Department). The recommended maximum contaminant level for HAAs is 0.060 mg/l; all samples were below the level. The number of sites used to calculate the quarterly averages increased from four to eight in the fourth quarter of 2012 (vertical red line). Kendall's  $\tau$  correlation was used because the data were not monotonic-linear; the correlation was significant.




Figure 2.28: Haloacetic acids (HAAs) quarterly average concentrations in the Bellingham water distribution system plotted by quarter (data provided by the City of Bellingham Public Works Department). The recommended maximum contaminant level for HAAs is 0.060 mg/l; all samples were below the level. The number of sites used to calculate the quarterly averages increased from four to eight in the fourth quarter of 2012 (vertical red line). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; the Quarter 1 and Quarter 2 correlations of HAAs with time were significant.

### 3 Tributary Monitoring

The major objective for the tributary monitoring was to provide baseline water quality data for the tributaries that flow into Lake Whatcom. Whatcom Creek was also sampled to provide baseline data for the lake's outlet. Monthly samples were collected in 2004–2006, 2010–2012, and 2014. The level of effort was reduced in 2007–2009, with samples collected twice each year. Monthly sampling was re-initiated in January 2016 and has continued through 2021.

#### 3.1 Site Descriptions

Samples were collected from Anderson, Austin, Blue Canyon, Brannian, Carpenter, Euclid, Millwheel, Olsen, Silver Beach, Smith, and Whatcom Creeks and the Park Place drain. The sampling locations for these sites are described in Appendix A.2 and shown on Figure A2, page 99.

#### 3.2 Field Sampling and Analytical Methods

The tributaries were sampled on October 12, November 16, and December 16, 2020; and January 13 & 20, February 3, March 10, April 13, May 11, June 9, July 13, August 10, and September 14, 2021. All samples were collected during daylight hours, typically between 10:00 am and 3:00 pm. The analytical and sampling procedures are summarized in Tables 2.1 & 3.1 (pages 18 & 61). Table 3.2 (page 62) summarizes missing data from the 2020/2021 sampling season.

A YSI ProDO LDO field meter was used to measure temperature and dissolved oxygen in the field. Raw water and bacteriological samples were stored on ice and in the dark until they reached the laboratory. The bacteria samples were analyzed by the City of Bellingham and total organic carbon analyses were analyzed by AmTest<sup>25</sup> and by IWS.

---

<sup>25</sup> AmTest, 13600 Northeast 126th Place, Suite C, Kirkland, WA, 98034–8720.



### 3.3 Results and Discussion

The tributary data include field measurements (dissolved oxygen and temperature); laboratory analyses for ambient water quality parameters (ammonium,<sup>26</sup> nitrate/nitrite,<sup>27</sup> total nitrogen, soluble phosphate, total phosphorus, alkalinity, conductivity, pH, total suspended solids, and turbidity); bacteria counts; and total organic carbon measurements.

The 2020/2021 tributary data are summarized in Table 3.3 (page 63), with descriptive statistics for each site listed in Tables 3.4–3.15 (pages 64–75). The total organic carbon data are listed in Table 3.16 (page 76). Because of missing samples during the 2020/2021 field season due to insufficient flow in creeks to sample (see Table 3.2), the summary statistics for these sites are biased toward water quality conditions present during spring, fall, and winter, with less representation of summer conditions.

Historical tributary data from 2004 to the present are plotted in Appendix B.4 (Figures B146–B184, pages 251–289). These figures include a dashed (blue) horizontal line that shows the median value for Smith Creek and a solid (red) horizontal line that shows the median value for each site. Smith Creek was chosen as a reference because it is a major tributary to the lake and has a history of being relatively unpolluted.

In Table 3.3, the “typical ranges” for alkalinity, conductivity, total suspended solids, ammonium, and soluble phosphate were derived from historic water quality data for Lake Whatcom tributaries that flow through predominantly forested portions of the watershed (Anderson, Brannian, Olsen, and Smith Creeks). The temperature, dissolved oxygen, and pH ranges were based on WAC 173-201A, Tables 200 (1)(c), 200(1)(d), and 200 (1)(g) for salmonid spawning, rearing, and migration, with the qualification that the single monthly grab samples from the Lake Whatcom tributaries may not show the lowest 1-day minimum dissolved oxygen or the maximum 7-day temperature. The turbidity range was based on historic watershed data and WAC 173-201A Table 200 (1)(e), which limits anthropogenic contributions to no more than 5 NTU over background. The coliform

<sup>26</sup>Nearly all ammonia ( $\text{NH}_4^+$ ) is ionized to ammonium ( $\text{NH}_3$ ) in surface water. Earlier IWS reports used “ammonia” and “ammonium” interchangeably; we now use “ammonium” to indicate that the data represent the concentration of ionized ammonia.

<sup>27</sup>Nitrate and nitrite were analyzed together because nitrite concentrations are very low in surface water. For simplicity, nitrate/nitrite will be referred to as “nitrate” in this document.

range was based on the WAC 173-201A Table 200 (2)(b) standard for *Primary Contact Recreation* in place prior to 2021.<sup>28</sup> The total phosphorus range was based on the lake nutrient criteria action value for the Coast Range, Puget Lowlands, and Northern Rockies Ecoregions listed from WAC 173-201A-230, Table 230(1). The lake nutrient criteria require collecting multiple samples from the epilimnion during summer, so the total phosphorus range in Table 3.3 can only be used as a general reference.

Water temperatures and dissolved oxygen concentrations followed typical seasonal cycles, with most sites having colder temperatures and higher oxygen concentrations during the winter, and warmer temperatures and lower oxygen concentrations during the summer (Figures B146–B151). Whatcom Creek had higher temperatures and slightly lower oxygen concentrations than most other sites, reflecting the influence of Lake Whatcom (Figures B146 and B149). Corresponding with high water temperatures in the lake, the July water temperature in Whatcom Creek (26.2°C) was the highest stream temperature value recorded by the Institute at any tributary site that we monitor since 2004.

The residential tributaries (Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain) often have slightly elevated temperatures and lower dissolved oxygen concentrations (Figures B148 and B151). But the dry conditions meant that some of the residential sites could not be sampled during late summer when high temperatures and low oxygen concentrations are most common.

Most of the tributaries in the Lake Whatcom watershed had relatively low concentrations of dissolved solids, indicated by conductivities  $\leq 100 \mu\text{S}$  and alkalinities  $\leq 30 \text{ mg/L}$  (Table 3.3; Figures B152–B160). Sites that did not match this description included some residential tributaries (Silver Beach Creek and the Park Place drain) and Blue Canyon Creek, which drains an area rich in soluble minerals. Most sites also had low total suspended solids concentrations ( $\leq 5 \text{ mg/L}$ ) and low turbidities ( $\leq 5 \text{ NTU}$ ) except during periods of high precipitation and runoff (Figures B161–B166). The only site that had consistently high solids and turbidity values was Millwheel Creek, which is often turbid due to disturbed sediments in an upstream pond.

The median ammonium concentrations were generally low ( $\leq 10 \mu\text{g-N/L}$ ) except in Millwheel Creek (Table 3.3; Figures B167–B169). Ammonium does not persist long in oxygenated surface waters. When present in streams, it usually indicates

---

<sup>28</sup>See Section 2.3.7, page 15 for discussion.

a near-by source such as an upstream wetland with anaerobic soils or a pollution source.

Most of the tributaries had lower total nitrogen and nitrate concentrations than Smith Creek (Figures B170– B175). The relatively high nitrate and total nitrogen concentrations in Smith Creek are probably due to the presence of nitrogen-fixing alders (*Alnus rubra*) in the riparian zone upstream from the sampling site. High nitrate and total nitrogen concentrations are not necessarily an indication of water pollution, and low nitrate concentrations actually favor the growth of nuisance Cyanobacteria. The exceptionally low nitrate concentrations in Whatcom Creek (Figure B170) reflect algal uptake of nitrogen in the lake.

Soluble inorganic phosphate is quickly removed from surface water by biota, so high concentrations of soluble phosphate usually indicate a near-by source such as an anaerobic wetland or a pollution source. The median 2020/2021 soluble phosphate concentrations were  $\leq 10$   $\mu\text{g-P/L}$  at all sites except Silver Beach Creek and the Park Place drain (Table 3.3). The historical data indicate that although soluble phosphate concentrations were generally low, nearly all sites have had a few high peaks, and high concentrations were common in residential streams.

Total phosphorus concentrations were higher than soluble phosphate concentrations (Figures B176–B181). The median 2020/2021 concentrations were  $\leq 20$   $\mu\text{g-P/L}$  at all sites except Millwheel Creek, Silver Beach Creek, and the Park Place drain (Table 3.3). As with soluble phosphate, nearly all sites have had occasional high total phosphorus peaks.

High coliform counts are an indicator of residential pollution (Table 3.3; Figures B182–B184). Although most of the sites had relatively low coliform counts during 2020/2021, three sites exceeded a geometric mean of 100 cfu/100 mL (Millwheel Creek, Silver Beach Creek and the Park Place drain). Seven sites (Olsen, Carpenter, Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain) had more than 10% of the samples that exceeded 200 cfu/100 mL. Several of the small residential tributaries could not be sampled during the late summer (see Table 3.2), when coliform counts are often higher, so these sites may have exceeded the coliform criteria by a greater margin than what is indicated in the summary tables.

The total organic carbon concentrations from February and July 2021 are included in Table 3.16 (page 76). Several of the residential sites (Millwheel and Silver Beach Creeks and the Park Place drain) had slightly elevated ( $\geq 3$  mg/L) total

organic carbon concentrations. Blue Canyon, Brannian, Carpenter, Euclid, Mill-wheel, and Silver Beach could not be sampled in July 2021 because of insufficient flow (Table 3.2). The paired samples analyzed by IWS and AmTest were very similar, with a median difference of 0.3 mg/L. Larger differences could have been caused by small particulates that were unevenly distributed in the split samples or differences in analytical methodologies.

Parameter	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
DO -field	•	•	•	•	•	•	•	•	•	•	•	•
pH - lab	•	•	•	•	•	•	•	•	•	•	•	•
Temp - field	•	•	•	•	•	•	•	•	•	•	•	•
Cond - lab	•	•	•	•	•	•	•	•	•	•	•	•
Alkalinity	•	•	•	•	•	•	•	•	•	•	•	•
Ammonium	•	•	•	•	•	•	•	•	•	•	•	•
Nitrate/nitrite	•	•	•	•	•	•	•	•	•	•	•	•
T. nitrogen	•	•	•	•	•	•	•	•	•	•	•	•
Sol. phosphate	•	•	•	•	•	•	•	•	•	•	•	•
T. phosphorus	•	•	•	•	•	•	•	•	•	•	•	•
T. susp. solids	•	•	•	•	•	•	•	•	•	•	•	•
Turbidity	•	•	•	•	•	•	•	•	•	•	•	•
T. organic carbon		•					•					
Bacteria (City)	•	•	•	•	•	•	•	•	•	•	•	•

Table 3.1: Lake Whatcom proposed tributary monitoring schedule. All field and laboratory methods are summarized in Table 2.1.

Month	Sample Summary	Comments
October 2020	No missing data	
November 2020	No missing data	
December 2020	No missing data	
January 2021	No missing data	
February 2021	No missing data	
March 2021	No missing data	
April 2021	No missing data	
May 2021	No missing data	
June 2021	No missing data	
July 2021	All field and laboratory data missing for Blue Canyon, Brannian, Carpenter, Euclid, Mill-wheel, and Silver Beach Creeks	insufficient flow
August 2021	All field and laboratory data missing for Blue Canyon, Brannian, Carpenter, Euclid, Mill-wheel, and Silver Beach Creeks	insufficient flow
September 2021	All field and laboratory data missing for Blue Canyon, Brannian, Carpenter, Euclid, Mill-wheel, and Silver Beach Creeks	insufficient flow

Table 3.2: Summary of missing tributary data due to sampling or laboratory issues.

	Typical range	Anderson	Austin	Brannian	Olsen	Smith	Whatcom
Alkalinity	med. $\leq 30$ mg/L	yes	yes	yes	yes	yes	yes
Conductivity	med. $\leq 100$ $\mu$ S	yes	yes	yes	yes	yes	yes
D. oxygen <sup>†</sup>	min. $\geq 8.0$ mg/L	no	yes	yes	yes	yes	yes
pH	6.5–8.5	yes	yes	yes	yes	yes	yes
Temperature <sup>†</sup>	max. $\leq 17.5$ C	yes	yes	yes	yes	yes	no
T. susp. solids	med. $\leq 5$ mg/L	yes	yes	yes	yes	yes	yes
Turbidity	med. $\leq 5$ NTU	yes	yes	yes	yes	yes	yes
Ammonium	med. $\leq 10$ $\mu$ g-N/L	yes	yes	yes	yes	yes	yes
Sol. phosphate	med. $\leq 10$ $\mu$ g-P/L	yes	yes	yes	yes	yes	yes
T. phosphorus	med. $\leq 20$ $\mu$ g-P/L	yes	yes	yes	yes	yes	yes
F. coliforms	gmean $\leq 100$ cfu	yes	yes	yes	yes	yes	yes
	max. 10% $> 200$ cfu	yes	yes	yes	no	yes	yes

	Typical range	Blue Canyon	Carpenter	Euclid	Mill- wheel	Park Place	Silver Beach
Alkalinity	med. $\leq 30$ mg/L	no	yes	yes	yes	no	no
Conductivity	med. $\leq 100$ $\mu$ S	no	yes	yes	yes	no	no
D. oxygen <sup>†</sup>	min. $\geq 8.0$ mg/L	yes	yes	yes	no	no	yes
pH	6.5–8.5	yes	yes	yes	yes	yes	yes
Temperature <sup>†</sup>	max. $\leq 17.5$ C	yes	yes	yes	yes	yes	yes
T. susp. solids	med. $\leq 5$ mg/L	yes	yes	yes	no	yes	yes
Turbidity	med. $\leq 5$ NTU	yes	yes	yes	no	yes	yes
Ammonium	med. $\leq 10$ $\mu$ g-N/L	yes	yes	yes	no	yes	yes
Sol. phosphate	med. $\leq 10$ $\mu$ g-P/L	yes	yes	yes	yes	no	no
T. phosphorus	med. $\leq 20$ $\mu$ g-P/L	yes	yes	yes	no	no	no
F. coliforms	gmean $\leq 100$ cfu	yes	yes	yes	no	no	no
	Max. 10% $> 200$ cfu	yes	no	no	no	no	no

<sup>†</sup>Many of the residential creeks were not sampled during part of the summer due to low flow, which is when water temperatures are usually high and dissolved oxygen concentrations low.

Table 3.3: Comparison of October 2020-September 2021 water quality in Lake Whatcom tributaries (“no” indicates that the site does not fall within the water quality ranges or meet the criteria described on page 57).

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	12.9	17.8	18.8	26.3
Conductivity (μS/cm)	47.4	56.7	57.5	67.8
Dissolved oxygen (mg/L)	7.6	10.4	10.3	12.1
pH	6.7	6.9	6.9	7.2
Temperature (°C)	5.3	9.2	9.3	14.1
Total suspended solids (mg/L)	<2	<2	<2	3.7
Turbidity (NTU)	0.3	1.1	1.1	2.3
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	22.7
Nitrogen, nitrate/nitrite (μg-N/L)	308.2	571.8	556.3	799.5
Nitrogen, total (μg-N/L)	460.1	635.8	661.3	988.6
Phosphorus, soluble (μg-P/L)	<5	7.3	7.1	16.0
Phosphorus, total (μg-P/L)	6.7	12.7	15.7	31.0
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 8)	2	28	28	270

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.4: Summary of Anderson Creek water quality data, October 2020-September 2021.



Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	10.8	17.5	22.6	39.4
Conductivity (μS/cm)	42.1	62.6	82.9	144.2
Dissolved oxygen (mg/L)	9.4	11.7	11.4	13.1
pH	7.0	7.4	7.5	7.7
Temperature (°C)	4.8	8.6	9.7	16.8
Total suspended solids (mg/L)	<2	<2	6.0	33.2
Turbidity (NTU)	0.4	1.4	2.6	11.2
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	21.7
Nitrogen, nitrate/nitrite (μg-N/L)	106.6	676.5	684.5	1390.8
Nitrogen, total (μg-N/L)	171.3	712	759.1	1653.1
Phosphorus, soluble (μg-P/L)	<5	5.7	5.2	9.3
Phosphorus, total (μg-P/L)	8.4	11.6	13	24.5
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 8)	4	99	46	244

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.5: Summary of Austin Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	53.5	143.3	139	186.2
Conductivity (μS/cm)	142.1	308	296.7	362
Dissolved oxygen (mg/L)	11.0	11.9	11.9	12.9
pH	7.9	8.3	8.3	8.5
Temperature (°C)	5.7	8.0	8.0	11.3
Total suspended solids (mg/L)	<2	3.1	3.7	10.0
Turbidity (NTU)	0.7	1.6	1.9	5.5
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	<10
Nitrogen, nitrate/nitrite (μg-N/L)	280.6	780.3	1071.8	2680.2
Nitrogen, total (μg-N/L)	283.9	732.7	982.3	1907.4
Phosphorus, soluble (μg-P/L)	<5	6.0	<5	11.1
Phosphorus, total (μg-P/L)	<5	9.6	10.2	14.9
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 0)	<1	1	2	130

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.6: Summary of Blue Canyon Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	6.2	9.1	10.2	17.1
Conductivity (μS/cm)	34.6	41.8	42.6	52.2
Dissolved oxygen (mg/L)	9.3	11.8	11.5	12.7
pH	6.8	7.0	7.0	7.1
Temperature (°C)	4.9	7.3	7.7	11.4
Total suspended solids (mg/L)	<2	<2	2.7	12.0
Turbidity (NTU)	0.6	1	1.6	5.2
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	<10
Nitrogen, nitrate/nitrite (μg-N/L)	250.5	887.5	976.6	1744.7
Nitrogen, total (μg-N/L)	320.8	1128.3	1065.7	1944.9
Phosphorus, soluble (μg-P/L)	<5	<5	<5	6.2
Phosphorus, total (μg-P/L)	5.1	7.3	8.2	12.9
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 0)	1	7	6	64

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.7: Summary of Brannian Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	13.3	22.7	24.9	45.1
Conductivity (μS/cm)	52.5	69.6	74.1	107.5
Dissolved oxygen (mg/L)	10.7	12.1	11.9	12.7
pH	7.2	7.6	7.5	7.9
Temperature (°C)	5.7	7.2	8.1	11.9
Total suspended solids (mg/L)	<2	2.8	4.8	14.5
Turbidity (NTU)	0.7	3.9	3.9	8.4
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	21.3
Nitrogen, nitrate/nitrite (μg-N/L)	293.5	811.5	896.7	1398.1
Nitrogen, total (μg-N/L)	415.6	1186.5	1111.5	1890.3
Phosphorus, soluble (μg-P/L)	<5	7.0	6.0	11.1
Phosphorus, total (μg-P/L)	6.9	13.5	18.0	37.0
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 11)	4	22	28	460

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.8: Summary of Carpenter Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.4	27	30.6	46.7
Conductivity (μS/cm)	38.2	87.4	88	128.8
Dissolved oxygen (mg/L)	8.5	11.3	11.0	12.3
pH	7.1	7.2	7.2	7.3
Temperature (°C)	6.6	8.2	9.0	12.8
Total suspended solids (mg/L)	<2	<2	2.8	7.2
Turbidity (NTU)	0.8	1.4	2.2	4.7
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	10.0
Nitrogen, nitrate/nitrite (μg-N/L)	108.0	475.0	449.1	905.3
Nitrogen, total (μg-N/L)	174.6	689.3	580.3	1124.6
Phosphorus, soluble (μg-P/L)	<5	6.5	5.6	9.6
Phosphorus, total (μg-P/L)	7.2	14.8	15.2	22.6
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 11)	1	32	20	230

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.9: Summary of Euclid Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	16.4	28.3	34.2	64.1
Conductivity (μS/cm)	58.1	91.4	95.2	145.6
Dissolved oxygen (mg/L)	4.6	11.0	9.9	12.2
pH	7.0	7.2	7.2	7.5
Temperature (°C)	6.5	7.9	10.4	18.2
Total suspended solids (mg/L)	2.8	7.3	6.9	11.8
Turbidity (NTU)	5.4	7.6	7.8	10.4
Nitrogen, ammonium (μg-N/L)	<10	12.1	25.9	107.2
Nitrogen, nitrate/nitrite (μg-N/L)	<20	433.0	555.5	1184.0
Nitrogen, total (μg-N/L)	350.0	1013.0	897.2	1526.6
Phosphorus, soluble (μg-P/L)	<5	9.5	11.3	35.8
Phosphorus, total (μg-P/L)	25.5	36.7	49.9	119.5
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 22)	40	92	127	2500

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.10: Summary of Millwheel Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	9.5	19.2	26.7	50.6
Conductivity (μS/cm)	36.7	61.2	76.2	130.4
Dissolved oxygen (mg/L)	9.6	11.9	11.5	12.8
pH	7.1	7.5	7.6	8.0
Temperature (°C)	5.3	8.0	9.5	16.6
Total suspended solids (mg/L)	<2	3.3	13.0	79.0
Turbidity (NTU)	0.3	1.5	5.1	26.7
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	10.2
Nitrogen, nitrate/nitrite (μg-N/L)	300.6	651.1	789.6	1694.8
Nitrogen, total (μg-N/L)	164.2	746.3	925.4	2027.3
Phosphorus, soluble (μg-P/L)	<5	6.0	6.3	13.5
Phosphorus, total (μg-P/L)	7.9	12.9	17.6	46.6
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 25)	<1	54	40	2000

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.11: Summary of Olsen Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	50.2	80.3	77.9	104.3
Conductivity (μS/cm)	149.9	231.6	216.6	268.5
Dissolved oxygen (mg/L)	0.3	10.6	8.7	11.9
pH	7.2	7.7	7.7	8.1
Temperature (°C)	7.4	11.4	12.2	18.7
Total suspended solids (mg/L)	<2	2.0	2.2	5.7
Turbidity (NTU)	1.7	3.0	3.2	5.1
Nitrogen, ammonium (μg-N/L)	<10	<10	10.6	22.4
Nitrogen, nitrate/nitrite (μg-N/L)	434.7	664.9	762.4	1148.4
Nitrogen, total (μg-N/L)	539.7	708.4	881.9	1476.2
Phosphorus, soluble (μg-P/L)	<5	13.9	13.9	33.6
Phosphorus, total (μg-P/L)	15.9	23.6	24.9	33.6
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 33)	7	91	116	1200

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.12: Summary of Park Place outlet water quality data, October 2020-September 2021.



Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	35.6	57.4	64.6	113.1
Conductivity (μS/cm)	100.8	165.6	166.8	263.6
Dissolved oxygen (mg/L)	10.4	11.8	11.6	12.5
pH	7.5	7.8	7.8	8.2
Temperature (°C)	6.1	8.0	8.8	12.1
Total suspended solids (mg/L)	<2	3.2	2.9	4.9
Turbidity (NTU)	1.5	2.4	3.1	5.6
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	15.2
Nitrogen, nitrate/nitrite (μg-N/L)	230.8	536.6	650.2	1374.9
Nitrogen, total (μg-N/L)	412.4	643.5	901	1819.1
Phosphorus, soluble (μg-P/L)	<5	13.3	11.8	19.2
Phosphorus, total (μg-P/L)	17.8	26.3	25.5	33.8
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 50)	3	165	122	2100

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.13: Summary of Silver Beach Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	11.1	16.7	21.3	36.6
Conductivity (μS/cm)	42.9	59	67.2	105.9
Dissolved oxygen (mg/L)	9.9	12.1	11.8	13.2
pH	7.2	7.5	7.5	7.8
Temperature (°C)	5.0	7.9	9.2	15.7
Total suspended solids (mg/L)	<2	2.9	5.1	18.8
Turbidity (NTU)	0.1	1.3	2.0	7.0
Nitrogen, ammonium (μg-N/L)	<10	<10	<10	15.5
Nitrogen, nitrate/nitrite (μg-N/L)	359.2	1033.6	1090.2	2095.5
Nitrogen, total (μg-N/L)	251.1	1166.6	1189	2403.9
Phosphorus, soluble (μg-P/L)	<5	<5	<5	7.5
Phosphorus, total (μg-P/L)	6.0	8.2	10.3	18.9
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 0)	<1	7	8	110

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3.14: Summary of Smith Creek water quality data, October 2020-September 2021.

Variable	Min.	Med. <sup>†</sup>	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	19.4	21.7	21.7	24.2
Conductivity ( $\mu$ S/cm)	57.6	61	61.3	64.8
Dissolved oxygen (mg/L)	8.6	10.8	10.6	12.3
pH	7.4	7.6	7.6	7.9
Temperature ( $^{\circ}$ C)	6.6	14.5	14.5	26.2
Total suspended solids (mg/L)	<2	<2	2.0	5.5
Turbidity (NTU)	0.4	0.8	0.8	1.2
Nitrogen, ammonium ( $\mu$ g-N/L)	<10	<10	<10	34.5
Nitrogen, nitrate/nitrite ( $\mu$ g-N/L)	<20	100.3	121.6	292.5
Nitrogen, total ( $\mu$ g-N/L)	<100	278.5	272.3	482.4
Phosphorus, soluble ( $\mu$ g-P/L)	<5	<5	<5	6.4
Phosphorus, total ( $\mu$ g-P/L)	6.5	9.3	10	14.9
Coliforms, fecal (cfu/100 mL) <sup>‡</sup> (Percent of samples >200 cfu/100 mL = 0)	<1	5	6	96

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1  $\Rightarrow$  1).

Table 3.15: Summary of Whatcom Creek water quality data, October 2020-September 2021.

Site	Date	TOC-AM (mg/L)	TOC-IWS (mg/L)	Date	TOC-AM (mg/L)	TOC-IWS (mg/L)
Anderson	Feb 3, 2021	2.4	2.2	Jul 13, 2021	1.5	1.2
Austin (lower)	Feb 3, 2021	2.7	2.3	Jul 13, 2021	2.2	1.7
Blue Canyon	Feb 3, 2021	3.8	3.6	Jul 13, 2021	insufficient flow	
Brannian	Feb 3, 2021	2.2	1.9	Jul 13, 2021	insufficient flow	
Carpenter	Feb 3, 2021	5.0	4.8	Jul 13, 2021	insufficient flow	
Euclid	Feb 3, 2021	4.1	3.9	Jul 13, 2021	insufficient flow	
Millwheel	Feb 3, 2021	4.4	4.3	Jul 13, 2021	insufficient flow	
Olsen	Feb 3, 2021	3.8	3.5	Jul 13, 2021	2.7	2.3
Park Place	Feb 3, 2021	5.3	4.7	Jul 13, 2021	3.0	2.5
Silver Beach	Feb 3, 2021	5.4	5.9	Jul 13, 2021	insufficient flow	
Smith	Feb 3, 2021	4.3	3.0	Jul 13, 2021	2.1	2.0
Whatcom	Feb 3, 2021	2.2	1.9	Jul 13, 2021	3.4	2.6

Table 3.16: Lake Whatcom 2021 tributary total organic carbon data. February and July samples were split and analyzed by AmTest (TOC-AM) and IWS (TOC-IWS).

## 4 Storm Water Monitoring

### 4.1 Hydrograph Monitoring

### 4.2 Field Sampling and Analytical Methods

Recording hydrographs are installed in Austin Creek and Smith Creek; the data are plotted in Figures 4.1–4.2 (pages 82–83). The location of each hydrograph is described in Appendix A.2 (page 94). All hydrograph data, including data from previous years, are online at [www.wvu.edu/iws](http://www.wvu.edu/iws). Field notes, comments on missing data, and rating curves for each water year are available upon request to the City of Bellingham or Institute for Watershed Studies. The rating curves were generated using Aquarius rating curve software (Aquatic Informatics, 2014; <https://aquaticinformatics.com>). The field discharge and stage height measurements are plotted in Figures 4.3–4.4 (pages 84–85) and the Aquarius rating curve equations are listed in Tables 4.1–4.2 (pages 79–80). All results are reported as Pacific Standard Time, without Daylight Saving Time adjustment.

### 4.3 Site Descriptions

The 2020/2021 storm water sampling focused on Austin, Euclid, and Silver Beach Creeks (Figure A2, page 99). Earlier storm water sampling in the Lake Whatcom watershed summarized in previous annual reports (see Section 5.2, page 90).

### 4.4 Field Sampling and Analytical Methods

Two storm events were sampled in Austin Creek, three in Euclid Creek, and four in Silver Beach Creek (Table 4.3, page 81). The samples were collected using time-paced ISCO samplers provided by the City of Bellingham and analyzed for total suspended solids, total phosphorus, soluble reactive phosphorus, total nitrogen, and nitrate/nitrite<sup>29</sup> as described in Table 2.1 (page 18).

---

<sup>29</sup>Nitrate and nitrite were analyzed together because nitrite concentrations are very low in surface water and require low level analytical techniques to measure accurately. For simplicity, nitrate/nitrite will be referred to as “nitrate” in this document.

Dry summers have a direct impact on stream flow (base flow), which is supported by soil water and groundwater. As illustrated in the hydrographs for the Lake Whatcom watershed (Figures 4.1–4.2, pages 82–83), stream discharge decreased over the course of the summer as soils dried out and groundwater levels declined due to low rainfall and high levels of evapotranspiration from vegetation. Moreover, most late summer rainfall goes into replenishing soil water (storage) rather than direct runoff into streams. Lower summer stream flows and groundwater levels will also reduce runoff into the lake. When coupled with higher summer lake withdrawals and lake evaporation, the lake level will usually drop over the course of the summer, reaching a minimum in late fall.

As indicated in Table 3.2 (page 62), many of the smaller tributaries to Lake Whatcom were dry, or nearly dry during late summer. Storm water data are used by the City as part of their watershed modeling program and will be reported directly to the City to be incorporated into the model. Storm water data are available upon request to the City of Bellingham or the Institute for Watershed Studies.

Stage Height (ft)	Discharge Equations
0.23–0.31	discharge = $175.741 \times \text{stage}^{4.808}$
0.31–0.60	discharge = $8.932 \times \text{stage}^{2.264}$
0.60–0.95	discharge = $11.907 \times \text{stage}^{2.827}$
0.95–1.41	discharge = $11.952 \times \text{stage}^{2.897}$
1.41–2.28	discharge = $13.526 \times \text{stage}^{2.537}$
2.28–3.61	discharge = $18.155 \times \text{stage}^{2.180}$

Table 4.1: Austin Creek rating curves for WY2021 (October 1, 2020-September 30, 2021); equations generated by Aquarius software (Aquatic Informatics, 2014).

Stage Height (ft)	Discharge Equations
1.58–1.63	discharge = $1.2^{-27} \times \text{stage}^{125.566}$
1.63–2.24	discharge = $0.010 \times \text{stage}^{7.920}$
2.24–3.08	discharge = $0.034 \times \text{stage}^{6.460}$
3.08–3.89	discharge = $0.344 \times \text{stage}^{4.398}$
3.89–4.67	discharge = $1.458 \times \text{stage}^{3.336}$

Table 4.2: Smith Creek rating curves for WY2021 (October 1, 2020-September 30, 2021); equations generated by Aquarius software (Aquatic Informatics, 2014).



<i>Austin Creek</i>			
Start Date	Start Time	End Date	End time
Dec 21, 2020	07:00	Dec 21, 2020	13:00
Jan 2, 2021	14:00	Jan 4, 2021	08:00

<i>Euclid Creek</i>			
Start Date	Start Time	End Date	End time
Nov 12, 2020	18:00	Nov 13, 2020	13:15
Dec 8, 2020	03:00	Dec 9, 2020	10:00
Feb 21, 2021	20:30	Feb 22, 2021	10:00

<i>Silver Beach Creek</i>			
Start Date	Start Time	End Date	End time
Nov 3, 2020	09:15	Nov 4, 2020	07:45
Nov 17, 2020	19:45	Nov 18, 2020	05:45
Dec 8, 2020	00:30	Dec 9, 2020	10:30
Feb 21, 2021	20:30	Feb 22, 2021	10:00

Table 4.3: Summary of 2020-2021 storm event sampling dates for Austin, Euclid, and Silver Beach Creeks.




Figure 4.1: Austin Creek hydrograph for WY2021 (October 1, 2020–September 30, 2021). Data were recorded at 15 minute intervals.




Figure 4.2: Smith Creek hydrograph for WY2021 (October 1, 2020–September 30, 2021). Data were recorded at 15 minute intervals.




Figure 4.3: Aquarius rating curve for Austin Creek. The triangles (▲) show the fitted rating curve for WY2021. The circles show data collected in WY2021 (●), WY2020 (●), WY2019 (●). The red circles (●) show data collected prior to WY2019 that were used for the rating curve because they help define the high end of the rating curve.




Figure 4.4: Aquarius rating curve for Smith Creek. The triangles (▲) show the fitted rating curve for WY2021. The circles show data collected in WY2021 (●), WY2020 (●), WY2019 (●). The red circles (●) show data collected prior to WY2019 that were used for the rating curve because they help define the high end of the rating curve.

## 5 References and Related Reports

### 5.1 Cited References

- Aquatic Informatics. 2014. AQUARIUS rating curve software. Aquatic Informatics, Vancouver, British Columbia, Canada.
- APHA. 2017. Standard Methods for the Examination of Water and Wastewater, 23rd Edition. American Public Health Association, American Water Works Association, and Water Environment Federation, Washington, DC.
- Ashurst, S. 2003. Microcosm study of the accumulation of benzo(a)pyrene by Lake Whatcom phytoplankton. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.
- Baribeau, H., S. Krasner, R. Chinn, and P. Singer. 2005. Impact of biomass on the stability of HAAs and THMs in a simulated distribution system. *Journal–American Water Works Association* 97:69–81
- Bittner, C. W. 1993. The response of Lake Whatcom bacterioplankton to nutrient enrichment. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.
- Deacon, J. 2015. Determining biologically available phosphorus in storm water entering Lake Whatcom, WA using the dual culture diffusion apparatus. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.
- Fee, E. 1976. The vertical and seasonal distribution of chlorophyll in lakes of the Experimental Lakes Area, northwestern Ontario: Implications for primary production estimates. *Limnology and Oceanography* 21: 767-783.
- Grigorescu, A., R. Hozalski. 2010. Modeling HAA biodegradation in biofilters and distribution systems. *Journal–American Water Works Association* 102: 67–80.
- Groce, S. 2011. Soils as a Source of Bioavailable Phosphorus in the Lake Whatcom Watershed. M. S. thesis, Huxley College of the Environment, Western Washington University, Bellingham, WA.

- Johengen, T., G. Smith, D. Schar, H. Purcel, D. Loewensteiner, Z. Epperson, M. Tamburri, G. Meadows, S. Green, F. Yousef, J. Anderson. 2016. Performance verification statement for Hach Hydrolab DS5X and HL4 dissolved oxygen sensor. Alliance for Coastal Technologies, Alliance for Coastal Technologies, ACT VS16-05. DOI: <http://dx.doi.org/10.25607/OBP-298>.
- Liang, C-W. 1994. Impact of Soil and Phosphorus Enrichment on Lake Whatcom Periphytic Algae. M. S. thesis, Huxley College of the Environment, Western Washington University, Bellingham, WA.
- Lind, O. T. 1985. Handbook of Common Methods in Limnology, 2nd Edition. Kendall/Hunt Publishing Co., Dubuque, IA.
- Matthews, R. A. and E. DeLuna, 2008. Metalimnetic oxygen and ammonium maxima in Lake Whatcom, Washington (USA). Northwest Science 82:18–29.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, K. Beeler, and G. B. Matthews. 2015. Lake Whatcom Monitoring Project, 2013/2014 Final Report, February 26, 2015. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2012. Lake Whatcom Monitoring Project, 2010/2011 Final Report, February 24, 2012. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2008. Lake Whatcom Monitoring Project 2006–2007 Final Report. Final Report prepared for the City of Bellingham Public Works Department, April, 2008, Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2005. Lake Whatcom Monitoring Project 2003–2004 Final Report. Final Report prepared for the City of Bellingham Public Works Department, March, 2005, Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2004. Lake Whatcom Monitoring Project 2002–2003 Final Report. Final Report prepared for the City of Bellingham Public Works Department, March, 2004, Bellingham, WA.

- Matthews, R., M. Hilles, and G. Pelletier. 2002a. Determining trophic state in Lake Whatcom, Washington (USA), a soft water lake exhibiting seasonal nitrogen limitation. *Hydrobiologia* 468:107–121.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2002b. Lake Whatcom Monitoring Project 2000–2001 Final Report. Final Report prepared for the City of Bellingham Public Works Department, March, 2002, Bellingham, WA.
- McDonald, K. R. 1994. Nutrient limitation of phytoplankton in Lake Whatcom. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.
- Mitchell, R., G. Gabrisch, and R. Matthews. 2010. Lake Whatcom Bathymetry and Morphology. Report prepared for the City of Bellingham Public Works Department, December 2, 2010, Bellingham, WA.
- Oliver, S. K., S. M. Colling, P. A. Soranno, T. Wagner, E. H. Stanley, J. R. Jones, C. A. Stow, and N. R. Lottig. 2017. Unexpected stasis in a changing world: lake nutrient and chlorophyll trends since 1990. *Global Change Biology* 23(12): 5455–5467.
- Pelletier, G. 1998. Dissolved oxygen in Lake Whatcom. Trend in the depletion of hypolimnetic oxygen in basin I, 1983–1997. Washington State Department of Ecology Report #98–313, Olympia, WA.
- Potter, B. and F. Wimsatt. 2009. Method 415.3, Rev. 1.2: Determination of Total Organic Carbon and Specific UV Absorbance at 254 nm in Source Water and Drinking Water. U. S. Environmental Protection Agency, Washington, DC.
- Rantz, S.E., et al. (1982). Measurement and Computation of Streamflow: Volume 1. Measurement of Stage and Discharge. Geological Survey Water-Supply Paper #2175, U. S. Government Printing Office, Washington, D. C.
- Rogora, M., S. Arisci, and A. Marchetto. 2012. The role of nitrogen deposition in the recent nitrate decline in lakes and rivers in Northern Italy. *Science of the Total Environment* 417–418: 214–223.



- Sickman, J. O., J. M. Melack, and D. W. Clow. 2003. Evidence for nutrient enrichment of high-elevation lakes in the Sierra Nevada, California. *Limnol. Oceanogr.* 48(5):1885–1892
- Sung, W., B. Reilly-Matthews, D. K. O’Day, and K. Horrigan. 2000. Modeling DBP Formation. *J. Amer. Water Works Assoc.* 92:5–53.
- Waller, K., D. Driscoll, J. Lynch, D. Newcomb, and K. Roy. 2012. Long-term recovery of lakes in the Adirondack region of New York to decreases in acid deposition. *Atmospheric Environment* 46:56–64.
- Wetzel, R. G. 2001. *Limnology, Third Edition*. Academic Press, San Diego, CA.
- WOW. 2004. Water on the Web - Monitoring Minnesota Lakes on the Internet and Training Water Science Technicians for the Future - A National On-line Curriculum using Advanced Technologies and Real-Time Data (<http://WaterOnTheWeb.org>). Authors: Munson, B., R. Axler, C. Hagley, G. Host, G. Merrick, and C. Richards. University of Minnesota-Duluth, Duluth, MN 55812.
- YSI. 2017. EXO User Manual, Revision G, April 2017. YSI Incorporated, Yellow Springs, OH.

## 5.2 Related Reports

The following is a list of annual reports and special project reports produced by the Institute for Watershed Studies since 1987 as part of the Lake Whatcom monitoring program sponsored by the City of Bellingham and Western Washington University. Many of the reports are available online at [www.wwu.edu/iws](http://www.wwu.edu/iws) (follow links to the Lake Whatcom project under Lake Studies); older reports are available in the IWS library and through the city of Bellingham Public Works Department. This list does not include research reports, student projects, or publications that were not prepared specifically for the City of Bellingham. Contact IWS for information about additional Lake Whatcom publications.

### Annual monitoring reports:

- Matthews, R. A., A. Strecker, M. Hilles, J. Pickens, R. Mitchell, and G. B. Matthews. 2020. Lake Whatcom Monitoring Project, 2018/2019 Final Report, February 6, 2020. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Pickens, R. Mitchell, and G. B. Matthews. 2019. Lake Whatcom Monitoring Project, 2017/2018 Final Report, February 26, 2019. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Pickens, R. Mitchell, K. Beeler, and G. B. Matthews. 2018. Lake Whatcom Monitoring Project, 2016/2017 Final Report, February 23, 2018. Report to the City of Bellingham, WA. 0
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, K. Beeler, and G. B. Matthews. 2017. Lake Whatcom Monitoring Project, 2015/2016 Final Report, February 21, 2017. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, K. Beeler, and G. B. Matthews. 2016. Lake Whatcom Monitoring Project, 2014/2015 Final Report, February 23, 2016. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, K. Beeler, and G. B. Matthews. 2015. Lake Whatcom Monitoring Project, 2013/2014 Final Report, February 26, 2015. Report to the City of Bellingham, WA.

- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2014. Lake Whatcom Monitoring Project, 2012/2013 Final Report, March 6, 2014. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2013. Lake Whatcom Monitoring Project, 2011/2012 Final Report, March 8, 2013. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2012. Lake Whatcom Monitoring Project, 2010/2011 Final Report, February 24, 2012. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2009/2010 Final Report, March 1, 2011. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2008/2009 Final Report, March 10, 2010. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2007/2008 Final Report, March 19, 2009. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2006/2007 Final Report, April 2, 2008. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2005/2006 Final Report, April 11, 2007. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2004/2005 Final Report, March 30, 2006. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2003/2004 Final Report, March 15, 2005. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2002/2003 Final Report, April 5, 2004. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2001/2002 Final Report, April 21, 2003. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2000/2001 Final Report, March 15, 2002. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 1999/2000 Final Report, March 23, 2001. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 1998/99 Final Report, March 15, 2000. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1997/98 Final Report, April 12, 1999. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1996/97 Final Report, February 10, 1998. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1995/96 Final Report, March 24, 1997. Report to the City of Bellingham, WA.

Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1994/95 Final Report, February 9, 1996. Report to the City of Bellingham, WA.

Matthews, R. A. and G. B. Matthews. Lake Whatcom Monitoring Project, 1993–1994 Final Report, March 2, 1995. Report to the City of Bellingham, WA.

Matthews, R. and G. Matthews. Lake Whatcom Monitoring Project, 1992–1993 Final Report, January 31, 1994. Report to the City of Bellingham, WA.

Matthews, R. and G. Matthews. Lake Whatcom Monitoring Project, 1991–1992 Final Report, March 19, 1993. Report to the City of Bellingham, WA.

Rector, J. M. and R. A. Matthews. Lake Whatcom Monitoring Program, August 1987 Final Report. Institute for Watershed Studies Report, Western Washington University, Bellingham, WA.

Strecker, A., R. A. Matthews, M. Hilles, J. Pickens, R. Mitchell, and G. B. Matthews. 2021. Lake Whatcom Monitoring Project, 2019/2020 Final Report, February 24, 2021. Report to the City of Bellingham, WA.

**Other Lake Whatcom reports:**

Matthews, R. A., M. Hilles and J. Vandersypen. Austin Creek and Beaver Creek Sampling Project, October 11, 2005. Report to the City of Bellingham, WA.

Matthews, R. A. Relationship between Drinking Water Treatment Chemical Usage and Lake Whatcom water Quality and Algal Data, October 4, 2004. Report to the City of Bellingham, WA.

Matthews, R. A. Strawberry Sill Water Quality Analysis, March 19, 2004. Report to the City of Bellingham, WA.

Matthews, R. A., M. Saunders, M A. Hilles, and J. Vandersypen. Park Place Wet Pond Monitoring Project, 1994–2000 Summary Report, February 2, 2001. Report to the City of Bellingham, WA.

Carpenter, M. R., C. A. Suczek, and R. A. Matthews. Mirror Lake Sedimentation Study Summary Report, February, 1992. Report to the City of Bellingham, WA.

Walker, S., R. Matthews, and G. Matthews. Lake Whatcom Storm Runoff Project, Final Report, January 13, 1992. Report to the City of Bellingham, WA.

Creahan, K., T. Loranger, B. Gall, D. Brakke, and R. Matthews. Lake Whatcom Watershed Management Plan, December, 1986, revised July, 1987. Institute for Watershed Studies Report, Western Washington University, Bellingham, WA.

## A Site Descriptions

Figures A1–A2 (pages 98–99) show the locations of the current monitoring sites and Table A1 (page 97) lists the approximate GPS coordinates for the lake and creek sites. All site descriptions, including text descriptions and GPS coordinates, are approximate. For detailed information about sampling locations, contact IWS.

### A.1 Lake Whatcom Monitoring Sites

**Site 1** is located in the north central portion of basin 1 along a straight line from the Bloedel Donovan boat launch to the house located at 171 E. North Shore Rd. The depth at Site 1 should be at least 25 meters; samples are collected from the surface to 20 m.

**Site 2** is located in the south central portion of basin 2 just west of the intersection of a line joining the boat house at 73 Strawberry Point and the point of Geneva sill. Samples are collected from the surface to 20 m.

The **Intake Site** location is omitted from this report at the City’s request.

**Site 3** is located in the northern portion of basin 3, mid-basin just north of a line between the old railroad bridge and Lakewood. The depth at Site 3 should be at least 80 m; samples are collected from the surface to 75 m.

**Site 4** is located in the southern portion of basin 3, mid-basin, and just north of South Bay. The depth at Site 4 should be at least 90 m; samples are collected from the surface to 90 m.

### A.2 Tributary Monitoring Sites

**Anderson Creek** samples are collected using a sampling pole from the downstream side of the South Bay Rd. bridge. The Anderson Creek hydrograph<sup>30</sup> is mounted in the stilling well on the east side of Anderson Creek, directly adjacent

---

<sup>30</sup>This hydrograph is no longer maintained by IWS; data are available on the USGS web site at [http://waterdata.usgs.gov/nwis/inventory?agency\\_code=USGS&site\\_no=12201950](http://waterdata.usgs.gov/nwis/inventory?agency_code=USGS&site_no=12201950).

to the bridge over Anderson Creek (South Bay Rd.), approximately 0.5 km from the mouth of the creek.

The **Austin Creek** hydrograph gauge and sampling site is located approximately 15 m downstream from Lake Whatcom Blvd. From October 2004 through September 2006, three additional sampling sites were sampled in the Austin Creek watershed, so for clarification, the gauged site has been renamed **Lower Austin Creek**.

**Blue Canyon Creek** samples are collected downstream from the culvert under Blue Canyon Rd. in the second of three small streams that cross the road. During conditions of low flow or high lake levels, samples are collected, if possible, approximately 7 m upstream from the road crossing.

**Brannian Creek** samples are collected using a sampling pole from the downstream side of South Bay Rd., approximately 40 m upstream from the USGS hydrograph gauge. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Carpenter Creek** samples are collected approximately 7 m upstream from North Shore Dr. near the USGS hydrograph gauge. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Euclid Ave.** samples are collected from a small tributary off Euclid Avenue near the USGS hydrograph gauge. The site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Millwheel Creek** samples are collected approximately 8 m upstream from Flynn St. near the USGS hydrograph gauge. The creek is unnamed on most topographic maps, but has been called “Millwheel Creek” by residents of the watershed due to its proximity to the old mill pond. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Olsen Creek** samples are collected upstream from North Shore Dr., approximately 3 meters upstream from the bridge. This site was added in October 2004 as part of the 2004–2006 monthly creek monitoring project.

**Park Place** samples are collected from the storm drain that empties into Lake Whatcom at Park Place Ln. Samples from this site include outlet flow from the Park Place storm water treatment facility.

**Silver Beach Creek** samples are collected approximately 75 m upstream from the culvert under North Shore Rd., just upstream from the USGS hydrograph gauge.

The **Smith Creek** hydrograph is mounted on the south wall of a sandstone bluff directly underneath the bridge over Smith Creek (North Shore Rd.) approximately 1 km upstream from the mouth of the creek. Water samples are collected near the old bridge site, at the end of North Shore Rd. approximately 500 m downstream from the IWS hydrograph gauge. During periods of low flow, Smith Creek is sampled approximately 15 m downstream from the IWS hydrograph gauge.

**Whatcom Creek** samples are collected approximately 2 m downstream from the foot bridge below the Lake Whatcom outlet spillway. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

### **A.3 Storm Water Monitoring Sites**

The 2020/2021 storm water monitoring program focused on collecting storm runoff data from Austin, Euclid, and Silver Beach Creeks. Austin Creek samples are collected approximately 15 m downstream from Lake Whatcom Blvd. Euclid Creek samples are collected from a small tributary located off Euclid Avenue near the USGS hydrograph gauge. Silver Beach Creek samples are collected approximately 75 m upstream from the culvert under North Shore Rd., just upstream from the USGS hydrograph gauge. For information about other storm water sites that have been monitored by IWS, refer to the annual reports listed in Section [5.2](#) (page [90](#)).



Lake Sites	Latitude (°N)	Longitude (°W)
Site 1	48.760	-122.411
Intake	(GPS omitted)	
Site 2	48.743	-122.382
Site 3	48.738	-122.336
Site 4	48.695	-122.304

Tributary/Stormwater Sites	Latitude (°N)	Longitude (°W)
Anderson	48.673	-122.268
Austin (lower)	48.713	-122.331
Blue Canyon	48.685	-122.283
Brannian	48.669	-122.279
Carpenter	48.754	-122.354
Euclid	48.748	-122.410
Millwheel	48.755	-122.416
Olsen	48.751	-122.354
Park Place	48.769	-122.409
Silver Beach	48.769	-122.407
Smith	48.732	-122.309
Whatcom	48.757	-122.422

Table A1: Approximate GPS coordinates for Lake Whatcom sampling sites.




Figure A1: Lake Whatcom lake sampling sites. Basemap created using data from Western Washington University, Skagit County, the Nooksack Tribe, and the City of Bellingham.




Figure A2: Lake Whatcom tributary and storm water sampling sites. Basemap created using data from Western Washington University, Skagit County, the Nooksack Tribe, and the City of Bellingham.

## **B Long-Term Water Quality Figures**

The current and historic Lake Whatcom water quality data are plotted on the following pages. Detection limits and abbreviations for each parameter are listed in Table 2.1 (page 18).

The historic detection limits for each parameter were estimated based on an analysis of historic detection ranges, instrument limitations, and analyst judgment on the lowest repeatable concentration for each test. Over time, some analytical techniques have improved so that current detection limits are lower than the historic limits listed in Table 2.1, page 18). Because the Lake Whatcom data set includes long-term monitoring data that have been collected using a variety of analytical techniques, this report sets conservative historic detection limits to allow comparisons between all years.

In the Lake Whatcom report, unless indicated, no data substitutions are used for below detection values (“bdl” data). Instead, we identify summary statistics that include bdl values, and, if appropriate, discuss the implications of including these values in the analysis.

Because of the length of the data record, many of the figures reflect trends related to improvements in analytical techniques over time and the introduction of increasingly sensitive field equipment (see, for example, Figures B81–B85, pages 183–187, which show the effect of using increasingly sensitive conductivity probes). These changes generally result in a reduction in analytical variability, and sometimes result in lower detection limits.

## **B.1 Monthly YSI Profiles**




Figure B1: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, October 7, 2020.




Figure B2: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, October 7, 2020.




Figure B3: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, October 7, 2020.






Figure B4: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, October 5, 2020.




Figure B5: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, October 5, 2020.




Figure B6: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, November 9, 2020.




Figure B7: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, November 9, 2020.




Figure B8: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, November 9, 2020.




Figure B9: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, November 2, 2020.




Figure B10: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, November 2, 2020.




Figure B11: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, December 9, 2020.






Figure B12: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, December 9, 2020.




Figure B13: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, December 9, 2020.




Figure B14: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, December 2, 2020.




Figure B15: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, December 2, 2020.




Figure B16: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, February 27, 2021.




Figure B17: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, February 24, 2021.




Figure B18: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, February 27, 2021.




Figure B19: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, February 17, 2021.






Figure B20: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, February 17, 2021.




Figure B21: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, April 8, 2021.




Figure B22: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, April 8, 2021.




Figure B23: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, April 8, 2021.




Figure B24: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, April 6, 2021.




Figure B25: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, April 6, 2021.




Figure B26: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, May 10, 2021.




Figure B27: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, May 6, 2021.






Figure B28: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, May 10, 2021.




Figure B29: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, May 4, 2021.




Figure B30: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, May 4, 2021.




Figure B31: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, June 3, 2021.




Figure B32: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, June 3, 2021.




Figure B33: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, June 3, 2021.




Figure B34: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, June 1, 2021.




Figure B35: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, June 1, 2021.






Figure B36: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, July 6, 2021.




Figure B37: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, July 8, 2021.




Figure B38: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, July 8, 2021.




Figure B39: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, July 8, 2021.




Figure B40: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, July 6, 2021.




Figure B41: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, August 9, 2021.




Figure B42: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, August 9, 2021.




Figure B43: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, August 9, 2021.






Figure B44: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, August 11, 2021.




Figure B45: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, August 11, 2021.




Figure B46: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, September 9, 2021.




Figure B47: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, September 9, 2021.




Figure B48: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, September 9, 2021.




Figure B49: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, September 7, 2021.




Figure B50: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, September 7, 2021.




Figure B51: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, October 7, 2021.






Figure B52: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, October 7, 2021.




Figure B53: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, October 7, 2021.




Figure B54: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, October 19, 2021.




Figure B55: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, October 12, 2021.




Figure B56: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, November 9, 2021.




Figure B57: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, November 9, 2021.




Figure B58: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, November 9, 2021.




Figure B59: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, November 2, 2021.






Figure B60: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, November 2, 2021.




Figure B61: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 1, December 9, 2021.




Figure B62: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 2, December 9, 2021.




Figure B63: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for the Intake, December 9, 2021.




Figure B64: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 3, December 2, 2021.




Figure B65: Lake Whatcom water column profiles showing temperature, pH, conductivity, and dissolved oxygen for Site 4, December 2, 2021.

## **B.2 Long-term YSI/Hydrolab Data (1988-present)**




Figure B66: Lake Whatcom historic temperature data for Site 1.






Figure B67: Lake Whatcom historic temperature data for Site 2.




Figure B68: Lake Whatcom historic temperature data for the Intake.




Figure B69: Lake Whatcom historic temperature data for Site 3.




Figure B70: Lake Whatcom historic temperature data for Site 4.




Figure B71: Lake Whatcom historic dissolved oxygen data for Site 1.




Figure B72: Lake Whatcom historic dissolved oxygen data for Site 2.




Figure B73: Lake Whatcom historic dissolved oxygen data for the Intake. See discussion of the low dissolved oxygen value in Matthews et al. (2014).




Figure B74: Lake Whatcom historic dissolved oxygen data for Site 3.






Figure B75: Lake Whatcom historic dissolved oxygen data for Site 4.




Figure B76: Lake Whatcom historic pH data for Site 1.




Figure B77: Lake Whatcom historic pH data for Site 2.




Figure B78: Lake Whatcom historic pH data for the Intake.




Figure B79: Lake Whatcom historic pH data for Site 3.




Figure B80: Lake Whatcom historic pH data for Site 4.




Figure B81: Lake Whatcom historic conductivity data for Site 1. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B82: Lake Whatcom historic conductivity data for Site 2. The decreasing conductivity trend is the result of changing to more sensitive equipment.






Figure B83: Lake Whatcom historic conductivity data for the Intake. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B84: Lake Whatcom historic conductivity data for Site 3. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B85: Lake Whatcom historic conductivity data for Site 4. The decreasing conductivity trend is the result of changing to more sensitive equipment.

### **B.3 Long-term Water Quality Data (1988-present)**




Figure B86: Lake Whatcom alkalinity data for Site 1.




Figure B87: Lake Whatcom alkalinity data for Site 2.




Figure B88: Lake Whatcom alkalinity data for the Intake site.




Figure B89: Lake Whatcom alkalinity data for Site 3.






Figure B90: Lake Whatcom alkalinity data for Site 4.




Figure B91: Lake Whatcom turbidity data for Site 1.




Figure B92: Lake Whatcom turbidity data for Site 2.




Figure B93: Lake Whatcom turbidity data for the Intake site.




Figure B94: Lake Whatcom turbidity data for Site 3.




Figure B95: Lake Whatcom turbidity data for Site 4.




Figure B96: Lake Whatcom ammonium data for Site 1.




Figure B97: Lake Whatcom ammonium data for Site 2.






Figure B98: Lake Whatcom ammonium data for the Intake site.




Figure B99: Lake Whatcom ammonium data for Site 3.




Figure B100: Lake Whatcom ammonium data for Site 4.




Figure B101: Lake Whatcom nitrate/nitrite data for Site 1.




Figure B102: Lake Whatcom nitrate/nitrite data for Site 2.




Figure B103: Lake Whatcom nitrate/nitrite data for the Intake site.




Figure B104: Lake Whatcom nitrate/nitrite data for Site 3.




Figure B105: Lake Whatcom nitrate/nitrite data for Site 4.






Figure B106: Lake Whatcom total nitrogen data for Site 1.




Figure B107: Lake Whatcom total nitrogen data for Site 2.




Figure B108: Lake Whatcom total nitrogen data for the Intake site.




Figure B109: Lake Whatcom total nitrogen data for Site 3.




Figure B110: Lake Whatcom total nitrogen data for Site 4.




Figure B111: Lake Whatcom soluble phosphate data for Site 1.




Figure B112: Lake Whatcom soluble phosphate data for Site 2.




Figure B113: Lake Whatcom soluble phosphate data for the Intake site.






Figure B114: Lake Whatcom soluble phosphate data for Site 3.




Figure B115: Lake Whatcom soluble phosphate data for Site 4.




Figure B116: Lake Whatcom total phosphorus data for Site 1.




Figure B117: Lake Whatcom total phosphorus data for Site 2.




Figure B118: Lake Whatcom total phosphorus data for the Intake site.




Figure B119: Lake Whatcom total phosphorus data for Site 3.




Figure B120: Lake Whatcom total phosphorus data for Site 4.




Figure B121: Lake Whatcom chlorophyll data for Site 1.






Figure B122: Lake Whatcom chlorophyll data for Site 2.




Figure B123: Lake Whatcom chlorophyll data for the Intake site.




Figure B124: Lake Whatcom chlorophyll data for Site 3.




Figure B125: Lake Whatcom chlorophyll data for Site 4.




Figure B126: Lake Whatcom Secchi depths for Site 1.




Figure B127: Lake Whatcom Secchi depths for Site 2.




Figure B128: Lake Whatcom Secchi depths for the Intake site.




Figure B129: Lake Whatcom Secchi depths for Site 3.






Figure B130: Lake Whatcom Secchi depths for Site 4.




Figure B131: Lake Whatcom fecal coliform data for Site 1.




Figure B132: Lake Whatcom fecal coliform data for Site 2.




Figure B133: Lake Whatcom fecal coliform data for the Intake site.




Figure B134: Lake Whatcom fecal coliform data for Site 3.




Figure B135: Lake Whatcom fecal coliform data for Site 4.




Figure B136: Lake Whatcom plankton data for Site 1.




Figure B137: Lake Whatcom plankton data for Site 2.






Figure B138: Lake Whatcom plankton data for the Intake Site.




Figure B139: Lake Whatcom plankton data for Site 3.




Figure B140: Lake Whatcom plankton data for Site 4.




Figure B141: Lake Whatcom plankton data for Site 1, with Chrysophyta omitted to show remaining plankton groups.




Figure B142: Lake Whatcom plankton data for Site 2, with Chrysophyta omitted to show remaining plankton groups.




Figure B143: Lake Whatcom plankton data for the Intake Site, with Chrysophyta omitted to show remaining plankton groups.




Figure B144: Lake Whatcom plankton data for Site 3, with Chrysophyta omitted to show remaining plankton groups.




Figure B145: Lake Whatcom plankton data for Site 4, with Chrysophyta omitted to show remaining plankton groups.



## **B.4 Lake Whatcom Tributary Data (2004-present)**

The figures in this appendix include the monthly or biannual baseline data collected from 2004 through the current monitoring period. Each figure includes a dashed (blue) horizontal line that shows the median value for Smith Creek and a solid (red) horizontal line that shows the median value for each creek. Smith Creek was chosen as a reference because it is a major tributary to the lake and has a history of being relatively unpolluted. The figures were scaled to include all but extreme outliers; off-scale outliers are listed in Table [B1](#) (page [250](#)).

Site	Date	Parameter	Concentration
Anderson	January 10, 2006	Total susp. solids	168.8 mg/L
Austin	January 10, 2006	Total susp. solids	166.5 mg/L
Brannian	March 3, 2014	Total phosphorus	349.8 $\mu\text{g-P/L}$
	March 3, 2014	Total susp. solids	328.5 mg/L
Millwheel	February 8, 2005	Ammonium	569.4 $\mu\text{g-N/L}$
	February 8, 2005	Soluble phosphate	116.5 $\mu\text{g-P/L}$
	July 11, 2011	Ammonium	291.7 $\mu\text{g-N/L}$
	October 12, 2011	Total phosphorus	521.8 $\mu\text{g-P/L}$
	September 12, 2012	Ammonium	837.7 $\mu\text{g-N/L}$
	September 12, 2012	Total phosphorus	452.2 $\mu\text{g-P/L}$
	July 8, 2014	Total phosphorus	788.2 $\mu\text{g-P/L}$
	July 8, 2014	Soluble phosphate	165.1 $\mu\text{g-P/L}$
	July 8, 2014	Ammonium	1956.4 $\mu\text{g-N/L}$
	September 9, 2014	Total phosphorus	263.5 $\mu\text{g-P/L}$
	October 9, 2018	Total phosphorus	1,342 $\mu\text{g-P/L}$
	July 12, 2019	Total phosphorus	292.0 $\mu\text{g-P/L}$
	July 12, 2019	Ammonium	291.0 $\mu\text{g-N/L}$
Olsen	January 10, 2006	Total susp. solids	166.9 mg/L
Park Place	August 1, 2006	F. coliforms	18,000 cfu/100 mL
	July 18, 2017	F. coliforms	19,000 cfu/100 mL
	May 14, 2019	Ammonium	693.3 $\mu\text{g-N/L}$
	May 14, 2019	Soluble phosphate	111.8 $\mu\text{g-P/L}$
Silver Beach	August 1, 2006	F. coliforms	12,000 cfu/100 mL

Table B1: List of outliers omitted from Figures B146–B184 to preserve scale.




Figure B146: Temperature data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B147: Temperature data Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B148: Temperature data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B149: Dissolved oxygen data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B150: Dissolved oxygen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B151: Dissolved oxygen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B152: Tributary pH data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B153: Tributary pH data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B154: Tributary pH data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B155: Conductivity data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B156: Conductivity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B157: Conductivity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B158: Alkalinity data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B159: Alkalinity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B160: Alkalinity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B161: Total suspended solids data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B162: Total suspended solids data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B163: Total suspended solids data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B164: Turbidity data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B165: Turbidity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B166: Turbidity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B167: Ammonium data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B168: Ammonium data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B169: Ammonium data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B170: Nitrate/nitrite data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B171: Nitrate/nitrite data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B172: Nitrate/nitrite data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B173: Total nitrogen data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B174: Total nitrogen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B175: Total nitrogen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B176: Soluble phosphate data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B177: Soluble phosphate data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B178: Soluble phosphate data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B179: Total phosphorus data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B180: Total phosphorus data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B181: Total phosphorus data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B182: Fecal coliform data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B183: Fecal coliform data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B184: Fecal coliform data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.

## C Quality Control

### C.1 Performance Evaluation Reports

In order to maintain a high degree of accuracy and confidence in the water quality data all personnel associated with this project were trained according to standard operating procedures for the methods listed in Table 2.1 (page 18). Single-blind quality control tests were conducted as part of the IWS laboratory certification process (Table C1).

### C.2 Laboratory Duplicates, Spikes, and Check Standards

Ten percent of all samples analyzed in the laboratory were duplicated to measure analytical precision. Sample matrix spikes were analyzed during each analytical run to evaluate analyte recovery for the nutrient analyses (ammonium, nitrate/nitrite, total nitrogen, soluble reactive phosphate, and total phosphorus). External check standards were analyzed during each analytical run to evaluate measurement precision and accuracy.<sup>31</sup> The quality control results for laboratory duplicates, matrix spikes, and check standards are plotted in control charts (Figures C1–C30, pages 293–322).

### C.3 Field Duplicates

Ten percent of all samples collected in the field were duplicated to measure sample replication (Figures C31–C48, pages 323–340). Samples collected using field meters (conductivity, dissolved oxygen, and pH) were evaluated using water samples collected from the same depth as the field meter measurement. The absolute mean difference for the field duplicates was calculated as follows:

$$\text{Absolute mean difference} = \frac{\sum |\text{Original Sample} - \text{Duplicate Sample}|}{\text{number of duplicate pairs}}$$

---

<sup>31</sup>External check standards are not available for all analytes.

	Reported Value	Assigned Value	Acceptance Limits	Test Result
Specific conductivity ( $\mu\text{S}/\text{cm}$ at $25^\circ\text{C}$ )	413	413	372–454	accept
Total alkalinity (mg/L as $\text{CaCO}_3$ )	139	142	121–163	accept
Ammonium nitrogen, manual (mg-N/L)	5.98	5.52	4.29–6.77	accept
Ammonium nitrogen, auto (mg-N/L)	5.46	5.52	4.29–6.77	accept
Nitrate/nitrite nitrogen, auto (mg-N/L)	7.31	7.10	5.86–8.27	accept
Nitrite nitrogen, auto (mg-N/L)	0.51	0.50	0.37–0.63	accept
Organic carbon, dissolved (mg/L)	2.90	2.96	2.60–3.36	accept
	5.09	5.26	4.66–5.79	accept
Organic carbon, total (mg/L)	14.6	15.2	12.0–18.5	accept
Orthophosphate, manual (mg-P/L)	2.25	2.26	1.92–2.60	accept
Orthophosphate, auto (mg-P/L)	2.27	2.26	1.92–2.60	accept
Total phosphorus, manual (mg-P/L)	3.95	3.95	3.26–4.61	accept
Total phosphorus, auto (mg-P/L)	3.93	3.95	3.26–4.61	accept
pH	8.39	8.40	8.20–8.60	accept
Solids, non-filterable (mg/L)	33.3	38.6	28.7–45.1	accept
Turbidity (NTU)	9.09	9.83	7.96–11.7	accept

Table C1: Single-blind quality control results, WP–270 (12/18/2020); all results were within acceptance limits. IWS is applying for certification for total and dissolved organic carbon analyses; the 2018 performance evaluation results for dissolved organic carbon (WP–114, 05/31/2018; WP–116, 11/29/2018) are reported here.

Year	Month	Analyte	Type	Testing or Training	Value
2019	June	pH	lab duplicate	training	-0.6
	October	orthophosphate	spike	training	55
	October	turbidity	lab duplicate	training	0.77
	November	orthophosphate	spike	training	136
	December	orthophosphate	lab duplicate	training	5.05
2020	January	nitrate/nitrite	lab duplicate	training	-135.36
	January	nitrate/nitrite	lab duplicate	training	124.92
	February	total phosphorus	lab duplicate	training	46.22
	July	dissolved oxygen	lab duplicate	training	0.38
	September	ammonium	lab duplicate	training	90.43
	November	orthophosphate	lab duplicate	testing	6.37
2021	December	nitrate/nitrite	check	testing	34
	January	alkalinity	lab duplicate	testing	-1
	January	total nitrogen	lab duplicate	testing	-129.43
	January	total phosphorus	spike	testing	146
	February	alkalinity	lab duplicate	testing	-2
	February	total suspended solids	lab duplicate	testing	-8.25
	April	alkalinity	check	testing	0.09
	April	conductivity	lab duplicate	testing	-2.8
	May	dissolved oxygen	lab duplicate	testing	0.34
	July	nitrate/nitrite	spike	testing	67
July	nitrate/nitrite	spike	testing	63	
September	total nitrogen	lab duplicate	testing	-152.35	

Table C2: Data in this table denote quality control values that exceeded  $\pm 4$  std. dev. from the training mean. Unplotted points were included in QC calculations, but were not plotted to preserve plotting scale (Figures C1–C30).





Figure C1: Alkalinity laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C2: Alkalinity high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C3: Alkalinity low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C4: Chlorophyll laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.







Figure C5: Conductivity laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C6: Dissolved oxygen laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C7: Nitrogen (ammonium) laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C8: Nitrogen (ammonium) spike recoveries for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of spike data.





Figure C9: Nitrogen (ammonium) high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C10: Nitrogen (ammonium) low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C11: Nitrogen (nitrate/nitrite) laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C12: Nitrogen (nitrate/nitrite) spike recoveries for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of spike data.







Figure C13: Nitrogen (nitrate/nitrite) high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C14: Nitrogen (nitrate/nitrite) low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C15: Nitrogen (total) laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C16: Nitrogen (total) spike recoveries for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of spike data.





Figure C17: Nitrogen (total) high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C18: Nitrogen (total) low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C19: Laboratory pH duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C20: Phosphorus (soluble reactive phosphate) laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.







Figure C21: Phosphorus (soluble reactive phosphate) spike recoveries for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of spike data.





Figure C22: Phosphorus (soluble reactive phosphate) high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C23: Phosphorus (soluble reactive phosphate) low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C24: Phosphorus (total) laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C25: Phosphorus (total) spike recoveries for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of spike data.





Figure C26: Phosphorus (total) high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C27: Phosphorus (total) low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.







Figure C28: Total suspended solids laboratory duplicates for the Lake Whatcom monitoring program (tributary and storm water samples). Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Total Suspended Solids Check Standards, Training Data



Total Suspended Solids Check Standards, Test Data

Figure C29: Total suspended solids check standards for the Lake Whatcom monitoring program (tributary and storm water samples). Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of check standard data.





Figure C30: Turbidity laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.




Figure C31: Alkalinity field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship.




Figure C32: Alkalinity field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship.




Figure C33: Chlorophyll field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship. The outlier identified likely represents slight differences in sample depth of the field duplicate near the steep thermocline.




Figure C34: Conductivity field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship.




Figure C35: Dissolved oxygen field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship. The labeled outlier was collected when extreme gradients were present. Field meter samples were collected at true depth; Winkler samples were measured in the laboratory from samples collected using a marked line, which will be slightly shallower than true depth. There was a systematic bias between the Winkler and field meter results, with the Winkler results  $\sim 0.5$  mg/L lower than the field meter. This is within typical ranges for Winkler vs. field meter comparisons (Johengen, et al., 2016).




Figure C36: Dissolved oxygen field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship. There was a systematic bias between the Winkler and field meter results, with the Winkler results  $\sim 0.5$  mg/L lower than the field meter. This is within typical ranges for Winkler vs. field meter comparisons (Johengen, et al., 2016). Park Place is often an outlier because the sample is taken from a manhole, which can result in low readings from the field meter.






Figure C37: Nitrogen (ammonium) field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit. The outlier identified likely represents slight differences in sample depth of the field duplicate near the steep thermocline. The high degree of scatter is due to the low concentrations of the samples.




Figure C38: Nitrogen (ammonium) field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit. The high degree of scatter is due to the low concentrations of the samples.




Figure C39: Nitrogen (nitrate/nitrite) field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit.




Figure C40: Nitrogen (nitrate/nitrite) field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit.




Figure C41: Nitrogen (total) field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship. The scatter around the line is within the range of the laboratory QC standards (Figure C15). The October outlier likely represents slight differences in sample depth of the field duplicate near the steep thermocline.




Figure C42: Nitrogen (total) field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship.




Figure C43: Field duplicates for pH from the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship. A systematic bias was observed for pH values  $\leq 7.0$ . This bias may be the result of the time lag between field sampling and laboratory measurements, temperature differences, and/or the effects of gas exchange once the sample is exposed to air.




Figure C44: Phosphorus (soluble reactive phosphate) field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit. The high degree of scatter is due to the low concentrations of the samples.






Figure C45: Phosphorus (total) field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit. The outlier identified likely represents slight differences in sample depth of the field duplicate near the steep thermocline. The high degree of scatter is due to the low concentrations of the samples.




Figure C46: Phosphorus (total) field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit. The outlier identified likely represents sediment contamination of the duplicate sample.




Figure C47: Total suspended solids field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship; horizontal reference line shows current detection limit. The outliers identified likely represent sediment contamination of the duplicate sample.




Figure C48: Turbidity field duplicates for the 2020/2021 Lake Whatcom monitoring program (lake samples). Diagonal reference line shows 1:1 relationship.




Figure C49: Turbidity field duplicates for the 2020/2021 Lake Whatcom monitoring program (tributary samples). Diagonal reference line shows 1:1 relationship. The outlier identified likely represents sediment contamination of the duplicate sample.

## D Lake Whatcom Online Data

The following **readme** file describes the electronic data posted at the IWS web site ([www.wvu.edu/iws](http://www.wvu.edu/iws)) and additional data available from IWS. Please contact the Director of the Institute for Watershed Studies if you have questions or trouble accessing the online data.

```
*****
* README FILE - LAKE WHATCOM ONLINE DATA
* THIS FILE WAS UPDATED DECEMBER 20, 2021
*****
Most of the Lake Whatcom water quality data are available in
electronic format at the IWS website (http://www.wvu.edu/iws) or from
the IWS Director.
```

The historic and current detection limits and abbreviations for each parameter are listed in the annual reports. The historic detection limits for each parameter were estimated based on recommended lower detection ranges, instrument limitations, and analyst judgment on the lowest repeatable concentration for each test. Over time, some analytical techniques have improved so that current detection limits are usually lower than historic detection limits. Because the Lake Whatcom data set includes long-term monitoring data, which have been collected using a variety of analytical techniques, this report sets conservative detection limits to allow comparisons between years.

All files are comma-separated ascii data files. The code "NA" has been entered into all empty cells in the ascii data files to fill in unsampled dates and depths, missing data, etc. Questions about missing data should be directed to the IWS Director.

Unless otherwise indicated, the electronic data files have NOT been censored to flag or otherwise identify below detection and above detection values. As a result, the ascii files may contain negative values due to linear extrapolation of the standards regression curve for below detection data. It is essential that any statistical or analytical results that are generated using these data be reviewed by someone familiar with statistical uncertainty associated with uncensored data.

\*\*\*\*\*

\* ONLINE LAKE DATA FILES:

\*\*\*\*\*

Hydrolab/YSI data

1988\_hl.csv, 1989\_hl.csv, 1990\_hl.csv, 1991\_hl.csv, 1992\_hl.csv  
1993\_hl.csv, 1994\_hl.csv, 1995\_hl.csv, 1996\_hl.csv, 1997\_hl.csv  
1998\_hl.csv, 1999\_hl.csv, 2000\_hl.csv, 2001\_hl.csv, 2002\_hl.csv  
2003\_hl.csv, 2004\_hl.csv, 2005\_hl.csv, 2006\_hl.csv, 2007\_hl.csv  
2008\_hl.csv, 2009\_hl.csv, 2010\_hl.csv, 2011\_hl.csv, 2012\_hl.csv  
2013\_hl.csv, 2014\_hl.csv, 2015\_hl.csv, 2016\_hl.csv, 2017\_hl.csv  
2018\_hl.csv, 2019\_hl.csv, 2020\_hl.csv, 2021\_hl.csv

Water quality data

1988\_wq.csv, 1989\_wq.csv, 1990\_wq.csv, 1991\_wq.csv, 1992\_wq.csv  
1993\_wq.csv, 1994\_wq.csv, 1995\_wq.csv, 1996\_wq.csv, 1997\_wq.csv  
1998\_wq.csv, 1999\_wq.csv, 2000\_wq.csv, 2001\_wq.csv, 2002\_wq.csv  
2003\_wq.csv, 2004\_wq.csv, 2005\_wq.csv, 2006\_wq.csv, 2007\_wq.csv  
2008\_wq.csv, 2009\_wq.csv, 2010\_wq.csv, 2011\_wq.csv, 2012\_wq.csv  
2013\_wq.csv, 2014\_wq.csv, 2015\_wq.csv, 2016\_wq.csv, 2017\_wq.csv  
2018\_wq.csv, 2019\_wq.csv, 2020\_wq.csv, 2021\_wq.csv

Plankton counts

plankton.csv

The \*\_hl.csv files include: site, depth (m), month, day, year, temp (temperature, C), pH, cond (conductivity, uS/cm), do (dissolved oxygen, mg/L), lcond (lab conductivity qc, uS/cm), secchi (secchi depth, m).

The \*\_wq.csv files include: site, depth (m), month, day, year, alk (alkalinity, mg/L as CaCO3), turb (turbidity, NTU), nh3 (ammonium, ug-N/L), tn (total persulfate nitrogen, ug-N/L), nos (nitrate/nitrite, ug-N/L), srp (soluble reactive phosphate, ug-P/L), tp (total persulfate phosphorus, ug-P/L), chl (chlorophyll, ug/L).

The plankton.csv file includes: site, depth (m), month, day, year, zoop (zooplankton, #/L), chry (chrysophyta, #/L), cyan (cyano-bacteria, #/L), chlo (chlorophyta, #/L), pyrr (pyrrophyta, #/L).

\*\*\*\*\*

\* ONLINE HYDROGRAPH DATA FILES:

\*\*\*\*\*

WY1998.csv, WY1999.csv, WY2000\_rev.csv (rev. 3/8/2012), WY2001.csv,  
WY2002.csv, WY2003.csv, WY2004\_rev.csv (rev. 6/21/2006), WY2005.csv,  
WY2006.csv, WY2007.csv (rev. July 31, 2008), WY2008.csv, WY2009.csv,

WY2010.csv, WY2011.csv, WY2012.csv, WY2013.csv, WY2014.csv, WY2015.csv  
WY2016.csv, WY2017.csv, WY2018.csv, WY2019.csv, WY2020.csv, WY2021.csv

The WY\*.csv files include: month, day, year, hour, min, sec, ander.g (anderson gauge height, ft), ander.cfs (anderson discharge, cfs), austin.g (austin gauge height, ft), austin.cfs (austin discharge, cfs), smith.g (smith gauge height, ft), smith.cfs (smith discharge, cfs). Anderson Creek hydrograph data were deleted in WY2000\_rev.csv due to uncertainty about the gauge height; Anderson Creek data are available for WY1998, WY1999, and WY2001-WY2007. Beginning with WY2002, the variable "time" replaced "hour, min, sec," with time reported daily on a 24-hr basis. Data are reported as Pacific Standard Time without Daylight Saving Time adjustment.

\*\*\*\*\*

\* STORM WATER AND TRIBUTARY DATA FILES

\*\*\*\*\*

The storm water and tributary data include composite and grab samples from numerous sites in the Lake Whatcom watershed (1994--present), representing a variety of study objectives and sampling intensities over time. The electronic data files are not posted online, but may be obtained by contacting the Institute for Watershed Studies.

\*\*\*\*\*

\* SITE CODES

\* ALL FILES - INCLUDES DISCONTINUED SITES AND OFF-LINE DATA

\*\*\*\*\*

The site codes in the data are as follows:

11 = Lake Whatcom Site 1  
21 = Lake Whatcom Intake site  
22 = Lake Whatcom Site 2  
31 = Lake Whatcom Site 3  
32 = Lake Whatcom Site 4  
33 = Strawberry Sill site S1  
34 = Strawberry Sill site S2  
35 = Strawberry Sill site S3

AlabamaVault inlet	= Alabama canister vault inlet
AlabamaVault outlet	= Alabama canister vault outlet
Brentwood inlet	= Brentwood wet pond inlet
Brentwood outlet	= Brentwood wet pond outlet
ParkPlace cell1	= Park Place wet pond cell 1
ParkPlace cell2	= Park Place wet pond cell 2
ParkPlace cell3	= Park Place wet pond cell 3
ParkPlace inlet	= Park Place wet pond inlet



ParkPlace outlet = Park Place wet pond outlet  
Parkstone\_swale inlet = Parkstone grass swale inlet  
Parkstone\_swale outlet = Parkstone grass swale outlet  
Parkstone\_pond inlet = Parkstone wet pond inlet  
Parkstone\_pond outlet = Parkstone wet pond outlet  
SouthCampus inlet = South Campus storm water facility inlet  
SouthCampus outletE = South Campus storm water facility east outlet  
SouthCampus outletW = South Campus storm water facility west outlet  
Sylvan inlet = Sylvan storm drain inlet  
Sylvan outlet = Sylvan storm drain outlet  
Wetland outlet = Grace Lane wetland

CW1 = Smith Creek (see alternate code below)  
CW2 = Silver Beach Creek (see alternate code below)  
CW3 = Park Place drain (see alternate code below)  
CW4 = Blue Canyon Creek (see alternate code below)  
CW5 = Anderson Creek (see alternate code below)  
CW6 = Wildwood Creek (discontinued in 2004)  
CW7 = Austin Creek (see alternate code below)

The following tributary site codes were used for the expanded 2004–2006 tributary monitoring project

AND = Anderson Creek (same location as CW5 above)  
BEA1 = Austin.Beaver.confluence  
AUS = Austin.lower (same location as CW7 above)  
BEA2 = Austin.upper  
BEA3 = Beaver.upper  
BLU = BlueCanyon (same location as CW4 above)  
BRA = Brannian  
CAR = Carpenter  
EUC = Euclid  
MIL = Millwheel  
OLS = Olsen  
PAR = ParkPlace (same location as CW3 above)  
SIL = SilverBeach (same location as CW2 above)  
SMI = Smith (same location as CW1 above)  
WHA = Whatcom

\*\*\*\*\*  
\* VERIFICATION PROCESS FOR THE LAKE WHATCOM DATA FILES

\*\*\*\*\*  
During the summer of 1998 the Institute for Watershed Studies began creating an electronic data file that would contain long term data records for Lake Whatcom. These data were to be included with annual Lake Whatcom monitoring reports. This was the first attempt to make a long-term Lake Whatcom data record available to the public. Because these data had been generated using different quality control plans over the years, a comprehensive re-verification process was done.

The re-verification started with printing a copy of the entire data file and checking 5% of all entries against historic laboratory bench sheets and field notebooks. If an error was found, the entire set of values for that analysis were reviewed for the sampling period containing the error. Corrections were noted in the printed copy and entered into the electronic file; all entries were dated and initialed in the archive copy.

Next, all data were plotted and descriptive statistics (e.g., minimum, maximum) were computed to identify outliers and unusual results. All outliers and unusual data were verified against original bench sheets. A summary of decisions pertaining to these data is presented below. All verification actions were entered into the printed copy, dated, and initialed by the IWS director.

The following is a partial list of the changes made to the verified Lake Whatcom data files. For detailed information refer to the data verification archive files in the IWS library.

Specific Deletions: 1) Rows containing only missing values were deleted. 2) All lab conductivity for February 1993 were deleted for cause: meter inadequate for low conductivity readings (borrowed Huxley's student meter). 3) All Hydrolab conductivity from April - December 1993 were deleted for cause: Hydrolab probe slowly lost sensitivity. Probe was replaced and Hydrolab was reconditioned prior to the February 1994 sampling. 4) All 1993 Hydrolab dissolved oxygen data less than or equal to 2.6 mg/L were deleted for cause: Hydrolab probe lost sensitivity at low oxygen concentrations. Probe was replaced and Hydrolab was reconditioned prior to February 1994 sampling. 5) All srp and tp data were deleted (entered as "missing" in 1989) from the July 10, 1989 wq data due to sample contamination in at least three samples. 6) December 2, 1991, Site 3, 0 m conductivity point deleted due to inconsistency with adjacent points. 7) December 15, 1993, Site 4, 80 m lab conductivity point deleted because matching

field conductivity data are absent and point is inconsistent with all other lab conductivity points. 8) November 4, 1991, Site 2, 17-20 m, conductivity points deleted due to evidence of equipment problems related to depth. 9) February 2, 1990, Site 1, 20 m, soluble reactive phosphate and total phosphorus points deleted due to evidence of sample contamination. 10) August 6, 1990, Site 1, 0 m, soluble reactive phosphate and total phosphorus points deleted due to evidence of sample contamination. 11) October 5, 1992, Site 3, 80 m, all data deleted due to evidence of sample contamination in turbidity, ammonium, and total phosphorus results. 12) August 31, 1992, Site 3, 5 m, soluble reactive phosphate and total phosphorus data deleted due to probable coding error. 13) All total Kjeldahl nitrogen data were removed from the historic record. This was not due to errors with the data but rather on-going confusion over which records contained total persulfate nitrogen and which contained total Kjeldahl nitrogen. The current historic record contains only total persulfate nitrogen. Total Kjeldahl nitrogen data were retained in the IWS data base, but not in the long-term Lake Whatcom data files.

\*\*\*\*\*

\* ROUTINE DATA VERIFICATION PROCESS

\*\*\*\*\*

1994-present: The Lake Whatcom data are verified using a four step method: 1) The results are reviewed as they are generated. Outliers are checked for possible analytical or computational errors. This step is completed by the Laboratory Analyst and IWS Laboratory Supervisor. 2) The results are reviewed monthly or quarterly and sent to the City. Unusual results are identified. This step is completed by the IWS Director. 3) The results are reviewed on an annual basis and discussed in the Lake Whatcom Monitoring Program Final Report. Unusual results are identified, and explained, if possible. This step is completed by the IWS Director, IWS Laboratory Supervisor, and Laboratory Analyst. 4) Single-blind quality control samples, laboratory duplicates, and field duplicates are analyzed as specified in the Lake Whatcom Monitoring Program contract and in the IWS Laboratory Certification requirements. Unusual results that suggest instrumentation or analytical problems are reported to the IWS Director and City. The results from these analyses are summarized in the annual report.

1987-1993: The lake data were reviewed as above except that the IWS Director's responsibilities were delegated to the Principle Investigator in charge of the lake monitoring contract.

Prior to 1987: Data were informally reviewed by the Laboratory Analyst and IWS Director. Laboratory and field duplicates were commonly included as part of the analysis process, but no formal (i.e., written) quality control program was in place. Laboratory logs were maintained for most analyses, so it is possible to verify data against original analytical results. It is also possible to review laboratory quality control results for some analyses.