

2013

On the Hörmander Classes of Bilinear Pseudodifferential Operators, II

Árpád Bényi

Western Washington University, arpad.benyi@wwu.edu

Frederic Bernicot

Universite de Nantes

Diego Maldonado

Kansas State University

Rodolfo H. (Rodolfo Humberto) Torres

University of Kansas

Virginia Naibo

Kansas State University

Follow this and additional works at: https://cedar.wwu.edu/math_facpubs



Part of the [Mathematics Commons](#)

Recommended Citation

Arpad Benyi, Frederic Bernicot, Diego Maldonado, Virginia Naibo, Rodolfo H.Torres, On the Hormander classes of bilinear pseudodifferential operators II, *Indiana Univ. Math. J.* 62 (2013), 1733-1764

This Article is brought to you for free and open access by the College of Science and Engineering at Western CEDAR. It has been accepted for inclusion in Mathematics by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

On the Hörmander Classes of Bilinear Pseudodifferential Operators II

ÁRPAD BÉNYI, FRÉDÉRIC BERNICOT, DIEGO MALDONADO,
VIRGINIA NAIBO & RODOLFO H. TORRES

ABSTRACT. Boundedness properties for pseudodifferential operators with symbols in the bilinear Hörmander classes of sufficiently negative order are proved. The results are obtained in the scale of Lebesgue spaces, and in some cases, end-point estimates involving weak-type spaces and BMO are provided as well. From the Lebesgue space estimates, Sobolev ones are then easily obtained using functional calculus and interpolation. In addition, it is shown that, in contrast with the linear case, operators associated with symbols of order zero may fail to be bounded on products of Lebesgue spaces.

1. INTRODUCTION

In this article, we continue the systematic study of the general Hörmander classes of bilinear pseudodifferential operators $BS_{\rho,\delta}^m$ (see the next section for definitions) started in [2]. While the work in [2] focussed mainly on basic properties related to the symbolic calculus of the bilinear pseudodifferential operators and some point-wise estimates for their kernels, the present work addresses boundedness properties on the full scale of Lebesgue spaces. The general properties developed in [2] will become very useful in this current work, and will allow us to provide a fairly complete range of results.

The literature on bilinear pseudodifferential operators continues to grow, and [2] also gives a historical account and motivations, as well as numerous references on the subject. We would like to reiterate here that most results so far have dealt with the cases $\rho = 1$ and $\rho = 0$. For the first value of ρ , the available boundedness and unboundedness results, and other properties of the classes $BS_{1,\delta}^0$, are

similar to the ones in the linear situation. They are closely tied to the (bilinear) Calderón-Zygmund theory, which was started by Coifman-Meyer in the 1970s (see, e.g., [13] and the references therein) and was further developed by Christ-Journé [11], Kenig-Stein [21], and Grafakos-Torres [15]. See also Bényi-Torres [3] and Maldonado-Naibo [23]. The value of $\rho = 0$, however, produces some surprises, and the possible theory deviates from the linear situation. In particular, the famous Calderón-Vaillancourt theorem [10] does not hold unless further properties on the symbols in $BS_{0,0}^0$ are imposed; see Bényi-Torres [4] and Bernicot-Shrivastava [9].

One important contribution for other values of ρ , almost the exception so far, is the recent work of Michalowski-Rule-Staubach [24]. Since, for example, the class $BS_{0,0}^0$ does not map $L^\infty \times L^2 \rightarrow L^2$, it was asked in [2] (and some answers were provided) about results of the form $X \times L^2 \rightarrow L^2$ with some functional space X smaller than L^∞ and symbols in $BS_{\rho,\delta}^0$. The question of whether the classes $BS_{\rho,\delta}^0$ produce operators that are bounded on some product of Lebesgue spaces when $0 \leq \delta < \rho$ was left unanswered in [2] (recall the keystone result that the linear class $S_{\rho,\delta}^0$ is bounded on L^2 , as proved by Hörmander [17]). Likewise, in [24], the authors asked about which negative values of $m = m(\rho)$ produce classes $BS_{\rho,\delta}^m$ for which the corresponding bilinear pseudodifferential operators are bounded from $L^{p_1} \times L^{p_2}$ into L^p with $1/p_1 + 1/p_2 = 1/p$ and $1 < p_1, p_2, p \leq \infty$. Here, we will expand and improve some of the results in [24] in several directions.

First, we will show that it is very much relevant to look at negative values of m when $\rho < 1$ because operators with symbols in the classes $BS_{\rho,\delta}^0$ may fail to be bounded on any product of Lebesgue spaces. This is proved in Theorem 2.1 below, thus answering in the negative the question left unanswered in [2]. Next, we show in Theorem 2.2 that the values of m provided in [24] can be taken much larger (smaller in absolute value). We succeed in doing so using kernel estimates and the symbolic calculus from [2], also used in [24], but adding arguments involving the complex interpolation of the classes $BS_{\rho,\delta}^m$. Moreover, bringing back the bilinear Calderón-Zygmund theory for sufficiently negative values of m , and using further interpolation arguments, we also obtain results outside the *Banach triangle*, that is, for $1/p_1 + 1/p_2 = 1/p$, but where $\frac{1}{2} < p < 1$. We also obtain appropriate weak-type end-point estimates at one end and a strong one at another. This last is the bilinear analog of a result of C. Fefferman, which was also a keystone in the understanding of linear pseudodifferential operators.

Fefferman [14] showed, in particular, that the linear classes $S_{\rho,0}^{-(1-\rho)n/2}$ (for $0 < \rho < 1$) map $L^\infty \rightarrow \text{BMO}$. The natural conjecture, then, is that $BS_{\rho,0}^{-(1-\rho)n}$ should map $L^\infty \times L^\infty \rightarrow \text{BMO}$, since often the role of n in the linear case is played by $2n$ in the bilinear setting. We are able to prove this conjecture in Theorem 2.4, at least for $0 < \rho < \frac{1}{2}$. Though we use some ideas from [14], new technical difficulties not present in the linear case need to be overcome. In fact, Fefferman used the result of Hörmander that operators with symbols in $S_{\rho,\delta}^0$ are bounded

on L^2 but, as Theorem 2.1 establishes, the analogous result for bilinear operators is false. Instead, we rely on the $L^2 \times L^2 \rightarrow L^2$ boundedness of certain classes of symbols as proved in Theorem 2.3.

The article is organized as follows. In the next section, we include the main definitions, some basic properties, and the precise statements of the main theorems. We also provide some further motivation and applications. The subsequent sections, Sections 3–7, contain the detailed proof of each of the main theorems in the order we list them, except that a series of technical lemmata used in the proof of Theorem 2.4 are postponed until Section 8. Section 9 contains some weighted versions of the results. Further remarks about the results and comparisons to other linear and bilinear ones are provided throughout the paper as well. Upper-case letters are used to label theorems corresponding to known results, while single numbers are used for theorems, lemmas, and corollaries that are proved in this article.

Unless otherwise indicated, the underlying space for the functional classes used will be Euclidean space \mathbb{R}^n . In particular, L^p will stand for $L^p(\mathbb{R}^n)$, and $W^{s,p}$ will stand for $W^{s,p}(\mathbb{R}^n)$, the Sobolev space of functions with “ s derivatives” in L^p . Their respective norms will be denoted $\|f\|_{L^p}$ and $\|f\|_{W^{s,p}}$. Finally, S will indicate the Schwartz class on \mathbb{R}^n .

Throughout, the symbol \lesssim will be used in inequalities where constants are independent of its left and right hand sides.

2. MAIN RESULTS

Let $\delta, \rho \geq 0$ and $m \in \mathbb{R}$. The Hörmander class $S_{\rho,\delta}^m$ (introduced in [17]) consists of symbols $\sigma = \sigma(x, \xi)$, $x, \xi \in \mathbb{R}^n$ such that, for all multi-indices α and β ,

$$\sup_{x, \xi \in \mathbb{R}^n} |\partial_x^\alpha \partial_\xi^\beta \sigma(x, \xi)| (1 + |\xi|)^{-m - \delta|\alpha| + \rho|\beta|} < \infty.$$

For each symbol σ , there is an associated linear pseudodifferential operator T_σ defined by

$$T_\sigma(f)(x) = \int_{\mathbb{R}^n} \sigma(x, \xi) \hat{f}(\xi) e^{ix \cdot \xi} d\xi, \quad f \in S,$$

where \hat{f} denotes the Fourier transform of f .

The bilinear counterpart of $S_{\rho,\delta}^m$ is denoted $BS_{\rho,\delta}^m$. Also, a bilinear symbol $\sigma(x, \xi, \eta)$, $x, \xi, \eta \in \mathbb{R}^n$, belongs to the bilinear Hörmander class $BS_{\rho,\delta}^m$ if, for all multi-indices α, β and γ ,

$$\sup_{x, \xi, \eta \in \mathbb{R}^n} |\partial_x^\alpha \partial_\xi^\beta \partial_\eta^\gamma \sigma(x, \xi, \eta)| (1 + |\xi| + |\eta|)^{-m - \delta|\alpha| + \rho(|\beta| + |\gamma|)} < \infty.$$

For $\sigma \in BS_{\rho,\delta}^m$ and non-negative integers K and N , define

$$\|\sigma\|_{K,N} := \sup_{\substack{|\alpha| \leq K \\ |\beta|, |\gamma| \leq N}} \sup_{x, \xi, \eta \in \mathbb{R}^n} |\partial_x^\alpha \partial_\xi^\beta \partial_\eta^\gamma \sigma(x, \xi, \eta)| (1 + |\xi| + |\eta|)^{-m - \delta|\alpha| + \rho(|\beta| + |\gamma|)}.$$

Then, the family of norms $\{\|\cdot\|_{K,N}\}_{K,N \in \mathbb{N}_0}$ turns $BS_{\rho,\delta}^m$ into a Fréchet space.

For $\sigma \in BS_{\rho,\delta}^m$, we consider the bilinear pseudodifferential operator defined by

$$T_\sigma(f, g)(x) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sigma(x, \xi, \eta) \hat{f}(\xi) \hat{g}(\eta) e^{ix \cdot (\xi + \eta)} d\xi d\eta, \quad f, g \in S.$$

We now proceed to state the new results in this article.

Theorem 2.1. *Let $0 \leq \rho < 1$, $0 \leq \delta \leq 1$, and $1 \leq p, p_1, p_2 < \infty$ such that $1/p = 1/p_1 + 1/p_2$. There exist symbols in $BS_{\rho,\delta}^0$ that give rise to unbounded operators from $L^{p_1} \times L^{p_2}$ into L^p .*

As mentioned in the [Introduction](#), the result in [Theorem 2.1](#) is in contrast with the fact that linear pseudodifferential operators of order zero do produce bounded operators on L^2 . The case $\rho = \delta = 0$ of [Theorem 2.1](#) was proved by Bényi and Torres in [\[4\]](#).

Theorem 2.2. *Let $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$, $1 \leq p_1, p_2 \leq \infty$, and p given by $1/p = 1/p_1 + 1/p_2$. If $\sigma \in BS_{\rho,\delta}^m$ where*

$$m < m(p_1, p_2) := n(\rho - 1) \left(\max \left\{ \frac{1}{2}, \frac{1}{p_1}, \frac{1}{p_2}, 1 - \frac{1}{p} \right\} + \max \left\{ \frac{1}{p} - 1, 0 \right\} \right),$$

then the following statements hold:

- (i) *If $p \geq 1$, then there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T_\sigma(f, g)\|_{L^p} \lesssim \|\sigma\|_{K,N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}$$

for all $f \in L^{p_1}$ and $g \in L^{p_2}$.

- (ii) *If $\rho > 0$, $p < 1$, $p_1 \neq 1$, and $p_2 \neq 1$, then there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T_\sigma(f, g)\|_{L^p} \lesssim \|\sigma\|_{K,N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}$$

for all $f \in L^{p_1}$ and $g \in L^{p_2}$.

- (iii) *If $\rho > 0$, $p < 1$, and $p_1 = 1$ or $p_2 = 1$, then there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T_\sigma(f, g)\|_{L^{p,\infty}} \lesssim \|\sigma\|_{K,N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}$$

for all $f \in L^{p_1}$ and $g \in L^{p_2}$.

When $p \geq 1$ (Banach case), [Theorem 2.2](#) improves the results in [\[24, Theorem 5.5\]](#) by Michalowski, Rule, and Staubach, which require

$$m < n(\rho - 1) \max \left\{ \frac{1}{2}, \left(\frac{2}{p_1} - \frac{1}{2} \right), \left(\frac{2}{p_2} - \frac{1}{2} \right), \left(\frac{3}{2} - \frac{2}{p} \right) \right\}.$$

This improvement is based on the following facts:

- (1) Bilinear pseudodifferential operators with symbols in the classes $BS_{\rho,\delta}^m$ with $m < n(\rho - 1)$ (as opposed to $m < \frac{3}{2}n(\rho - 1)$ used in [24]) are bounded from $L^\infty \times L^\infty$ into L^∞ , with norm bounded by the norm of the symbol (see also Remark 4.2).
- (2) Roughly speaking, the intermediate spaces in the complex interpolation of two bilinear Hörmander classes are other bilinear Hörmander classes.

When $\rho < 1$ (non-Banach case), the result of Theorem 2.2 relies on interpolation arguments using boundedness of operators in the Banach case and bilinear Calderón-Zygmund theory.

We note that the operator T_σ is defined *a priori* on $S \times S$. In Theorem 2.2, $T_\sigma(f, g)$ for $f \in L^{p_1}$ and $g \in L^{p_2}$ denotes the “value” given by a bounded extension of the operator, which exists and is unique in the cases $p_1 < \infty$ and $p_2 < \infty$, and is shown to exist when $p_1 = \infty$ or $p_2 = \infty$.

Theorem 2.3. *If $\sigma(x, \xi, \eta)$, $x, \xi, \eta \in \mathbb{R}^n$, is a bilinear symbol such that*

$$C(\sigma) := \sup_{|\beta| \leq [n/2]+1, |\alpha| \leq 2(2n+1)} \sup_{\xi, \gamma \in \mathbb{R}^n} \|\partial_\xi^\alpha \partial_\gamma^\beta \sigma(\gamma, \xi - \cdot, \cdot)\|_{L^2} < \infty,$$

then T_σ maps continuously $L^2 \times L^2$ into L^2 with $\|T_\sigma\|_{L^2 \times L^2 \rightarrow L^2} \lesssim C(\sigma)$.

Theorem 2.4. *If $\sigma \in BS_{\rho,0}^{n(\rho-1)}$, $0 \leq \rho < \frac{1}{2}$, then there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T_\sigma(f, g)\|_{\text{BMO}} \lesssim \|\sigma\|_{K,N} \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in S.$$

Theorem 2.4, which complements the endpoint $m = n(\rho - 1)$ for $p_1 = p_2 = \infty$ in Theorem 2.2, can be thought of as a bilinear counterpart (when $0 \leq \rho < \frac{1}{2}$ and $\delta = 0$) to the following linear result proved by Ch. Fefferman in [14].

Theorem A (Fefferman [14]). *If σ is a symbol in the linear Hörmander class $S_{\rho,\delta}^{-(n/2)(1-\rho)}$ with $0 \leq \delta < \rho < 1$, then T_σ maps L^∞ continuously into BMO.*

The proof of Theorem A uses the fact that the linear class $S_{\rho,\delta}^0$, $0 < \delta < \rho \leq 1$, maps L^2 continuously into L^2 . The bilinear counterpart of this result is false by Theorem 2.1. Our proof of Theorem 2.4 relies on Fefferman’s ideas and the result given by Theorem 2.3.

Next, we present a result concerning the boundedness properties of bilinear pseudodifferential operators on Lebesgue spaces with indices that satisfy the Sobolev scaling, as opposed to the Hölder scaling employed in the previous theorems.

Theorem 2.5. *Let $0 \leq \delta \leq 1$, $0 < \rho \leq 1$, $s \in (0, 2n)$, and $m_s := 2n(\rho - 1) - \rho s$. If $\sigma \in BS_{\rho,\delta}^{m_s}$, $m \leq m_s$, $1 < p_1, p_2 < \infty$, and $q > 0$ is given by $1/q = 1/p_1 + 1/p_1 - s/n$, then there exist $K, N \in \mathbb{N}$ such that*

$$\|T_\sigma(f, g)\|_{L^q} \lesssim \|\sigma\|_{K,N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}$$

for all $f \in L^{p_1}$ and $g \in L^{p_2}$.

We end this section by briefly featuring some remarks, motivations and applications in the next three subsections.

2.1. The operator norm, the number of derivatives, and complex interpolation of the classes of symbols. Theorems 2.2 and 2.5 state that the operator norm of T_σ , as a bounded operator from a product of Lebesgue spaces into another Lebesgue space, is controlled by $\|\sigma\|_{K,N}$ for some nonnegative integers K and N . Even though this is a consequence of the proof provided in each case, it can be shown to be a necessary condition. More precisely, we have the following result.

Lemma 2.6. *Let $0 < p \leq \infty$, $1 \leq p_1, p_2 < \infty$, $0 \leq \delta, \rho \leq 1$, and suppose T_σ is bounded from $L^{p_1} \times L^{p_2}$ into L^p for all $\sigma \in BS_{\rho,\delta}^m$. Then, there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T_\sigma\| \lesssim \|\sigma\|_{K,N} \quad \text{for all } \sigma \in BS_{\rho,\delta}^m.$$

Indeed, Lemma 2.6 is a consequence of the Closed Graph Theorem. Consider in $BS_{\rho,\delta}^m$ the topology induced by the family of norms $\{\|\cdot\|_{K,N}\}_{K,N \in \mathbb{N}_0}$, as defined above, which turns $BS_{\rho,\delta}^m$ into a Fréchet space. If T_σ is bounded from $L^{p_1} \times L^{p_2}$ into L^p for all $\sigma \in BS_{\rho,\delta}^m$, we can define the linear transformation

$$U : BS_{\rho,\delta}^m \rightarrow \mathcal{L}(L^{p_1} \times L^{p_2}, L^p), \quad U(\sigma) = T_\sigma,$$

where $\mathcal{L}(L^{p_1} \times L^{p_2}, L^p)$ denotes the quasi-Banach space (Banach space if $p \geq 1$) of all bilinear bounded operators from $L^{p_1} \times L^{p_2}$ into L^p endowed with the operator quasi-norm (norm if $p \geq 1$). If $\{(\sigma_k, T_{\sigma_k})\}_{k \in \mathbb{N}}$ is a sequence in the graph of U that converges to (σ, T) , for some $\sigma \in BS_{\rho,\delta}^m$ and $T \in \mathcal{L}(L^{p_1} \times L^{p_2}, L^p)$, then it easily follows that $T(f, g) = T_\sigma(f, g)$ for any $f, g \in \mathcal{S}(\mathbb{R}^n)$. Since T_σ and T are bilinear bounded operators from $L^{p_1} \times L^{p_2}$ into L^p , by density, we obtain that $T = T_\sigma$. Then, the graph of U is closed, and therefore, by the Closed Graph Theorem, U is continuous and the desired result follows.

In regards to the number of derivatives required for the symbols, we note that the following modified versions of the bilinear Hörmander classes can be considered: for $K, N \in \mathbb{N}_0$,

$$BS_{\rho,\delta,K,N}^m := \{\sigma(x, \xi, \eta) \in C^{K,N}(\mathbb{R}^{3n}) : \|\sigma\|_{K,N} < \infty\},$$

where $C^{K,N}(\mathbb{R}^{3n})$ means derivatives up to order K in x and up to order N in ξ and η . Then, $BS_{\rho,\delta,K,N}^m$ is a Banach space with norm $\|\cdot\|_{K,N}$ that contains $BS_{\rho,\delta}^m$ as a dense subset, and therefore the results of Theorem 2.2, 2.4, and 2.5 remain true if $BS_{\rho,\delta}^m$ is replaced with $BS_{\rho,\delta,K,N}^m$ for appropriate values of $K, N \in \mathbb{N}_0$, possibly depending on m, ρ and δ . We will not pursue in this paper the question regarding the minimum number of derivatives needed to achieve the results presented, though some estimates can be inferred from the proofs.

We close this subsection with a result on the complex interpolation of the classes $BS_{\rho,\rho,N,N}^m$ which will be useful in the proof of Theorem 2.2.

Lemma 2.7. *If $m_0, m_1 \in \mathbb{R}$, $0 \leq \rho < 1$, and $m = \theta m_0 + (1 - \theta)m_1$ for some $\theta \in (0, 1)$, then*

$$(BS_{\rho, \rho, N, N}^{m_0}, BS_{\rho, \rho, N, N}^{m_1})[\theta] = BS_{\rho, \rho, N, N}^m.$$

Indeed, the lemma follows by the same arguments as in the work of Päivärinta-Somersalo [27, Lemma 3.1], where the analogous result for the linear Hörmander classes is proved.

2.2. Leibniz-type rules. In terms of applications of the bilinear L^p -theory for the class $BS_{\rho, \delta}^m$, the results in this paper allow for enriched versions of the fractional Leibniz rule

$$(2.1) \quad \|fg\|_{W^{s,p}} \leq C(\|f\|_{W^{s,p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{p_1}} \|g\|_{W^{s,p_2}}),$$

where $s \geq 0$, $1/p = 1/p_1 + 1/p_2$, and $1 < p_1, p_2 < \infty$ (see Kato-Ponce [18], Christ-Weinstein [12], and Kenig-Ponce-Vega [19]).

Inequalities of the type (2.1) for pseudodifferential operators $T_\sigma(f, g)$ instead of the product fg ($\sigma \equiv 1$) can be easily obtained by following what is by now a well-known procedure that uses results going back to Coifman and Meyer and has become part of the folklore in the subject. The idea, as already used in [18], is to (smoothly) split the symbol into frequency regions where the derivatives can be distributed among the functions. (See also Semmes [28] and Gulisashvili-Kon [16] where both homogeneous and inhomogeneous derivatives were considered in similar fashion.)

Consider $\sigma \in BS_{\rho, \delta}^m$ and $\phi \in C^\infty(\mathbb{R})$ such that $0 \leq \phi \leq 1$, $\text{supp}(\phi) \subset [-2, 2]$, and $\phi(r) + \phi(1/r) = 1$ on $[0, \infty)$. For $s > 0$, the symbols σ_1 and σ_2 given by

$$\begin{aligned} \sigma_1(x, \xi, \eta) &= \sigma(x, \xi, \eta) \phi\left(\frac{1 + |\eta|^2}{1 + |\xi|^2}\right) (1 + |\xi|^2)^{-(m+s)/2}, \\ \sigma_2(x, \xi, \eta) &= \sigma(x, \xi, \eta) \phi\left(\frac{1 + |\xi|^2}{1 + |\eta|^2}\right) (1 + |\eta|^2)^{-(m+s)/2}, \end{aligned}$$

satisfy $\sigma_1, \sigma_2 \in BS_{\rho, \delta}^{-s}$, and the corresponding operators T_σ, T_{σ_1} and T_{σ_2} are related through

$$T_\sigma(f, g) = T_{\sigma_1}(J^{m+s}f, g) + T_{\sigma_2}(f, J^{m+s}g),$$

where J^{m+s} denotes the linear Fourier multiplier with symbol $(1 + |\cdot|^2)^{(m+s)/2}$. Thus, the boundedness properties on Lebesgue spaces of bilinear pseudodifferential operators given in Theorems 2.2 and 2.5 imply

$$(2.2) \quad \|T_\sigma(f, g)\|_{L^p} \leq C(\|f\|_{W^{m+s,p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{q_1}} \|g\|_{W^{m+s,q_2}}), \quad f, g \in S,$$

for appropriate values of p_1, p_2, q_1, q_2 , and s . We refer the reader to Bernicot *et al* [8] for additional Leibniz-type rules.

In the same spirit, using the functional rule

$$\partial_{x_i} T\sigma(f, g) = T_{\partial_{x_i}\sigma}(f, g) + T\sigma(\partial_{x_i}f, g) + T\sigma(f, \partial_{x_i}g),$$

the fact that $\sigma \in BS_{\rho,\delta}^m$ yields $\partial_{x_i}\sigma \in BS_{\rho,\delta}^{m+\delta}$, and bilinear complex interpolation, Theorem 2.2, and Theorem 2.5, all imply the following corollaries.

Corollary 2.8. *Let $0 \leq \delta \leq \rho \leq 1, \delta < 1, 1 \leq p_1, p_2 \leq \infty, p$ given by $1/p = 1/p_1 + 1/p_2$ and $m(p_1, p_2)$ as in Theorem 2.2. If $\sigma \in BS_{\rho,\delta}^m, m < m(p_1, p_2) - k\delta$ for some nonnegative integer k , and $r \in [0, k]$, then there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T\sigma(f, g)\|_{W^{r,p}} \lesssim \|\sigma\|_{K,N} \|f\|_{W^{r,p_1}} \|g\|_{W^{r,p_2}},$$

for all $f \in W^{r,p_1}$ and $g \in W^{r,p_2}$.

Corollary 2.9. *Let $0 \leq \delta \leq 1, 0 < \rho \leq 1, s \in (0, 2n), m_s = 2n(\rho - 1) - \rho$ as in Theorem 2.5, $1 < p_1, p_2 < \infty$, and $q > 0$ such that $1/q = 1/p_1 + 1/p_2 - s/n$. If $\sigma \in BS_{\rho,\delta}^m, m \leq m_s - k\delta$, for some nonnegative integer k , and $r \in [0, k]$, then there exist $K, N \in \mathbb{N}_0$ such that*

$$\|T\sigma(f, g)\|_{W^{r,q}} \lesssim \|\sigma\|_{K,N} \|f\|_{W^{r,p_1}} \|g\|_{W^{r,p_2}},$$

for all $f \in W^{r,p_1}$ and $g \in W^{r,p_2}$.

2.3. Applications to the scattering of PDEs. Consider the system of partial differential equations for $u = u(t, x), v = v(t, x)$, and $w = w(t, x), t \in \mathbb{R}, x \in \mathbb{R}^n$,

$$(2.3) \quad \begin{cases} \partial_t u + a(D)u = vw, & u(0, x) = 0, \\ \partial_t v + b(D)v = 0, & v(0, x) = f(x), \\ \partial_t w + c(D)w = 0, & w(0, x) = g(x), \end{cases}$$

where $a(D), b(D)$, and $c(D)$ are linear multipliers with symbols $a(\xi), b(\xi)$, and $c(\xi), \xi \in \mathbb{R}^n$, respectively. Then, formally,

$$\begin{aligned} v(t, x) &= \int_{\mathbb{R}^n} e^{-tb(\xi)} \hat{f}(\xi) e^{ix \cdot \xi} d\xi, \\ w(t, x) &= \int_{\mathbb{R}^n} e^{-tc(\eta)} \hat{g}(\eta) e^{ix \cdot \eta} d\eta, \end{aligned}$$

and

$$v(t, x)w(t, x) = \int_{\mathbb{R}^{2n}} e^{-t(b(\xi)+c(\eta))} \hat{f}(\xi)\hat{g}(\eta) e^{ix \cdot (\xi+\eta)} d\xi d\eta.$$

Another formal computation then yields $u(t, x) = (e^{-ta(D)}F(t, \cdot))(x)$, where

$$\begin{aligned} F(t, x) &= \int_0^t e^{sa(D)}(v(s, \cdot)w(s, \cdot))(x) \, ds \\ &= \int_{\mathbb{R}^{2n}} \left(\int_0^t e^{s(a(\xi+\eta)-b(\xi)-c(\eta))} \, ds \right) \hat{f}(\xi)\hat{g}(\eta)e^{ix \cdot (\xi+\eta)} \, d\xi \, d\eta. \end{aligned}$$

Therefore, if the phase function $\lambda(\xi, \eta) := a(\xi + \eta) - b(\xi) - c(\eta)$ does not vanish, we have

$$F(t, x) = T_{(e^{t\lambda})/\lambda}(f, g)(x).$$

As a consequence, assuming that $\lambda < 0$, the solution u of (2.3) scatters in the Sobolev space $W^{r,p}$ if

$$\lim_{t \rightarrow \infty} T_{(e^{t\lambda})/\lambda}(f, g) = T_{-\lambda^{-1}}(f, g) \in W^{r,p}.$$

According to Corollary 2.8, $T_{-\lambda^{-1}}$ is a bounded operator on Sobolev spaces if $-\lambda^{-1}$ belongs to $BS_{\rho,\delta}^m$ for suitable exponents. As an example, consider $b(D) = 1 - \Delta$ and $c(D) = |D|$. Then, for $a(D) = 0$, we get

$$-\lambda(\xi, \eta)^{-1} = (1 + |\xi|^2 + |\eta|)^{-1} \quad \text{and} \quad \lambda(\xi, \eta)^{-1}\varphi(\xi, \eta) \in BS_{1/2,0}^{-1},$$

for any smooth function φ such that $\varphi = 1$ away from the set $\{(\xi, \eta) : \eta = 0\}$. In the case that $a(D) = \Delta$, we get

$$-\lambda(\xi, \eta)^{-1} = (1 + |\xi + \eta|^2 + |\xi|^2 + |\eta|)^{-1} \quad \text{and} \quad \lambda(\xi, \eta)^{-1}\varphi(\xi, \eta) \in BS_{1,0}^{-2}.$$

When the phase function λ vanishes, the situation is more difficult. We refer the reader to [6, 7], where a more precise study has been developed to obtain bilinear dispersive estimates (instead of scattering properties).

3. PROOF OF THEOREM 2.1

As we will show, Theorem 2.1 follows from the case corresponding to $\rho = \delta = 0$, a scaling argument, and Lemma 2.6. We first need to recall the following result.

Theorem B (Bényi-Torres [4, Proposition 1]). *There exist x -independent symbols in $BS_{0,0}^0$ that give rise to unbounded operators from $L^{p_1} \times L^{p_2}$ into L^p for $1 \leq p_1, p_2, p < \infty$, $1/p = 1/p_1 + 1/p_2$.*

Proof of Theorem 2.1. Fix $\delta, \rho, p_1, p_2, p$ as in the hypothesis. Suppose, on the contrary, that T_σ is bounded from $L^{p_1} \times L^{p_2}$ into L^p for all $\sigma \in BS_{\rho,\delta}^0$.

Consider an x -independent symbol $\sigma \in BS_{\rho,\delta}^0$ and, for multi-indices β, γ , set

$$C_{\beta,\gamma}(\sigma) := \sup_{\xi, \eta \in \mathbb{R}^n} |\partial_\xi^\beta \partial_\eta^\gamma \sigma(\xi, \eta)| (1 + |\xi| + |\eta|)^{\rho(|\beta| + |\gamma|)}.$$

For $\lambda > 0$, define $\sigma_\lambda(\xi, \eta) := \sigma(\lambda\xi, \lambda\eta)$, $\xi, \eta \in \mathbb{R}^n$. Then, for all multi-indices β, γ and $0 < \lambda < 1$, we have

$$\begin{aligned} |\partial_\xi^\beta \partial_\eta^\gamma \sigma_\lambda(\xi, \eta)| &= \lambda^{|\beta|+|\gamma|} |\partial_\xi^\beta \partial_\eta^\gamma \sigma(\lambda\xi, \lambda\eta)| \\ &\leq \lambda^{(1-\rho)(|\beta|+|\gamma|)} C_{\beta,\gamma}(\sigma)(1 + |\xi| + |\eta|)^{-\rho(|\beta|+|\gamma|)}, \end{aligned}$$

giving

$$(3.1) \quad C_{\beta,\gamma}(\sigma_\lambda) \leq \lambda^{(1-\rho)(|\beta|+|\gamma|)} C_{\beta,\gamma}(\sigma).$$

Let $f, g \in S$, and define $f_\lambda(x) := f(x/\lambda)$ and $g_\lambda(x) := g(x/\lambda)$, $x \in \mathbb{R}^n$. Then,

$$\begin{aligned} T_\sigma(f, g)(x) &= \int_{\mathbb{R}^{2n}} \sigma(\xi, \eta) \hat{f}(\xi) \hat{g}(\eta) e^{ix \cdot (\xi + \eta)} \, d\xi \, d\eta \\ &= \int_{\mathbb{R}^{2n}} \sigma\left(\lambda \frac{\xi}{\lambda}, \lambda \frac{\eta}{\lambda}\right) \hat{f}\left(\lambda \frac{\xi}{\lambda}\right) \hat{g}\left(\lambda \frac{\eta}{\lambda}\right) e^{i\lambda x \cdot (\xi/\lambda + \eta/\lambda)} \, d\xi \, d\eta \\ &= \int_{\mathbb{R}^{2n}} \sigma_\lambda(\xi, \eta) \widehat{f_\lambda}(\xi) \widehat{g_\lambda}(\eta) e^{i\lambda x \cdot (\xi + \eta)} \, d\xi \, d\eta = T_{\sigma_\lambda}(f_\lambda, g_\lambda)(\lambda x). \end{aligned}$$

Let $K, N \in \mathbb{N}_0$ be given by Lemma 2.6 for the class $BS_{\rho,\delta}^0$, and without loss of generality, assume $K = N$. Then, using that $\|\sigma_\lambda\|_{N,N} = (\sup_{|\beta|, |\gamma| \leq N} C_{\beta,\gamma}(\sigma_\lambda))$, $1/p = 1/p_1 + 1/p_2$, and (3.1), we obtain

$$\begin{aligned} \|T_\sigma(f, g)\|_{L^p} &= \|T_{\sigma_\lambda}(f_\lambda, g_\lambda)(\lambda \cdot)\|_{L^p} = \lambda^{-n/p} \|T_{\sigma_\lambda}(f_\lambda, g_\lambda)\|_{L^p} \\ &\leq \lambda^{-n/p} \left(\sup_{|\beta|, |\gamma| \leq N} C_{\beta,\gamma}(\sigma_\lambda) \right) \|f_\lambda\|_{L^{p_1}} \|g_\lambda\|_{L^{p_2}} \\ &= \lambda^{-n/p+n/p_1+n/p_2} \left(\sup_{|\beta|, |\gamma| \leq N} C_{\beta,\gamma}(\sigma_\lambda) \right) \|f\|_{L^{p_1}} \|g\|_{L^{p_2}} \\ &\leq \left(\sup_{|\beta|, |\gamma| \leq N} \lambda^{(1-\rho)(|\beta|+|\gamma|)} C_{\beta,\gamma}(\sigma) \right) \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}, \end{aligned}$$

and letting $\lambda \rightarrow 0$, it follows that

$$(3.2) \quad \|T_\sigma(f, g)\|_{L^p} \lesssim C_{0,0}(\sigma) \|f\|_{L^{p_1}} \|g\|_{L^{p_2}} \quad f \in L^{p_1}, g \in L^{p_2}.$$

However, (3.2) cannot be true, since this contradicts Theorem B. Indeed, take $\sigma \in BS_{0,0}^0$ x -independent such that T_σ is not bounded from $L^{p_1} \times L^{p_2}$ into L^p , and let φ be an infinitely differentiable function in \mathbb{R}^{2n} supported in $|\xi, \eta| \leq 2$ and equal to one on $|\xi, \eta| \leq 1$. For each $\varepsilon > 0$, set $\sigma_\varepsilon(\xi, \eta) := \varphi(\varepsilon\xi, \varepsilon\eta)\sigma(\xi, \eta)$. Then, $\sigma_\varepsilon \in BS_{\rho,\delta}^0(\mathbb{R}^n)$ and $C_{0,0}(\sigma_\varepsilon) \leq C_{0,0}(\sigma)$ for all $\varepsilon > 0$. If (3.2) were true, we would have

$$\|T_{\sigma_\varepsilon}(f, g)\|_{L^p} \lesssim C_{0,0}(\sigma) \|f\|_{L^{p_1}} \|g\|_{L^{p_2}} \quad f, g \in S, \text{ for all } \varepsilon > 0.$$

As $\varepsilon \rightarrow 0$, $T_{\sigma_\varepsilon}(f, g) \rightarrow T_\sigma(f, g)$ pointwise; this and Fatou's Lemma yield

$$\|T_\sigma(f, g)\|_{L^p} \lesssim C_{0,0}(\sigma) \|f\|_{L^{p_1}} \|g\|_{L^{p_2}} \quad f, g \in S,$$

a contradiction. □

4. PROOF OF THEOREM 2.2

4.1. Preliminary results. We will use the following results in the proof of Theorem 2.2.

Theorem C (Symbolic calculus, Bényi-Maldonado-Naibo-Torres [2]). Assume that $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$, and $\sigma \in BS_{\rho,\delta}^m$. Then, for $j = 1, 2$, we have $T_\sigma^{*j} = T_{\sigma^{*j}}$, where $\sigma^{*j} \in BS_{\rho,\delta}^m$. Moreover, if $0 \leq \delta < \rho \leq 1$ and $\sigma \in BS_{\rho,\delta}^m$, then σ^{*1} and σ^{*2} have explicit asymptotic expansions.

Theorem D (Michalowski-Rule-Staubach [24, Theorem 5.5]). Let $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$, $1 \leq p_1, p_2, p \leq \infty$, $1/p = 1/p_1 + 1/p_2$, and

$$m < n(\rho - 1) \max \left\{ \frac{1}{2}, \left(\frac{2}{p_1} - \frac{1}{2} \right), \left(\frac{2}{p_2} - \frac{1}{2} \right), \left(\frac{3}{2} - \frac{2}{p} \right) \right\}.$$

If $\sigma \in BS_{\rho,\delta}^m$, then there exist $K, N \in \mathbb{N}_0$ such that

$$\|T_\sigma(f, g)\|_{L^p} \lesssim \|\sigma\|_{K,N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}.$$

Set $\tilde{m}(p_1, p_2) = n(\rho - 1) \max \{ \frac{1}{2}, (2/p_1 - \frac{1}{2}), (2/p_2 - \frac{1}{2}), (\frac{3}{2} - 2/p) \}$, and note that, when $p > 1$, we have $m(p_1, p_2) = n(\rho - 1) \max \{ \frac{1}{2}, 1/p_1, 1/p_2, (1 - 1/p) \}$. Referring to Figure 4.1, we then have that $m(p_1, p_2) = n(\rho - 1)(1/p_2)$ and $\tilde{m}(p_1, p_2) = n(\rho - 1)(2/p_2 - \frac{1}{2})$ in region I, $m(p_1, p_2) = n(\rho - 1)(1/p_1)$ and $\tilde{m}(p_1, p_2) = n(\rho - 1)(2/p_1 - \frac{1}{2})$ in region II, $m(p_1, p_2) = n(\rho - 1)(1 - 1/p)$ and $\tilde{m}(p_1, p_2) = n(\rho - 1)(\frac{3}{2} - 2/p)$ in region III, and $m(p_1, p_2) = \tilde{m}(p_1, p_2) = n(\rho - 1)\frac{1}{2}$ in region IV. Then, $\tilde{m} < m$ in regions I, II, and III, and therefore the Banach case of Theorem 2.2 is an improvement on Theorem D.

In the non-Banach case ($p < 1$), we will use bilinear Calderón-Zygmund theory to get the boundedness results stated in Theorem 2.2. Indeed, we have the following result.

Theorem 4.1 (Bilinear Calderón-Zygmund operators). Let $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$, $0 < \rho$, and set $m_{cz} := 2n(\rho - 1)$. If $\sigma \in BS_{\rho,\delta}^m$ and $m < m_{cz}$, then T_σ is a bilinear Calderón-Zygmund operator. As a consequence, the following mapping properties hold true for $1 \leq p_1, p_2 \leq \infty$, $\frac{1}{2} \leq p < \infty$, $1/p = 1/p_1 + 1/p_2$:

- (i) If $1 < p_1, p_2$, then there exist $K, N \in \mathbb{N}_0$ such that

$$\|T_\sigma(f, g)\|_{L^p} \lesssim \|\sigma\|_{K,N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}},$$

where L^{p_1} or L^{p_2} should be replaced by L_c^∞ (bounded functions with compact support) if $p_1 = \infty$ or $p_2 = \infty$, respectively;

(ii) If $p_1 = 1$ or $p_2 = 1$, then there exist $K, N \in \mathbb{N}$ such that

$$\|T_\sigma(f, g)\|_{L^{p_1, \infty}} \lesssim \|\sigma\|_{K, N} \|f\|_{L^{p_1}} \|g\|_{L^{p_2}},$$

where L^{p_1} or L^{p_2} should be replaced by L_c^∞ if $p_1 = \infty$ or $p_2 = \infty$, respectively;

(iii) There exist $K, N \in \mathbb{N}$ such that

$$\|T_\sigma(f, g)\|_{\text{BMO}} \lesssim \|\sigma\|_{K, N} \|f\|_{L^\infty} \|g\|_{L^\infty}$$

for $f, g \in L_c^\infty$;

(iv) Weighted versions of the above inequalities hold (see Section 9).

The results of Theorem 4.1 are consequences of the following estimates for the kernel of T_σ .

Theorem E. Let $\sigma \in BS_{\rho, \delta}^m$, $0 < \rho \leq 1$, $0 \leq \delta < 1$, $m \in \mathbb{R}$, and denote by $\mathcal{K}(x, y, z)$ the distributional kernel of the associated bilinear pseudodifferential operator T_σ . For $x, y, z \in \mathbb{R}^n$, set $S(x, y, z) = |x - y| + |x - z| + |y - z|$. Then,

(i) Given $\alpha, \beta, \gamma \in \mathbb{N}_0^n$, there exists $N_0 \in \mathbb{N}_0$ such that, for each $\ell \geq N_0$,

$$\sup_{(x, y, z): S(x, y, z) > 0} S(x, y, z)^\ell |D_x^\alpha D_y^\beta D_z^\gamma \mathcal{K}(x, y, z)| < \infty;$$

(ii) Suppose that σ has compact support in (ξ, η) uniformly in x . Then, \mathcal{K} is smooth, and, given $\alpha, \beta, \gamma \in \mathbb{N}_0^n$ and $N_0 \in \mathbb{N}_0$, there exists $C > 0$ such that, for all $x, y, z \in \mathbb{R}^n$ with $S(x, y, z) > 0$, we have

$$|D_x^\alpha D_y^\beta D_z^\gamma \mathcal{K}(x, y, z)| \leq C(1 + S(x, y, z))^{-N_0};$$

(iii) Suppose that $m + M + 2n < 0$ for some $M \in \mathbb{N}_0$. Then, \mathcal{K} is a bounded continuous function with bounded continuous derivatives of order $\leq M$;

(iv) Suppose that $m + M + 2n = 0$ for some $M \in \mathbb{N}_0$. Then, there exists a constant $C > 0$ such that, for all $x, y, z \in \mathbb{R}^n$ with $S(x, y, z) > 0$,

$$\sup_{|\alpha + \beta + \gamma| = M} |D_x^\alpha D_y^\beta D_z^\gamma \mathcal{K}(x, y, z)| \leq C |\log |S(x, y, z)||;$$

(v) Suppose that $m + M + 2n > 0$ for some $M \in \mathbb{N}_0$. Then, given $\alpha, \beta, \gamma \in \mathbb{N}_0^n$, there exists a positive constant C such that, for all $x, y, z \in \mathbb{R}^n$ with $S(x, y, z) > 0$,

$$\sup_{|\alpha + \beta + \gamma| = M} |\partial_x^\alpha \partial_y^\beta \partial_z^\gamma \mathcal{K}(x, y, z)| \leq CS(x, y, z)^{-(m + M + 2n)/\rho};$$

- (vi) Suppose that $m + \varepsilon + 2n > 0$ for some $\varepsilon \in (0, 1)$. Then, there exists a positive constant C such that, for all $x, y, z, u \in \mathbb{R}^n$ with $S(x, y, z) > 0$ and $|u| \leq S(x, y, z)$,

$$|\mathcal{K}(x, y, z) - \mathcal{K}(x + u, y, z)| + |\mathcal{K}(x, y, z) - \mathcal{K}(x, y + u, z)| \\ + |\mathcal{K}(x, y, z) - \mathcal{K}(x, y, z + u)| \leq C|u|^\varepsilon S(x, y, z)^{-(m+\varepsilon+2n)/\rho}.$$

All constants in the above inequalities depend linearly on $\|\sigma\|_{K,N}$ for some $K, N \in \mathbb{N}_0$.

We refer the reader to [2, Theorem 6] for the proofs of items (i)–(v) in Theorem E. Item (vi) corresponds to the “Hölder” version of item (v); also, its proof is analogous, and relies on estimates for linear kernels as presented in Alvarez-Hounie [1, Theorem 1.1].

Proof of Theorem 4.1. It is enough to prove the result for $\sigma \in BS_{\rho,\delta}^m$ and m such that $2n(\rho - 1) - t < m < 2n(\rho - 1) = m_{cz}$ for some small positive number t . Denote by $\mathcal{K}(x, y, z)$ the distributional kernel of the associated bilinear pseudodifferential operator T_σ . Using that $BS_{\rho,\delta}^m \subset BS_{\rho,\delta}^{m_{cz}}$, we have that part (v) of Theorem E applied to $BS_{\rho,\delta}^{m_{cz}}$ yields, with constants depending linearly on $\|\sigma\|_{N,N}$ for some $N \in \mathbb{N}_0$, that

$$|\mathcal{K}(x, y, z)| \lesssim \frac{1}{(|x - y| + |x - z| + |y - z|)^{2n}},$$

while part (vi) gives, again with constants depending linearly on $\|\sigma\|_{N,N}$ for some $N \in \mathbb{N}_0$, that

$$|\mathcal{K}(x, y, z) - \mathcal{K}(x + u, y, z)| + |\mathcal{K}(x, y, z) - \mathcal{K}(x, y + u, z)| \\ + |\mathcal{K}(x, y, z) - \mathcal{K}(x, y, z + u)| \lesssim \frac{|u|^\varepsilon}{(|x - y| + |x - z| + |y - z|)^{2n+\varepsilon}},$$

where $|u| \leq |x - y| + |x - z| + |y - z|$ and $\varepsilon \in (0, 1)$ has been chosen such that $(m + 2n + \varepsilon)/\rho = 2n + \varepsilon$ (which is possible since $2n(\rho - 1) - t < m < 2n(\rho - 1)$ for small enough $t > 0$). Moreover, since $m < m_{cz} < n(\rho - 1)/2$, Theorem D yields that there exists $N \in \mathbb{N}_0$ such that T_σ satisfies

$$\|T_\sigma(f, g)\|_{L^1} \lesssim \|\sigma\|_{N,N} \|f\|_{L^2} \|g\|_{L^2}.$$

We then conclude that T_σ is a bilinear Calderón-Zygmund operator for which the corresponding boundedness properties follow (see [15]). \square

4.2. Proof of Theorem 2.2. With these preliminary and technical results, we are now ready for the proof of our main result in this section.

Proof of Theorem 2.2. We first prove the theorem for $p_1 = p_2 = p = \infty$, in which case $m(p_1, p_2) = n(\rho - 1)$. Let $m < n(\rho - 1)$, $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$. Let $\{\psi_j\}_{j \in \mathbb{N}_0}$ be a partition of unity on \mathbb{R}^{2n} :

$$\sum_{j=0}^{\infty} \psi_j(\xi, \eta) = 1, \quad \xi, \eta \in \mathbb{R}^n,$$

such that ψ_0 is supported in $\{(\xi, \eta) \in \mathbb{R}^{2n} : |(\xi, \eta)| \leq 2\}$ and $\psi_j(\xi, \eta) = \psi(2^{-j}\xi, 2^{-j}\eta)$, for some $\psi \in C_0^\infty(\mathbb{R}^{2n})$ supported in $\{(\xi, \eta) \in \mathbb{R}^{2n} : |(\xi, \eta)| \sim 1\}$ for $j \in \mathbb{N}$. We decompose the symbol $\sigma(x, \xi, \eta)$ as

$$\sigma(x, \xi, \eta) = \sum_{j=0}^{\infty} \sigma_j(x, \xi, \eta),$$

where $\sigma_j(x, \xi, \eta) := \sigma(x, \xi, \eta)\psi_j(\xi, \eta)$. Then, $\|\sigma_j\|_{0,N} \lesssim \|\sigma\|_{0,N}$ for all $N \in \mathbb{N}_0$, and, by Lemma 6.1 (see Section 6),

$$\|T_{\sigma_j}(f, g)\|_{L^\infty} \lesssim \|\sigma\|_{0,2N} 2^{j(m+n(1-\rho))} \|f\|_{L^\infty} \|g\|_{L^\infty},$$

where $j \in \mathbb{N}_0$, $N > n$, $K \in \mathbb{N}_0$. Therefore, we have

$$\begin{aligned} \|T_\sigma(f, g)\|_{L^\infty} &\leq \sum_{j=0}^{\infty} \|T_{\sigma_j}(f, g)\|_{L^\infty} \lesssim \|\sigma\|_{0,2N} \sum_{j=0}^{\infty} 2^{j(m+n(1-\rho))} \|f\|_{L^\infty} \|g\|_{L^\infty} \\ &\lesssim \|\sigma\|_{0,2N} \|f\|_{L^\infty} \|g\|_{L^\infty}, \end{aligned}$$

where we have used that $m < n(\rho - 1)$. This proves the theorem for $p_1 = p_2 = p = \infty$. Note that the proof shows that there is an extension of T_σ that is bounded from $L^\infty \times L^\infty$ into L^∞ , namely

$$T_\sigma(f, g) = \sum_{j=0}^{\infty} T_{\sigma_j}(f, g)$$

where

$$T_{\sigma_j}(f, g)(x) = \int_{\mathbb{R}^{2n}} \mathcal{K}_j(x, x - y, x - z) f(y) g(z) \, dy \, dz,$$

with

$$\mathcal{K}_j(x, y, z) = \int_{\mathbb{R}^{2n}} \sigma_j(x, \xi, \eta) e^{i\xi \cdot y} e^{i\eta \cdot z} \, d\xi \, d\eta, \quad x, y, z \in \mathbb{R}^n.$$

We now proceed to prove the theorem in the general case. We recall that the boundedness properties in Lebesgue spaces for operators corresponding to the class $BS_{1,\delta}^0$ for $0 \leq \delta < 1$ are well known (see the [Introduction](#)); therefore, we

will work with $\rho < 1$. Moreover, since $BS_{\rho,\delta}^m \subset BS_{\rho,\rho}^m$ for $\delta \leq \rho$, we will assume $\delta = \rho$, $0 \leq \rho < 1$. Define on $BS_{\rho,\rho}^m \times L^{p_1} \times L^{p_2}$ the operator T given by

$$T(\sigma, f, g) := T_\sigma(f, g).$$

In the following, we use the notation $T : BS_{\rho,\rho}^m \times X \times Y \rightarrow Z$ to express the fact that T maps continuously from $BS_{\rho,\rho}^m \times X \times Y$ into Z : there exists $N \in \mathbb{N}_0$, possibly depending on m and ρ , such that $\|T(\sigma, f, g)\|_Z \lesssim \|\sigma\|_{N,N} \|f\|_X \|g\|_Y$, for all $\sigma \in BS_{\rho,\rho}^m$, $f \in X$, $g \in Y$.




FIGURE 4.1. Value of $m(p_1, p_2)$ as given by Theorem 2.2

We first prove (i) (case $p > 1$). The case $p_1 = p_2 = p = \infty$ proved above, and Theorem C, yield

- $T : BS_{\rho,\rho}^m \times L^\infty \times L^\infty \rightarrow L^\infty$ for $m < n(\rho - 1)$ (point $(0, 0)$ in Figure 4.1),
- $T : BS_{\rho,\rho}^m \times L^1 \times L^\infty \rightarrow L^1$ for $m < n(\rho - 1)$ (point $(1, 0)$ in Figure 4.1),
- $T : BS_{\rho,\rho}^m \times L^\infty \times L^1 \rightarrow L^1$ for $m < n(\rho - 1)$ (point $(0, 1)$ in Figure 4.1).

Moreover, by Theorem D we have

- $T : BS_{\rho,\rho}^m \times L^2 \times L^2 \rightarrow L^1$ for $m < (n/2)(\rho - 1)$ (point $(\frac{1}{2}, \frac{1}{2})$ in Figure 4.1),
- $T : BS_{\rho,\rho}^m \times L^2 \times L^\infty \rightarrow L^2$ for $m < (n/2)(\rho - 1)$ (point $(\frac{1}{2}, 0)$ in Figure 4.1),
- $T : BS_{\rho,\rho}^m \times L^\infty \times L^2 \rightarrow L^2$ for $m < (n/2)(\rho - 1)$ (point $(0, \frac{1}{2})$ in Figure 4.1).

We now recall the following modified version of the bilinear Hörmander classes (see Section 2.1):

$$BS_{\rho,\rho,N,N}^m := \{ \sigma(x, \xi, \eta) \in C^N(\mathbb{R}^{3n}) : \|\sigma\|_{N,N} < \infty \}$$

where $N \in \mathbb{N}_0$ and, as always,

$$\|\sigma\|_{N,N} := \sup_{|\alpha|, |\beta|, |\gamma| \leq N} \sup_{x, \xi, \eta \in \mathbb{R}^n} |\partial_x^\alpha \partial_\xi^\beta \partial_\eta^\gamma \sigma(x, \xi, \eta)| (1 + |\xi| + |\eta|)^{-m - \rho(|\alpha| + \rho(|\beta| + |\gamma|))}.$$

Since $BS_{\rho,\rho}^m$ is dense in $BS_{\rho,\rho,N,N}^m$, the above mentioned endpoint results also hold if $BS_{\rho,\rho}^m$ is replaced with $BS_{\rho,\rho,N,N}^m$ for large enough N , possibly depending on ρ and m . Lemma 2.7 and trilinear complex interpolation (see the book of Bergh and Löfström [5, Theorem 4.4.1]) then yield the thesis of the theorem for p_1 and p_2 such that $(1/p_1, 1/p_2)$ is on the border of the triangle with vertices $(0, 0)$, $(0, 1)$, and $(1, 0)$.

The result for p_1 and p_2 such that $(1/p_1, 1/p_2)$ is in the interior of the triangle follows by bilinear complex interpolation since, as shown in Figure 4.1, $m(p_1, p_2)$ is constant along horizontal segments in region I, $m(p_1, p_2)$ is constant along vertical segments in region II, $m(p_1, p_2)$ is constant along diagonal segments in region III, and $m(p_1, p_2)$ is constant in region IV.

We now prove (ii) and (iii) (case $p < 1$). Here, we must assume $\rho > 0$. Theorem 4.1 yields

$$T : BS_{\rho,\rho}^m \times L^1 \times L^1 \rightarrow L^{1/2,\infty}, \text{ for } m < 2n(\rho - 1) \text{ (point } (1, 1) \text{ in Figure 4.1),}$$

which, together with the boundedness properties at the points $(1, 0)$ and $(0, 1)$ in Figure 4.1 (as stated above), Lemma 2.7, and trilinear complex interpolation, gives that

$$T : BS_{\rho,\rho}^m \times L^{p_1} \times L^{p_2} \rightarrow L^{p,\infty}, \quad m < m(p_1, p_2),$$

for $(1/p_1, 1/p_2)$ on the segments joining the points $(0, 1)$ to $(1, 1)$, $(1, 0)$ to $(1, 1)$, and $(\frac{1}{2}, \frac{1}{2})$ to $(1, 1)$, in Figure 4.1. This gives Part (iii).

For Part (ii), consider the shaded triangle as indicated in each case presented in Figure 4.2. The value $m(p_1, p_2)$ is constant (say, m) on the upper border of this triangle, which is given by two segments with equations

$$m = n(\rho - 1) \left(\frac{2}{p_1} + \frac{1}{p_2} - 1 \right)$$

(inside the triangle with vertices $(1, 1)$, $(\frac{1}{2}, \frac{1}{2})$, and $(1, 0)$) and

$$m = n(\rho - 1) \left(\frac{2}{p_2} + \frac{1}{p_1} - 1 \right)$$

(inside the triangle with vertices $(1, 1)$, $(\frac{1}{2}, \frac{1}{2})$, and $(0, 1)$). Then, Part (ii) follows by bilinear real interpolation, by using the weak-type estimates obtained above for the vertices of the shaded triangle. \square




FIGURE 4.2. Case $p < 1$ of Theorem 2.2

Remark 4.2. We note that the proof of Theorem 2.2 given for the case $p_1 = p_2 = p = \infty$ does not require any assumptions on the derivatives of the symbol σ in the space variables. This particular result is included in [24, Theorem 3.3], which yields boundedness properties in Lebesgue spaces of bilinear pseudo-differential operators with rough symbols in the space variables as a consequence of $|T_\sigma(f, g)|$ being pointwise bounded in terms of the Hardy-Littlewood maximal operator evaluated at f and g . For completeness, we have provided another proof of the case $p_1 = p_2 = \infty$ of Theorem 2.2 following the arguments of the corresponding linear result in [20].

5. PROOF OF THEOREM 2.3

In this section, we continue to use L^p and $\|\cdot\|_{L^p}$ to denote the Lebesgue space $L^p(\mathbb{R}^n)$ and its norm, respectively. Sometimes it will be necessary to make explicit the variable of integration (e.g., integration with respect to x), in which case we employ the notation $\|\cdot\|_{L^p(dx)}$.

Proof of Theorem 2.3. Without loss of generality, we may assume that the symbol σ has compact support in the frequency variables ξ and η . Otherwise, define $\sigma_\varepsilon(x, \xi, \eta) := \varphi(\varepsilon \xi, \varepsilon \eta) \sigma(x, \xi, \eta)$, where φ is a smooth function compactly

supported in $B(0, 1)$ such that $0 \leq \varphi \leq 1$ and $\varphi(0, 0) = 1$. It easily follows that $C(\sigma_\varepsilon) \lesssim C(\sigma)$ and that

$$\lim_{\varepsilon \rightarrow 0} T_{\sigma_\varepsilon}(f, g) = T_\sigma(f, g)$$

pointwise for f and g belonging to the class \mathcal{U} of functions whose Fourier transforms are in C_0^∞ . Assuming the result for symbols of compact support, by Fatou's lemma, we have

$$\|T_\sigma(f, g)\|_{L^2} \leq \liminf_{\varepsilon \rightarrow 0} \|T_{\sigma_\varepsilon}(f, g)\|_{L^2} \leq C(\sigma) \|f\|_{L^2} \|g\|_{L^2},$$

for $f, g \in \mathcal{U}$. Since \mathcal{U} is dense in L^2 , the desired result holds for non-compactly supported symbols as well.

Suppose first that σ is x -independent, and define $\tau(\xi, \eta) := \sigma(x, \xi, \eta)$. Then,

$$T_\tau(\widehat{f, g})(\xi) = \int_{\mathbb{R}^n} \hat{f}(\xi - \eta) \hat{g}(\eta) \tau(\xi - \eta, \eta) \, d\eta,$$

and the Cauchy-Schwarz inequality yields

$$|T_\tau(\widehat{f, g})(\xi)| \leq \left(\int_{\mathbb{R}^n} |\hat{f}(\xi - \eta)|^2 |\hat{g}(\eta)|^2 \, d\eta \right)^{1/2} \sup_{\xi \in \mathbb{R}^n} \|\tau(\xi - \cdot, \cdot)\|_{L^2}.$$

Integrating in ξ and using Plancherel's theorem, it follows that

$$(5.1) \quad \|T_\tau(f, g)\|_{L^2} \lesssim \|f\|_{L^2} \|g\|_{L^2} \sup_{\xi \in \mathbb{R}^n} \|\tau(\xi - \cdot, \cdot)\|_{L^2},$$

which implies the desired result.

Next, we continue working with an x -independent symbol $\tau(\xi, \eta)$ in order to get estimates that will be useful later for x -dependent symbols. Let Φ be a smooth function compactly supported in $B(0, \sqrt{n})$ such that $0 \leq \Phi \leq 1$ and

$$\sum_{k \in \mathbb{Z}^n} \Phi(k - x) = 1, \quad x \in \mathbb{R}^n.$$

For a function h defined in \mathbb{R}^n and $\ell \in \mathbb{Z}^n$, we set $h_\ell(x) := \Phi(x - \ell)h(x)$. We show that, for every $N \in \mathbb{N}$,

$$(5.2) \quad \|\Phi(\cdot - \ell)T_\tau(f, g)\|_{L^2} \lesssim \sup_{\xi \in \mathbb{R}^n, |\alpha| \leq 2N} \|\partial_\xi^\alpha \tau(\xi - \cdot, \cdot)\|_{L^2} \\ \times \sum_{j, k \in \mathbb{Z}^n} \frac{\|f_j\|_{L^2} \|g_k\|_{L^2}}{(1 + |\ell - j| + |\ell - k|)^N}$$

for all $\ell \in \mathbb{Z}^n$ and with constants independent of τ , ℓ , f , and g .

We have

$$\Phi(x - \ell)T_\tau(f, g)(x) = \sum_{j, k \in \mathbb{Z}^n} \Phi(x - \ell)T_\tau(f_j, g_k)(x), \quad x \in \mathbb{R}^n,$$

and therefore (5.2) will follow from the estimate

$$(5.3) \quad \|\Phi(\cdot - \ell)T_\tau(f_j, g_k)\|_{L^2} \lesssim \sup_{\xi \in \mathbb{R}^n, |\alpha| \leq 2N} \|\partial_\xi^\alpha \tau(\xi - \cdot, \cdot)\|_{L^2} \times \frac{\|f_j\|_{L^2} \|g_k\|_{L^2}}{(1 + |\ell - j| + |\ell - k|)^N}.$$

Fix $\ell \in \mathbb{Z}^n$. When $j, k \in \mathbb{Z}^n$ are such that $|\ell - j| + |\ell - k| \leq 10$, we apply (5.1):

$$\begin{aligned} \|\Phi(\cdot - \ell)T_\tau(f_j, g_k)\|_{L^2} &\leq \|T_\tau(f_j, g_k)\|_{L^2} \\ &\lesssim \sup_{\xi \in \mathbb{R}^n} \|\tau(\xi - \cdot, \cdot)\|_{L^2} \|f_j\|_{L^2} \|g_k\|_{L^2} \\ &\lesssim \sup_{\xi \in \mathbb{R}^n} \|\tau(\xi - \cdot, \cdot)\|_{L^2} \frac{\|f_j\|_{L^2} \|g_k\|_{L^2}}{(1 + |\ell - j| + |\ell - k|)^N} \end{aligned}$$

for every integer N . Therefore, (5.3) holds.

We now consider j and k such that $|\ell - j| + |\ell - k| \geq 10$, and without loss of generality, we assume that $|\ell - j| \geq |\ell - k|$. Then,

$$\begin{aligned} &T_\tau(f_j, g_k)(x) \\ &= \int_{\mathbb{R}^{2n}} \left(\int_{\mathbb{R}^{2n}} e^{i(\xi \cdot (x-y) + \eta \cdot (x-z))} \tau(\xi, \eta) \, d\xi \, d\eta \right) f_j(y) g_k(z) \, dy \, dz \\ &= \int_{\mathbb{R}^{2n}} \left(\int_{\mathbb{R}^{2n}} e^{i(\xi \cdot (x-y) + \eta \cdot (x-z))} (1 - \Delta_\xi)^N \tau(\xi, \eta) \, d\xi \, d\eta \right) \frac{f_j(y) g_k(z) \, dy \, dz}{(1 + |x - y|^2)^N} \\ &= \int_{\mathbb{R}^{2n}} \mathcal{F}_{2n}((1 - \Delta_\xi)^N \tau)(y - x, z - x) \frac{f_j(y) g_k(z) \, dy \, dz}{(1 + |x - y|^2)^N} \end{aligned}$$

where \mathcal{F}_{2n} denotes the Fourier transform in \mathbb{R}^{2n} . Multiplying by $\Phi(x - \ell)$ and using the Sobolev embedding $W^{P,2} \subset L^\infty$ for any $P > n/2$, by fixing $x \in \mathbb{R}^n$, we have that

$$\begin{aligned} &|\Phi(x - \ell)T_\tau(f_j, g_k)(x)| \\ &\lesssim \sup_{a \in \mathbb{R}^n} \left| \Phi(a - \ell) \int_{\mathbb{R}^{2n}} \mathcal{F}_{2n}((1 - \Delta_\xi)^N \tau)(y - x, z - x) \frac{f_j(y) g_k(z)}{(1 + |a - y|^2)^N} \, dy \, dz \right| \\ &\lesssim \sup_{|\beta| \leq P} \left\| \int_{\mathbb{R}^{2n}} \mathcal{F}_{2n}((1 - \Delta_\xi)^N \tau)(y - x, z - x) \right. \\ &\quad \left. \times (\partial_a^\beta \mathcal{Y}_{\ell, N})(a, y) f_j(y) g_k(z) \, dy \, dz \right\|_{L^2(da)}, \end{aligned}$$

where $\gamma_{\ell,N}(a, \gamma) := \Phi(a - \ell)/(1 + |a - \gamma|^2)^N$. Therefore,

$$\begin{aligned} & \|\Phi(\cdot - \ell)T_\tau(f_j, g_k)\|_{L^2} \\ & \lesssim \sup_{|\beta| \leq P} \left\| \int_{\mathbb{R}^{2n}} \mathcal{F}_{2n}((1 - \Delta_\xi)^N \tau)(\gamma - x, z - x) \right. \\ & \qquad \qquad \qquad \times (\partial_a^\beta \gamma_{\ell,N})(a, \gamma) f_j(\gamma) g_k(z) \, d\gamma \, dz \Big\|_{L^2(da \, dx)} \\ & = \sup_{|\beta| \leq P} \left\| T_{((1 - \Delta_\xi)^N \tau)} \left(\partial_a^\beta \left(\frac{\Phi(a - \ell)}{(1 + |a - \cdot|^2)^N} \right) f_j(\cdot), g_k(\cdot) \right) (x) \right\|_{L^2(da \, dx)}. \end{aligned}$$

Applying (5.1) to $T_{((1 - \Delta_\xi)^N \tau)}$ then yields

$$\begin{aligned} & \|\Phi(\cdot - \ell)T_\tau(f_j, g_k)\|_{L^2} \\ & \lesssim \sup_{\xi \in \mathbb{R}^n} \|(1 - \Delta_\xi)^N \tau(\xi - \cdot, \cdot)\|_{L^2} \\ & \qquad \times \sup_{|\beta| \leq P} \left\| \partial_a^\beta \left(\frac{\Phi(a - \ell)}{(1 + |a - \gamma|^2)^N} \right) f_j(\gamma) \right\|_{L^2(da \, d\gamma)} \|g_k\|_{L^2} \\ & \lesssim \sup_{\xi \in \mathbb{R}^n} \|(1 - \Delta_\xi)^N \tau(\xi - \cdot, \cdot)\|_{L^2} \frac{\|f_j\|_{L^2} \|g_k\|_{L^2}}{(1 + \|\ell - j\|^2)^N}, \end{aligned}$$

giving (5.3), where we have used that

$$\left\| \partial_a^\beta \left(\frac{\Phi(a - \ell)}{(1 + |a - \gamma|^2)^N} \right) \right\|_{L^2(da)} \lesssim \frac{1}{(1 + |\ell - j|^2)^N}, \quad \gamma \in B(j, \sqrt{n}).$$

Consider now an x -dependent symbol. Let

$$U_\gamma(f, g)(x) := \int_{\mathbb{R}^{2n}} e^{ix \cdot (\xi + \eta)} \hat{f}(\xi) \hat{g}(\eta) \sigma(\gamma, \xi, \eta) \, d\xi \, d\eta, \quad x, \gamma \in \mathbb{R}^n.$$

Then, we have $T_\sigma(f, g)(x) = U(x)(f, g)(x)$. Fixing $x \in \mathbb{R}^n$, $\ell \in \mathbb{Z}^n$, and using the Sobolev embedding $W^{s,2} \hookrightarrow L^\infty$ for an integer $s > n/2$, we get

$$\begin{aligned} |\Phi(x - \ell)T_\sigma(f, g)(x)| & \leq \sup_{\gamma \in \mathbb{R}^n} |\Phi(\gamma - \ell)U_\gamma(f, g)(x)| \\ & \leq \sum_{|\beta| \leq s} \|\partial_\gamma^\beta (\Phi(\gamma - \ell)U_\gamma(f, g)(x))\|_{L^2(d\gamma)} \\ & \lesssim \sum_{|\beta| \leq s} \|\chi_{B(\ell)}(\gamma) \partial_\gamma^\beta U_\gamma(f, g)(x)\|_{L^2(d\gamma)}, \end{aligned}$$

where $B(\ell) = B(\ell, \sqrt{n})$. Let $\tilde{\Phi}$ be a smooth function supported in $B(0, \sqrt{n})$ such that $\tilde{\Phi}\Phi = \Phi$. Multiplying by $\tilde{\Phi}(x - \ell)$, integrating in x , and using Fubini's Theorem, we obtain

$$\begin{aligned} & \|\Phi(\cdot - \ell)T_\sigma(f, g)\|_{L^2} \\ & \lesssim \sum_{|\beta| \leq s} \|\chi_{B(\ell)}(\mathcal{Y})\| \chi_{B(\ell)}(x) \tilde{\Phi}(x - \ell) \partial_y^\beta U_y(f, g)(x) \|_{L^2(dx)} \|_{L^2(d\mathcal{Y})}. \end{aligned}$$

For each $\beta \in \mathbb{N}_0^n$, $|\beta| \leq s$, and $\mathcal{Y} \in \mathbb{R}^n$, we look at $\partial_y^\beta U_y$ as the bilinear multiplier defined by the x -independent symbol $\tau_y^\beta(\xi, \eta) := \partial_y^\beta \sigma(\mathcal{Y}, \xi, \eta)$. Then, applying (5.2), which also holds if on its left-hand side Φ is replaced by $\tilde{\Phi}$, we deduce

$$\begin{aligned} & \|\Phi(\cdot - \ell)T_\sigma(f, g)\|_{L^2} \\ & \lesssim \sum_{|\beta| \leq s} \sum_{j, k \in \mathbb{Z}^n} \sup_{\xi \in \mathbb{R}^n, |\alpha| \leq 2N} \sup_{\mathcal{Y} \in \mathbb{R}^n} \|\partial_\xi^\alpha \tau_y^\beta(\xi - \cdot, \cdot)\|_{L^2} \frac{\|f_j\|_{L^2} \|g_k\|_{L^2}}{(1 + |\ell - j| + |\ell - k|)^N}, \end{aligned}$$

which implies

$$\|\Phi(\cdot - \ell)T_\sigma(f, g)\|_{L^2} \lesssim C(\sigma) \sum_{j, k \in \mathbb{Z}^n} \frac{\|f_j\|_{L^2} \|g_k\|_{L^2}}{(1 + |\ell - j| + |\ell - k|)^N},$$

with $C(\sigma) := \sup_{|\beta| \leq s, |\alpha| \leq 2N} \sup_{\xi, \mathcal{Y} \in \mathbb{R}^n} \|\partial_\xi^\alpha \partial_y^\beta \sigma(\mathcal{Y}, \xi - \cdot, \cdot)\|_{L^2}$.

Using Hölder's inequality, we then obtain that

$$\begin{aligned} \|\Phi(\cdot - \ell)T_\sigma(f, g)\|_{L^2}^2 & \lesssim C(\sigma)^2 \sum_{j, k \in \mathbb{Z}^n} \frac{\|f_j\|_{L^2}^2 \|g_k\|_{L^2}^2}{(1 + |\ell - j| + |\ell - k|)^N} \\ & \quad \times \sum_{j, k \in \mathbb{Z}^n} \frac{1}{(1 + |\ell - j| + |\ell - k|)^N}. \end{aligned}$$

Choosing $N > 2n$, the second sum on the right-hand side is finite, and after summing over $\ell \in \mathbb{Z}^n$, we conclude that

$$\begin{aligned} \sum_{\ell \in \mathbb{Z}^n} \|T_\sigma(f, g)_\ell\|_{L^2}^2 & = \sum_{\ell \in \mathbb{Z}^n} \|\Phi(\cdot - \ell)T_\sigma(f, g)\|_{L^2}^2 \\ & \lesssim C(\sigma)^2 \sum_{j \in \mathbb{Z}^n} \|f_j\|_{L^2}^2 \sum_{k \in \mathbb{Z}^n} \|g_k\|_{L^2}^2. \end{aligned}$$

The desired result follows by taking $N = 2n + 1$, $s = [n/2] + 1$, and noting that $\|h\|_{L^2}^2 \sim \sum_j \|h_j\|^2$. □

6. PROOF OF THEOREM 2.4

The following lemmas, whose proofs are included in Section 8, will be used to prove Theorem 2.4.

Lemma 6.1. *Let $m \in \mathbb{R}$, $0 \leq \delta, \rho \leq 1$, $\sigma \in BS_{\rho, \delta}^m$, and $N > n$. Then,*

(a) *If $0 < R \leq 1$ and $\text{supp}(\sigma) \subset \{(x, \xi, \eta) : |\xi| + |\eta| \leq R\}$, then*

$$\|T_\sigma(f, g)\|_{L^\infty} \lesssim R^{2n} \|\sigma\|_{0, 2N} \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in L^\infty;$$

(b) *If $R \geq 1$ and $\text{supp}(\sigma) \subset \{R \leq |\xi| + |\eta| \leq 4R\}$, then*

$$\|T_\sigma(f, g)\|_{L^\infty} \lesssim R^{(1-\rho)n+m} \|\sigma\|_{0, 2N} \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in L^\infty.$$

Lemma 6.2. *Let $Q \subset \mathbb{R}^n$ be a cube with diameter d and $\sigma \in BS_{\rho, \delta}^m$ with $m = n(\rho - 1)$, $0 \leq \delta, \rho \leq 1$, such that $\text{supp}(\sigma) \subset \{(x, \xi, \eta) : |\xi| + |\eta| \leq d^{-1}\}$. Then, for every $N > n$,*

$$\frac{1}{|Q|} \int |T_\sigma(f, g)(x) - T_\sigma(f, g)_Q| \, dx \lesssim \|\sigma\|_{1, 2N} \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in S,$$

with constants depending only on n, N, ρ , and δ . Here, $T_\sigma(f, g)_Q$ is the average of $T_\sigma(f, g)$ over Q .

Lemma 6.3. *Let $d > 0$ and $\sigma \in BS_{\rho, \delta}^m$, $m = n(\rho - 1)$, $0 \leq \delta, \rho \leq 1$, such that $\text{supp}(\sigma) \subset \{(x, \xi, \eta) : |\xi| + |\eta| \geq d^{-1}\}$. Let $\phi \in S$, $\phi \geq 0$, and $\text{supp}(\hat{\phi}) \subset \{z \in \mathbb{R}^n : |z| \leq \frac{1}{8}d^{-\rho}\}$. For $f, g \in S$, define*

$$R(f, g)(x) := \phi^2(x)T_\sigma(f, g)(x) - T_\sigma(\phi f, \phi g)(x).$$

Then, for every $N > n$, we have

$$\|R(f, g)\|_{L^\infty} \lesssim \|\sigma\|_{0, 2N+1} \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in S,$$

with constants only depending on n, N, ρ , and δ .

Remark 6.4. The proofs of the above lemmas show that $BS_{\rho, \delta}^m$ can be replaced by $BS_{\rho, \delta, 0, 2N}^m$, $BS_{\rho, \delta, 1, 2N}^m$, and $BS_{\rho, \delta, 0, 2N+1}^m$, respectively (see the definition of these spaces in Section 2.1).

Proof of Theorem 2.4. Given $\sigma \in BS_{\rho, 0}^m$, with $m = n(\rho - 1)$, we need to prove

$$(6.1) \quad \frac{1}{|Q|} \int_Q |T_\sigma(f, g)(x) - T_\sigma(f, g)_Q| \, dx \lesssim \|f\|_{L^\infty} \|g\|_{L^\infty},$$

for all cubes $Q \subset \mathbb{R}^n$ and $f, g \in S$.

Let Q be a cube with diameter d , and assume first that $d \leq 1$. We write

$$\sigma(x, \xi, \eta) = \sigma(x, \xi, \eta)(1 - \theta(\xi, \eta)) + \sigma(x, \xi, \eta)\theta(\xi, \eta) =: \sigma_1 + \sigma_2,$$

where $\theta : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a smooth, non-negative function, $\theta(\xi, \eta) = \tilde{\theta}(d\xi, d\eta)$ with $\text{supp}(\tilde{\theta}) \subset \{(\xi, \eta) \in \mathbb{R}^n \times \mathbb{R}^n : |\xi| + |\eta| \geq 1\}$, and $\tilde{\theta} \equiv 1$ in $\{(\xi, \eta) \in \mathbb{R}^n \times \mathbb{R}^n : |\xi| + |\eta| \geq 2\}$. Since $d \leq 1$, then $\sigma_1, \sigma_2 \in BS_{\rho,0}^m$ satisfy

$$(6.2) \quad \|\sigma_j\|_{K,M} \lesssim \|\sigma\|_{K,M}, \quad K, M \in \mathbb{N}_0, \quad j = 1, 2,$$

with constants independent of d and σ .

Let ϕ be as in Lemma 6.3, that is, $\phi \in S$, $\phi \geq 0$, and $\text{supp}(\hat{\phi}) \subset \{z \in \mathbb{R}^n : |z| \leq d^{-\rho}/8\}$. In addition, we assume $\phi \equiv 1$ on Q , and in accordance with the uncertainty principle, we choose ϕ such that $\|\phi\|_{L^2} \leq d^{n\rho/2}$. For $x \in Q$, we have

$$\begin{aligned} T_\sigma(f, g)(x) &= T_{\sigma_1}(f, g)(x) + T_{\sigma_2}(f, g)(x) \\ &= T_{\sigma_1}(f, g)(x) + \phi^2(x)T_{\sigma_2}(f, g)(x) \\ &= T_{\sigma_1}(f, g)(x) + T_{\sigma_2}(\phi f, \phi g)(x) + R(f, g)(x), \end{aligned}$$

where $R(f, g)(x) = \phi^2(x)T_{\sigma_2}(f, g)(x) - T_{\sigma_2}(\phi f, \phi g)(x)$.

In order to get (6.1), it is enough to prove the inequality

$$(6.3) \quad \|T_{\sigma_2}(\phi f, \phi g)\|_{L^1(Q)} \lesssim \|\sigma_2\|_{K,M} d^n \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in S,$$

for some $K, M \in \mathbb{N}_0$. Indeed, using (6.3) and Lemmas 6.2 and 6.3, for $N > n$, we write

$$\begin{aligned} &\frac{1}{|Q|} \int_Q |T_\sigma(f, g)(x) - T_\sigma(f, g)_Q| \, dx \\ &\leq \frac{1}{|Q|} \int_Q |T_{\sigma_1}(f, g)(x) - T_{\sigma_1}(f, g)_Q| \, dx \\ &\quad + \frac{2}{|Q|} \|T_{\sigma_2}(\phi f, \phi g)\|_{L^1(Q)} + 2\|R(f, g)\|_{L^\infty} \\ &\lesssim (\|\sigma_1\|_{1,2N} + \|\sigma_2\|_{K,M} + \|\sigma_2\|_{0,2N+1}) \|f\|_{L^\infty} \|g\|_{L^\infty}, \end{aligned}$$

and therefore (6.1) holds when the diameter of Q is less than or equal to 1.

In turn, (6.3) will follow from

$$(6.4) \quad \|T_{\sigma_2}(\phi f, \phi g)\|_{L^2} \lesssim \|\sigma_2\|_{K,M} d^{n/2} \|f\|_{L^\infty} \|g\|_{L^\infty}, \quad f, g \in S,$$

since

$$\begin{aligned} \frac{1}{|Q|} \|T_{\sigma_2}(\phi f, \phi g)\|_{L^1(Q)} &\leq \frac{1}{|Q|^{1/2}} \|T_{\sigma_2}(\phi f, \phi g)\|_{L^2(Q)} \\ &\leq \frac{1}{|Q|^{1/2}} \|T_{\sigma_2}(\phi f, \phi g)\|_{L^2}. \end{aligned}$$

Moreover, because ϕ satisfies $\|\phi\|_{L^2} \lesssim d^{\rho n/2}$, (6.4) can be reduced to proving that

$$\|T_{\sigma_2}\|_{L^2 \times L^2 \rightarrow L^2} \lesssim \|\sigma_2\|_{K,M} d^{n/2-\rho n}.$$

By Theorem 2.3, the support of σ_2 , and the fact that $\sigma_2 \in BS_{\rho,0}^m$ with $m = n(\rho - 1)$ and $0 < \rho < \frac{1}{2}$, we obtain

$$\begin{aligned} \|T_{\sigma_2}\|_{L^2 \times L^2 \rightarrow L^2} &\lesssim \sup_{\substack{|\beta| \leq [n/2]+1 \\ |\alpha| \leq 2(2n+1)}} \sup_{\gamma, \xi \in \mathbb{R}^n} \|\partial_{\xi}^{\alpha} \partial_{\gamma}^{\beta} \sigma_2(\gamma, \xi - \cdot, \cdot)\|_{L^2} \\ &\lesssim \|\sigma_2\|_{K,M} \sup_{\xi \in \mathbb{R}^n} \|\chi_{\{|\xi-\eta|+|\eta| \geq d^{-1}\}}(\xi, \eta) (1 + |\xi - \eta| + |\eta|)^m\|_{L^2(d\eta)} \\ &\lesssim \|\sigma_2\|_{K,M} \left[\left(\int_{|\eta| \geq d^{-1}} |\eta|^{2m} d\eta \right)^{1/2} + \left(\int_{|\eta| \leq d^{-1}} d^{-2m} d\eta \right)^{1/2} \right] \\ &\lesssim \|\sigma_2\|_{K,M} d^{-m-n/2} = \|\sigma_2\|_{K,M} d^{n/2-\rho n}, \end{aligned}$$

where we have taken $K = [n/2] + 1$ and $M = 2(2n + 1)$.

The case $d > 1$ follows using the decomposition of σ with $\theta = \tilde{\theta}$ and then proceeding analogously, but applying to the term corresponding to T_{σ_1} Lemma 6.1 instead of Lemma 6.2. \square

7. PROOF OF THEOREM 2.5

For $s > 0$, we recall the bilinear fractional integral operator of order $s > 0$, introduced in Kenig-Stein [21], defined by

$$I_s(f, g)(x) := \int_{\mathbb{R}^{2n}} \frac{f(y)g(z)}{(|x - y| + |x - z|)^{2n-s}} dy dz, \quad x \in \mathbb{R}^n.$$

It easily follows that

$$|I_s(f, g)(x)| \leq I_{s_1}(|f|)(x) I_{s_2}(|g|)(x), \quad x \in \mathbb{R}^n, \quad s_1 + s_2 = s,$$

where

$$I_{\tau}(h)(x) = \int_{\mathbb{R}^n} \frac{h(y)}{|x - y|^{n-\tau}} dy, \quad 0 < \tau < n,$$

is the linear fractional integral. The boundedness properties of I_{τ} , $0 < \tau < n$, and Hölder's inequality imply that I_s is bounded from $L^{p_1} \times L^{p_2}$ into L^p with $1/p = 1/p_1 + 1/p_2 - s/n$, $0 < s < 2n$, $1 < p_1, p_2 < \infty$, $q > 0$.

We now observe that if $\sigma \in BS_{\rho,\delta}^m$, $m \leq 2n(\rho - 1) - \rho s$, $0 < s < 2n$, then part (v) of Theorem E implies that

$$(7.1) \quad |T_{\sigma}(f, g)(x)| \lesssim |I_s(f, g)(x)|.$$

Therefore, Theorem 2.5 follows from this inequality and the boundedness properties of I_s . The case $\rho = 1$ of Theorem 2.5 was treated in [8].

8. PROOFS OF LEMMAS FROM SECTION 6

Proof of Lemma 6.1. We have

$$(8.1) \quad T_\sigma(f, g)(x) = \int_{\mathbb{R}^{2n}} \mathcal{K}(x, x - y, x - z) f(y) g(z) \, dy \, dz,$$

where

$$\mathcal{K}(x, y, z) = \int_{\mathbb{R}^{2n}} e^{i\xi \cdot y} e^{i\eta \cdot z} \sigma(x, \xi, \eta) \, d\xi \, d\eta = \mathcal{F}_{2n}^{-1}(\sigma(x, \cdot, \cdot))(y, z),$$

and \mathcal{F}_{2n}^{-1} denotes the inverse Fourier transform in \mathbb{R}^{2n} . Then, it is enough to show that, for $N > n$, $N \in \mathbb{N}_0$,

$$(8.2) \quad \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^{2n}} |\mathcal{K}(x, y, z)| \, dy \, dz \lesssim R^{2n} \|\sigma\|_{0,2N}$$

and

$$(8.3) \quad \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^{2n}} |\mathcal{K}(x, y, z)| \, dy \, dz \lesssim R^{(1-\rho)n+m} \|\sigma\|_{0,2N},$$

for part (a) and part (b), respectively. (Note that this allows us to extend T_σ to a bounded operator form $L^\infty \times L^\infty$ into L^∞ by using the representation (8.1) to define $T_\sigma(f, g)$ for $f, g \in L^\infty$.)

Since σ is a smooth function with compact support in ξ and η , we have

$$(8.4) \quad \begin{aligned} & (1 + |(y, z)|^2)^N \mathcal{K}(x, y, z) \\ &= \int_{\mathbb{R}^{2n}} \sigma(x, \xi, \eta) (1 - \Delta_\xi - \Delta_\eta)^N (e^{i\xi \cdot y} e^{i\eta \cdot z}) \, d\xi \, d\eta \\ &= \int_{\mathbb{R}^{2n}} (1 - \Delta_\xi - \Delta_\eta)^N (\sigma(x, \xi, \eta)) e^{i\xi \cdot y} e^{i\eta \cdot z} \, d\xi \, d\eta \\ &= \mathcal{F}_{2n}^{-1}((1 - \Delta_\xi - \Delta_\eta)^N (\sigma(x, \cdot, \cdot)))(y, z), \end{aligned}$$

and similarly,

$$(8.5) \quad |(y, z)|^{2N} \mathcal{K}(x, y, z) = \mathcal{F}_{2n}^{-1}((-\Delta_\xi - \Delta_\eta)^N (\sigma(x, \cdot, \cdot)))(y, z).$$

For part (a), we use (8.4) and the fact that $R \leq 1$ to get

$$|\mathcal{K}(x, y, z)| \lesssim \frac{R^{2n} \|\sigma\|_{0,2N}}{(1 + |(y, z)|^2)^N},$$

and then (8.2) follows since $N > n$.

For part (b), we write

$$\int_{\mathbb{R}^{2n}} |\mathcal{K}(x, y, z)| \, dy \, dz = \int_{|y|+|z|\leq R^{-\rho}} |\mathcal{K}(x, y, z)| \, dy \, dz + \int_{|y|+|z|\geq R^{-\rho}} |\mathcal{K}(x, y, z)| \, dy \, dz.$$

Let us now estimate the first integral. By the Cauchy-Schwarz inequality, Plancherel’s identity, and the fact that $R \geq 1$, we have

$$\begin{aligned} & \left(\int_{|y|+|z|\leq R^{-\rho}} |\mathcal{K}(x, y, z)| \, dy \, dz \right)^2 \\ & \leq R^{-2\rho n} \int_{|y|+|z|\leq R^{-\rho}} |\mathcal{K}(x, y, z)|^2 \, dy \, dz \\ & \leq R^{-2\rho n} \int_{|\xi|+|\eta|\sim R} |\sigma(x, \xi, \eta)|^2 \, d\xi \, d\eta \\ & \leq \|\sigma\|_{0,0}^2 R^{-2\rho n} \int_{|\xi|+|\eta|\sim R} (1 + |\xi| + |\eta|)^{2m} \, d\xi \, d\eta \\ & \leq \|\sigma\|_{0,0}^2 R^{-2\rho n} R^{2m+2n} = \|\sigma\|_{0,0}^2 R^{2((1-\rho)n+m)}. \end{aligned}$$

Next, we estimate the second integral. Multiplying and dividing by $|(y, z)|^{2N}$, and using the Cauchy-Schwarz inequality, (8.5), Plancherel’s identity, and the facts that $N > n$ and $R \geq 1$, we have that

$$\begin{aligned} & \left(\int_{|y|+|z|\geq R^{-\rho}} |\mathcal{K}(x, y, z)| \, dy \, dz \right)^2 \\ & \leq \left(\int_{|y|+|z|\geq R^{-\rho}} \frac{1}{|(y, z)|^{4N}} \, dy \, dz \right) \\ & \quad \times \left(\int_{|y|+|z|\geq R^{-\rho}} | |(y, z)|^{2N} \mathcal{K}(x, y, z) |^2 \, dy \, dz \right) \\ & \leq R^{\rho(4N-2n)} \int_{|\xi|+|\eta|\sim R} |(-\Delta_\xi - \Delta_\eta)^N \sigma(x, \xi, \eta)|^2 \, d\xi \, d\eta \\ & \leq \|\sigma\|_{0,2N}^2 R^{\rho(4N-2n)} \int_{|\xi|+|\eta|\sim R} (1 + |\xi| + |\eta|)^{2(m-\rho 2N)} \, d\xi \, d\eta \\ & \leq \|\sigma\|_{0,2N}^2 R^{\rho(4N-2n)} R^{2(m-\rho 2N+n)} = \|\sigma\|_{0,2N}^2 R^{2((1-\rho)n+m)}. \end{aligned}$$

The last two computations give (8.3). □

Proof of Lemma 6.2. Let Q, d, N, m , and σ be as in the hypothesis. By definition,

$$T_\sigma(f, g)(x) = \int_{\mathbb{R}^{2n}} \sigma(x, \xi, \eta) \hat{f}(\xi) \hat{g}(\eta) e^{ix \cdot (\xi + \eta)} \, d\xi \, d\eta, \quad f, g \in \mathcal{S}.$$

Hence, for a fixed $j = 1, \dots, n$, the bilinear symbol $\tau = \tau(x, \xi, \eta)$ of the bilinear operator $\partial T_\sigma(f, g) / \partial x_j$ is given by

$$\tau(x, \xi, \eta) = i(\xi_j + \eta_j)\sigma(x, \xi, \eta) + \frac{\partial \sigma}{\partial x_j}(\xi, \xi, \eta).$$

Then, the symbol τ is also supported in $\{(x, \xi, \eta) : |\xi| + |\eta| \leq d^{-1}\}$, and we have $\tau \in BS_{\rho, \delta}^{m+\delta}$. Elementary computations show that, for $K, M \in \mathbb{N}_0$,

$$(8.6) \quad \|\tau\|_{K, M} \leq \max(1, d^{-1}) \|\sigma\|_{K+1, M},$$

where $\|\tau\|_{K, M}$ corresponds to a norm of τ as an element of $BS_{\rho, \delta}^{m+\delta}$, while $\|\sigma\|_{K+1, M}$ corresponds to a norm of σ as an element of $BS_{\rho, \delta}^m$. Then,

$$\begin{aligned} & \int_Q |T_\sigma(f, g)(x) - T_\sigma(f, g)_Q| dx \\ &= \frac{1}{|Q|} \int_Q \left| \int_Q (T_\sigma(f, g)(x) - T_\sigma(f, g)(y)) dy \right| dx \\ &\leq d|Q| \|\nabla T_\sigma(f, g)\|_{L^\infty} \lesssim d|Q| \|T_\tau(f, g)\|_{L^\infty} \\ &\leq d|Q| \min(1, d^{-2n}) \|\tau\|_{0, 2N} \|f\|_{L^\infty} \|g\|_{L^\infty} \\ &\lesssim |Q| \|\sigma\|_{1, 2N} \|f\|_{L^\infty} \|g\|_{L^\infty} \end{aligned} \quad (\text{by (8.6)}),$$

where we have used Lemma 6.1. The result follows. □

Proof of Lemma 6.3. Let d, N, m, ϕ , and σ be as in the hypothesis. We notice that the bilinear symbol $\theta(x, \xi, \eta)$ of R is given by

$$\theta(x, \xi, \eta) = \int_{\mathbb{R}^{2n}} e^{ix \cdot (y+z)} (\sigma(x, \xi, \eta) - \sigma(x, \xi + y, \eta + z)) \hat{\phi}(y) \hat{\phi}(z) dy dz.$$

We first assume that $d \leq 1$, and note that

$$\text{supp}(\theta) \subset \{(x, \xi, \eta) : |\xi| + |\eta| \geq \frac{1}{2}d^{-1}\}.$$

Consider a partition of unity of \mathbb{R}^{2n} given by $\{\psi_k\}_{k \in \mathbb{N}_0}$,

$$\sum_{k \geq 0} \psi_k(\xi, \eta) = 1, \quad \xi, \eta \in \mathbb{R}^n,$$

where $\psi_0 \in S(\mathbb{R}^{2n})$ is supported in the set

$$\{(\xi, \eta) : |\xi| + |\eta| \leq 2d^{-1}\},$$

and $\psi_k(\xi, \eta) = \psi(d2^{-k}\xi, d2^{-k}\eta)$ with $\psi \in S(\mathbb{R}^{2n})$ and $\text{supp}(\psi) \subset \{(\xi, \eta) : \frac{1}{2} \leq |\xi| + |\eta| \leq 2\}$ for $k \geq 1$. Then, $\text{supp}(\psi_k) \subset \{(\xi, \eta) : |\xi| + |\eta| \sim 2^k d^{-1}\}$ for $k \geq 1$, and

$$\theta(x, \xi, \eta) = \sum_{k \geq 0} \theta_k(x, \xi, \eta),$$

where $\theta_k(x, \xi, \eta) := \theta(x, \xi, \eta)\psi_k(\xi, \eta)$. We will show that, for all integers $M, k \geq 0$,

$$(8.7) \quad \|\theta_k\|_{0,M} \lesssim 2^{-\rho k} \|\sigma\|_{0,M+1},$$

with constants depending only on M, n, ρ , and δ .

Define R_k as the bilinear pseudo-differential operator with kernel θ_k . The lemma will follow from (8.7). Indeed,

$$\begin{aligned} \|R(f, g)\|_{L^\infty} &\leq \sum_{k \geq 0} \|R_k(f, g)\|_{L^\infty} \\ &\lesssim \sum_{k \geq 0} \|\theta_k\|_{0,2N} \|f\|_{L^\infty} \|g\|_{L^\infty} \quad (\text{by Lemma 6.1}) \\ &\lesssim \sum_{k \geq 0} 2^{-\rho k} \|\sigma\|_{0,2N+1} \|f\|_{L^\infty} \|g\|_{L^\infty} \quad (\text{by (8.7)}) \\ &\lesssim \|\sigma\|_{0,2N+1} \|f\|_{L^\infty} \|g\|_{L^\infty}. \end{aligned}$$

To prove (8.7), consider multi-indices β and γ such that $|\beta|, |\gamma| \leq M$. Since

$$\begin{aligned} &\theta_k(x, \xi, \eta) \\ &= \int_{\mathbb{R}^{2n}} e^{ix \cdot (\gamma+z)} \psi_k(\xi, \eta) (\sigma(x, \xi, \eta) - \sigma(x, \xi + \gamma, \eta + z)) \hat{\phi}(\gamma) \hat{\phi}(z) \, dy \, dz, \end{aligned}$$

we have

$$\begin{aligned} &(\partial_\xi^\beta \partial_\eta^\gamma \theta_k)(x, \xi, \eta) \\ &= \sum_{\lambda \leq \gamma, \omega \leq \beta} C_{\beta, \gamma, \omega, \lambda} (\partial_\xi^{\beta-\omega} \partial_\eta^{\gamma-\lambda} \psi)(d2^{-k}\xi, d2^{-k}\eta) (2^{-k}d)^{|\gamma-\lambda|+|\beta-\omega|} \\ &\times \int_{\mathbb{R}^{2n}} \hat{\phi}(\gamma) \hat{\phi}(z) e^{ix \cdot (\gamma+z)} ((\partial_\xi^\omega \partial_\eta^\lambda \sigma)(x, \xi, \eta) - (\partial_\xi^\omega \partial_\eta^\lambda \sigma)(x, \xi + \gamma, \eta + z)) \, dy \, dz. \end{aligned}$$

The mean value theorem gives

$$(\partial_\xi^\omega \partial_\eta^\lambda \sigma)(x, \xi, \eta) - (\partial_\xi^\omega \partial_\eta^\lambda \sigma)(x, \xi + \gamma, \eta + z) = (\nabla_\xi \partial_\xi^\omega \nabla_\eta \partial_\eta^\lambda \sigma)(x, \tilde{\xi}, \tilde{\eta}) \cdot (\gamma, z),$$

where $(\tilde{\xi}, \tilde{\eta}) = (\xi, \eta) + s(\gamma, z)$ for some $s \in (0, 1)$. Since $\sigma \in BS_{\rho, \delta}^m$, for

$$(\xi, \eta) \in \text{supp}(\psi_k) \cap \text{supp}(\theta) \text{ and } \gamma, z \in \text{supp}(\hat{\phi}),$$

we then have

$$\begin{aligned} & |(\partial_{\xi}^{\omega} \partial_{\eta}^{\lambda} \sigma)(x, \xi, \eta) - (\partial_{\xi}^{\omega} \partial_{\eta}^{\lambda} \sigma)(x, \xi + \gamma, \eta + z)| \\ & \leq \|\sigma\|_{0, M+1} (1 + |\tilde{\xi}| + |\tilde{\eta}|)^{m-\rho(|\omega|+|\lambda|+1)} |\gamma, z| \\ & \leq \|\sigma\|_{0, M+1} (1 + |\xi| + |\eta|)^{m-\rho(|\omega|+|\lambda|+1)} |\gamma, z|, \end{aligned}$$

where we have used that $|\tilde{\xi}| + |\tilde{\eta}| \approx |\xi| + |\eta|$, since $|\xi| + |\eta| \approx 2^k d^{-1}$ and $|\gamma| + |z| \leq d^{-\rho}/4 \leq d^{-1}/4$. Putting this all together, and using again that $2^k d^{-1} \geq 1$, $d \leq 1$, and $1 + |\xi| + |\eta| \approx |\xi| + |\eta| \approx 2^k d^{-1}$, we have

$$\begin{aligned} |\partial_{\xi}^{\beta} \partial_{\eta}^{\gamma} \theta_k(x, \xi, \eta)| & \lesssim \|\sigma\|_{0, M+1} (1 + |\xi| + |\eta|)^{m-\rho(|\gamma|+|\beta|)} (1 + 2^k d^{-1})^{-\rho} \\ & \quad \times \sum_{\lambda \leq \gamma, \omega \leq \beta} (2^{-k} d)^{(1-\rho)(|\gamma-\lambda|+|\beta-\omega|)} \\ & \lesssim \|\sigma\|_{0, M+1} (1 + |\xi| + |\eta|)^{m-\rho(|\gamma|+|\beta|)} 2^{-\rho k}, \end{aligned}$$

which gives (8.7). □

If $d > 1$, we then split θ as

$$\theta = \theta_1 + \theta_2,$$

where $\text{supp}(\theta_1) \subset \{(\xi, \eta) : |\xi| + |\eta| \leq 2\}$ (note that, in the case $d > 1$, we also have $|\gamma|, |z| \leq d^{-\rho}/8 \leq \frac{1}{8}$) and $\text{supp}(\theta_2) \subset \{(\xi, \eta) : |\xi| + |\eta| \geq 1\}$. A similar reasoning as above shows that $\|\theta_1\|_{0, M} \lesssim \|\sigma\|_{0, M+1}$. We then apply Lemma 6.1 to the bilinear pseudo-differential operator with symbol θ_1 , and reduce the analysis of θ_2 to the case $d = 1$.

9. WEIGHTED RESULTS

Given a weight w defined on \mathbb{R}^n and $p > 0$, the notation L_w^p will be used to refer to the weighted Lebesgue space of all functions $f : \mathbb{R}^n \rightarrow \mathbb{C}$ such that

$$\|f\|_{L_w^p} := \int_{\mathbb{R}^n} |f(x)|^p w(x) dx < \infty;$$

when $w \equiv 1$, we will continue to simply write L^p and $\|f\|_{L^p}$, respectively.

If w_1, w_2 are weights defined on \mathbb{R}^n , $1 \leq p_1, p_2 < \infty$, $q > 0$, and $w := w_1^{q/p_1} w_2^{q/p_2}$, we then say that (w_1, w_2) satisfies the $A_{(p_1, p_2), q}$ condition (or that (w_1, w_2) belongs to the bilinear Muckenhoupt class $A_{(p_1, p_2), q}$) if

$$\begin{aligned} & [(w_1, w_2)]_{A_{(p_1, p_2), q}} \\ & := \sup_B \left(\frac{1}{|B|} \int_B w(x) dx \right) \prod_{j=1}^2 \left(\frac{1}{|B|} \int_B w_j(x)^{1-p'_j} dx \right)^{q/p'_j} < \infty, \end{aligned}$$

where the supremum is taken over all Euclidean balls $B \subset \mathbb{R}^n$; when $p_j = 1$, $((1/|B|) \int_B w_j(x)^{1-p'_j} dx)^{1/p'_j}$ is understood as $(\inf_B w_j)^{-1}$.

The classes $A_{(p_1,p_2),q}$ are inspired by the classes of weights $A_{p,q}$, $1 \leq p, q < \infty$, defined by Muckenhoupt and Wheeden in [26] to study weighted norm inequalities for the fractional integral: a weight u defined on \mathbb{R}^n is in the class $A_{p,q}$ if

$$\sup_B \left(\frac{1}{|B|} \int_B u^{q/p} dx \right) \left(\frac{1}{|B|} \int_B u^{(1-p')} dx \right)^{q/p'} < \infty.$$

The classes $A_{(p_1,p_2),p^1}$ for $1/p = 1/p_1 + 1/p_2$ were introduced by Lerner *et al* in [22] to study characterizations of weights for boundedness properties of certain bilinear maximal functions and bilinear Calderón-Zygmund operators in weighted Lebesgue spaces. Likewise, as shown by Moen [25], the classes $A_{(p_1,p_2),q}$ characterize the weights rendering analogous bounds for bilinear fractional integral operators.

Theorem 4.1 and [22, Corollary 3.9] imply the following result.

Corollary 9.1. *Let $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$, $0 < \rho$, $m_{cz} = 2n(\rho - 1)$, $1 \leq p_1, p_2 < \infty$, and p given by $1/p = 1/p_1 + 1/p_2$. Suppose $\sigma \in BS_{\rho,\delta}^m$, $m < m_{cz}$, (w_1, w_2) satisfies the $A_{(p_1,p_2),p}$ condition, and that $w = w_1^{\sigma/p_1} w_2^{\sigma/p_2}$.*

(a) *If $1 < p_1, p_2 < \infty$, then there exists $K, N \in \mathbb{N}_0$ such that*

$$\|T_\sigma(f, g)\|_{L_w^p} \lesssim \|\sigma\|_{K,N} \|f\|_{L_{w_1}^{p_1}} \|g\|_{L_{w_2}^{p_2}}.$$

(b) *If $1 \leq p_1, p_2 < \infty$ and $p_1 = 1$ or $p_2 = 1$, then there exists $K, N \in \mathbb{N}_0$ such that*

$$\|T(f, g)\|_{L_w^{p,\infty}} \lesssim \|\sigma\|_{K,N} \|f\|_{L_{w_1}^{p_1}} \|g\|_{L_{w_2}^{p_2}}.$$

Inequality (7.1) and [25, Theorem 3.5] yield the following result.

Corollary 9.2 (Weighted version of Theorem 2.5). *Let $0 \leq \delta \leq 1$, $0 < \rho \leq 1$, $s \in (0, 2n)$, and $m_s := 2n(\rho - 1) - \rho s$. If $\sigma \in BS_{\rho,\delta}^{m_s}$, $m \leq m_s$, and $1/q = 1/p_1 + 1/p_2 - s/n$, $1 < p_1, p_2 < \infty$, then there exist nonnegative integers K and N such that*

$$\|T_\sigma(f, g)\|_{L_w^q} \lesssim \|\sigma\|_{K,N} \|f\|_{L_{w_1}^{p_1}} \|g\|_{L_{w_2}^{p_2}},$$

for $w := w_1^{q/p_1} w_2^{q/p_2}$ and pairs of weights (w_1, w_2) satisfying the $A_{(p_1,p_2),q}$ condition.

Acknowledgements. The first author is partially supported by a grant from the Simons Foundation (No. 246024). Partial NSF support under the following grants is also acknowledged: third author DMS 0901587, fourth author DMS 1101327, fifth author DMS 0800492 and DMS 1069015.

¹These classes were denoted by $A_{\vec{p}}$ in [22], with $\vec{p} = (p_1, p_2)$ determining $1/p = 1/p_1 + 1/p_2$.

REFERENCES

- [1] J. ÁLVAREZ and J. HOUNIE, *Estimates for the kernel and continuity properties of pseudo-differential operators*, Ark. Mat. **28** (1990), no. 1, 1–22.
<http://dx.doi.org/10.1007/BF02387364>. MR1049640 (91d:35255)
- [2] Á. BÉNYI, D. MALDONADO, V. NAIBO, and R. H. TORRES, *On the Hörmander classes of bilinear pseudodifferential operators*, Integral Equations Operator Theory **67** (2010), no. 3, 341–364. <http://dx.doi.org/10.1007/s00020-010-1782-y>. MR2660466 (2011g:47111)
- [3] Á. BÉNYI and R. H. TORRES, *Symbolic calculus and the transposes of bilinear pseudodifferential operators*, Comm. Partial Differential Equations **28** (2003), no. 5–6, 1161–1181.
<http://dx.doi.org/10.1081/PDE-120021190>. MR1986065 (2004d:35260)
- [4] ———, *Almost orthogonality and a class of bounded bilinear pseudodifferential operators*, Math. Res. Lett. **11** (2004), no. 1, 1–11.
<http://dx.doi.org/10.4310/MRL.2004.v11.n1.a1>. MR2046194 (2005a:35293)
- [5] J. BERGH and J. LÖFSTRÖM, *Interpolation Spaces: An Introduction*, Die Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 223, Springer-Verlag, Berlin, 1976. MR0482275 (58 #2349)
- [6] F. BERNICOT and P. GERMAIN, *Bilinear oscillatory integrals and boundedness for new bilinear multipliers*, Adv. Math. **225** (2010), no. 4, 1739–1785.
<http://dx.doi.org/10.1016/j.aim.2010.03.032>. MR2680189 (2011k:42025)
- [7] ———, *Bilinear dispersive estimates via space-time resonances, I: The one-dimensional case*, Anal. PDE **6** (2013), no. 3, 687–722. <http://dx.doi.org/10.2140/apde.2013.6.687>. MR3080192
- [8] F. BERNICOT, D. MALDONADO, K. MOEN, and V. NAIBO, *Bilinear Sobolev-Poincaré inequalities and Leibniz-type rules*, J. Geom. Anal. **24** (April 2014), no. 2, 1144–1180.
<http://dx.doi.org/10.1007/s12220-012-9367-4>.
- [9] F. BERNICOT and S. SHRIVASTAVA, *Boundedness of smooth bilinear square functions and applications to some bilinear pseudo-differential operators*, Indiana Univ. Math. J. **60** (2011), no. 1, 233–268. <http://dx.doi.org/10.1512/iumj.2011.60.4527>. MR2952417
- [10] A. P. CALDERÓN and R. VAILLANCOURT, *A class of bounded pseudo-differential operators*, Proc. Nat. Acad. Sci. U.S.A. **69** (1972), no. 5, 1185–1187.
<http://dx.doi.org/10.1073/pnas.69.5.1185>. MR0298480 (45 #7532)
- [11] F. M. CHRIST and J. L. JOURNÉ, *Polynomial growth estimates for multilinear singular integral operators*, Acta Math. **159** (1987), no. 1–2, 51–80.
<http://dx.doi.org/10.1007/BF02392554>. MR906525 (89a:42024)
- [12] F. M. CHRIST and M. I. WEINSTEIN, *Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation*, J. Funct. Anal. **100** (1991), no. 1, 87–109.
[http://dx.doi.org/10.1016/0022-1236\(91\)90103-C](http://dx.doi.org/10.1016/0022-1236(91)90103-C). MR1124294 (92h:35203)
- [13] R. R. COIFMAN and Y. MEYER, *Au delà des opérateurs pseudo-différentiels*, Astérisque, vol. 57, Société Mathématique de France, Paris, 1978 (French). With an English summary. MR518170 (81b:47061)
- [14] Ch. FEFFERMAN, *L^p bounds for pseudo-differential operators*, Israel J. Math. **14** (1973), no. 4, 413–417. <http://dx.doi.org/10.1007/BF02764718>. MR0336453 (49 #1227)
- [15] L. GRAFAKOS and R. H. TORRES, *Multilinear Calderón-Zygmund theory*, Adv. Math. **165** (2002), no. 1, 124–164.
<http://dx.doi.org/10.1006/aima.2001.2028>. MR1880324 (2002j:42029)
- [16] A. GULISASHVILI and M. A. KON, *Exact smoothing properties of Schrödinger semigroups*, Amer. J. Math. **118** (1996), no. 6, 1215–1248. MR1420922 (97m:47053)
- [17] L. HÖRMANDER, *Pseudo-differential operators and hypoelliptic equations*, Singular Integrals (Proc. Sympos. Pure Math., Vol. X, Chicago, Ill., 1966), Amer. Math. Soc., Providence, R.I., 1967, pp. 138–183. MR0383152 (52 #4033)

- [18] T. KATO and G. PONCE, *Commutator estimates and the Euler and Navier-Stokes equations*, Comm. Pure Appl. Math. **41** (1988), no. 7, 891–907.
<http://dx.doi.org/10.1002/cpa.3160410704>. MR951744 (90f:35162)
- [19] C. E. KENIG, G. PONCE, and L. VEGA, *Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle*, Comm. Pure Appl. Math. **46** (1993), no. 4, 527–620. <http://dx.doi.org/10.1002/cpa.3160460405>. MR1211741 (94h:35229)
- [20] C. E. KENIG and W. STAUBACH, *Ψ -pseudodifferential operators and estimates for maximal oscillatory integrals*, Studia Math. **183** (2007), no. 3, 249–258.
<http://dx.doi.org/10.4064/sm183-3-3>. MR2357989 (2009d:35361)
- [21] C. E. KENIG and E. M. STEIN, *Multilinear estimates and fractional integration*, Math. Res. Lett. **6** (1999), no. 1, 1–15.
<http://dx.doi.org/10.4310/MRL.1999.v6.n1.a1>. MR1682725 (2000k:42023a)
- [22] A. K. LERNER, Sh. OMBROSI, C. PÉREZ, R. H. TORRES, and R. TRUJILLO-GONZÁLEZ, *New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory*, Adv. Math. **220** (2009), no. 4, 1222–1264.
<http://dx.doi.org/10.1016/j.aim.2008.10.014>. MR2483720 (2010f:42024)
- [23] D. MALDONADO and V. NAIBO, *Weighted norm inequalities for paraproducts and bilinear pseudodifferential operators with mild regularity*, J. Fourier Anal. Appl. **15** (2009), no. 2, 218–261.
<http://dx.doi.org/10.1007/s00041-008-9029-x>. MR2500923 (2010b:42027)
- [24] N. MICHALOWSKI, D. RULE, and W. STAUBACH, *Multilinear pseudodifferential operators beyond Calderón-Zygmund theory*, J. Math. Anal. Appl. **414** (2014), no. 1, 149–165.
<http://dx.doi.org/10.1016/j.jmaa.2013.12.062>. MR3165300
- [25] K. MOEN, *Weighted inequalities for multilinear fractional integral operators*, Collect. Math. **60** (2009), no. 2, 213–238.
<http://dx.doi.org/10.1007/BF03191210>. MR2514845 (2010d:42028)
- [26] B. MUCKENHOUTP and R. WHEEDEN, *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc. **192** (1974), 261–274.
<http://dx.doi.org/10.2307/1996833>. MR0340523 (49 #5275)
- [27] L. PÄIVÄRINTA and E. SOMERSALO, *A generalization of the Calderón-Vaillancourt theorem to L^p and h^p* , Math. Nachr. **138** (1988), 145–156.
<http://dx.doi.org/10.1002/mana.19881380111>. MR975206 (89m:46043)
- [28] S. SEMMES, *Nonlinear Fourier analysis*, Bull. Amer. Math. Soc. (N.S.) **20** (1989), no. 1, 1–18.
<http://dx.doi.org/10.1090/S0273-0979-1989-15681-4>. MR959211 (89i:42001)

ÁRPÁD BÉNYI:
 Department of Mathematics
 Western Washington University
 516 High St
 Bellingham, WA 98225, USA
 E-MAIL: arpad.benyi@wwwu.edu

FRÉDÉRIC BERNICOT:
 CNRS-Université de Nantes
 Laboratoire Jean Leray
 2, rue de la Houssinière
 44322 Nantes Cedex 3, France
 E-MAIL: frederic.bernicot@univ-nantes.fr

DIEGO MALDONADO & VIRGINIA NAIBO:
 Department of Mathematics
 Kansas State University
 138 Cardwell Hall
 Manhattan, KS 66506-2602, USA
 E-MAIL: dmaldona@math.ksu.edu,
 vnaibo@math.ksu.edu

RODOLFO H. TORRES:
 Department of Mathematics
 University of Kansas
 405 Snow Hall, 1460 Jayhawk Blvd
 Lawrence, KS 66045-7594, USA
 E-MAIL: torres@math.ku.edu

KEY WORDS AND PHRASES: Bilinear pseudodifferential operators, bilinear Hormander classes, symbolic calculus, Calderon-Zygmund theory.

1991 MATHEMATICS SUBJECT CLASSIFICATION: 35S05, 47G30 (42B15, 42B20).

Received: September 6, 2012.