



Apr 4th, 1:45 PM - 2:00 PM

## Boundary Bay ambient monitoring program

Carrie Hightower

*Metro Vancouver Regional District, Canada, [carrie.hightower@metrovancover.org](mailto:carrie.hightower@metrovancover.org)*

Follow this and additional works at: <https://cedar.wvu.edu/ssec>

 Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

---

Hightower, Carrie, "Boundary Bay ambient monitoring program" (2018). *Salish Sea Ecosystem Conference*. 18.  
<https://cedar.wvu.edu/ssec/2018ssec/allsessions/18>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).



# Boundary Bay Ambient Monitoring Program

## PROGRAM ASSESSMENT

Carrie Hightower

ENVIRONMENTAL TECHNICIAN

Salish Sea Ecosystem Conference, April 4, 2018



**metrovancover**  
SERVICES AND SOLUTIONS FOR A LIVABLE REGION




# Program Definition

- Why Boundary Bay?
- Partnership Approach
- Monitored Media
- Monitoring Frequency

# Program Assessment

- Summarize the results of the water, sediment and biota studies
- Critically review the program
- Provide recommendations for future monitoring






# Marine Water Monitoring Stations

# Key Findings – Marine Water Quality

- Majority of results met the Boundary Bay Water Quality Objectives and BC Water Quality Guidelines
- Dissolved oxygen has been low from most sites but this is consistent with marine coastal waters
- Some fecal coliform levels exceedances at 2 sites
- Limited organics analysis but all results have been lower than the analytical detection limits and guidelines

# Key Findings – Marine Water Quality

- Water is well mixed with no significant difference observed at the 2 depths, between-site or temporally
- When clustered, mean concentrations for some parameters were higher in samples from nearfield sites than farfield
- Results from wet season sampling were higher than dry period sampling for all sites



# Sediment and Biota Monitoring Stations



# Key Findings - Sediment Monitoring

- Most measured substances with met objectives or guidelines
- The concentrations of metals and organics are highly dependent upon sediment particle size
- Only one site with fine sediment (>94% silt plus clay)



# Key Findings – Biota

- Caged mussels had 82% survival rate
- Tissue concentrations for contaminants met guidelines
- At near field sites, contaminant concentrations were higher in caged mussel tissues and lipid content lower
- Macro benthos showed high degree of variation but likely due to natural variability



# Recommendations

- Dry season water sampling can be eliminated and sampling sites reduced
- Sediment component is the least valuable and should be re-evaluated in the context of the bay's geomorphology and physical oceanography
- Caged mussel study should be repeated at 5 year intervals and data interpreted on an ecotoxicological concentration response perspective and compared with the US Mussel Watch database
- Benthic community characterization should be re-interpreted against data from similar areas in the Salish Sea

# Next Steps

- Additional study of biota data
- Water column experimentation with sampling plan and instrument mooring installation
- Caged mussel study to be repeated
- Sediment and macro benthos component to be re-examined



Questions?