



Western Washington University
Western CEDAR

Salish Sea Ecosystem Conference

2022 Salish Sea Ecosystem Conference
(Online)

Apr 27th, 4:30 PM - 5:00 PM

Characterizing Contaminant Concentrations in Priority Chinook Salmon Stocks Consumed by Resident Killer Whales in the Northeastern Pacific Ocean

Stephanie Holbert
Simon Fraser University

Follow this and additional works at: <https://cedar.wwu.edu/ssec>




Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Holbert, Stephanie, "Characterizing Contaminant Concentrations in Priority Chinook Salmon Stocks Consumed by Resident Killer Whales in the Northeastern Pacific Ocean" (2022). *Salish Sea Ecosystem Conference*. 156.

<https://cedar.wwu.edu/ssec/2022ssec/allsessions/156>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

Contaminant Profiles in Priority Chinook Salmon Stocks Consumed by Resident Killer Whales in the Northeastern Pacific Ocean




Stephanie Holbert^{1,2*}, Katerina Colbourne², Frank Gobas¹, Tanya M. Brown^{2*}

¹ School of Resource and Environmental Management, Simon Fraser University, Burnaby, British Columbia

² Pacific Science Enterprise Centre, Fisheries and Oceans Canada, West Vancouver, British Columbia

Contact email: stephanie_holbert@sfu.ca



SIMON FRASER UNIVERSITY
ENGAGING THE WORLD



Introduction

Southern Resident Killer Whales (*Orcinus orca*) SRKW

- 74 individuals (3 pods: J, K, and L) inhabiting transboundary waters of the Salish Sea down to southern California, US¹
- Listed as *Endangered* in Canada (2003) & United States (2005)
- Top three current threats²
 - Availability or quality of priority prey, Chinook salmon
 - Environmental contaminants
 - Physical and acoustical disturbance
- Northern Resident Killer Whale (NRKW) sympatric population with similar diet but annual population growth³
- Diet studies revealed most consumed Chinook stocks for SRKW and NRKW come from the following river systems^{4,5,6}:
 - Fraser River, Columbia River, Puget Sound, Skeena, and Vancouver Island

Chinook salmon (*Oncorhynchus tshawytscha*)

- Consumed by SRKW year-round, making up 80% of SRKW diet and close to 100% during the spring and summer^{5,6}
- 94% (15/16) Chinook populations in Fraser River at risk or endangered⁷
- Little information exists on the concentrations in priority contaminants of concern (COCs) in Canadian Chinook salmon stocks


Contaminants of Concern (COCs)

- PCBs, PBDEs, OCPs, dioxin/furans, HBCDD, alkylphenols, and chlorinated paraffins have been identified as contaminants of concern to SRKW and their primary prey, Chinook salmon⁸
- SRKW & NRKW exposed to environmental contaminants mainly through their prey
- Recent analyses show PCB & PBDE blubber concentrations in SRKW exceed marine mammal health thresholds^{9,10,11}
- Measuring contaminants in primary prey, Chinook salmon, allows for a larger sample size to infer contaminant exposure to R KWs & to account for confounding factors such as age, sex, diet, and migration

Objective

- Characterize 317 analytes from 7 contaminant classes and diet (stable isotope: $\delta^{13}C$ and $\delta^{15}N$) in nine priority Chinook salmon stocks consumed by SRKW and NRKW to better understand contaminant exposure in R KWs

Methods



Sample Collection & Processing

- Chinook salmon samples were collected during 2018 & 2019 in the Fraser River estuary, southwest coast and east coast of Vancouver Island in partnership with Albion Test Fishery, Pacheedaht First Nation, and the University of British Columbia
- Muscle tissue subsampled from heads for contaminants and stable isotopes ($\delta^{13}C$, $\delta^{15}N$)

Stock Identification and Selection

- Genetic stock identification via single nucleotide polymorphism (SNP) genotyping
- Based on SRKW & NRKW diet studies from DFO and NOAA the following nine priority Chinook stocks were selected for stable isotope and contaminant analyses:
 - East Coast Vancouver Island (ECVI), West Coast Vancouver Island (WCVI), Puget Sound, Columbia River, Skeena River, South Thompson River, Upper Fraser, Mid Fraser, and Harrison River (Lower Fraser)

Methods

Chemical Class	Uses	Status in Canada ¹³
PCBs	Electrical equipment, hydraulic systems	Banned in 1977
PBDEs	Flame retardants	Production banned; importation, use & sale regulated
OCPs	Pesticides	DDT & chlordanes banned in 1985 & 2003 respectively; others varying regulations
Dioxin/Furans	Pulp & paper mills	Elimination of use in progress
HBCDD	Brominated flame retardant	Banned in 2017
APs	Emulsifiers in detergents & pesticides	Amount of input into environment regulated
CPs	Plasticizers & flame retardants	Short-chain banned; MC & LC under evaluation

Contaminant Analyses

- 58 samples from 9 priority Chinook salmon stocks were analyzed for the following:
 - PCBs (159 congeners), PBDEs (40 congeners), OCPs (27 analytes) and dioxin/furans using GC-MR/MS
 - HBCDD and Alkylphenols (APs) using HPLC-MS/MS
 - Chlorinated paraffins (CPs) (short-, medium, and long-chain; 66 congeners) using high-resolution analytical methods: GC-HR/MS, LC-MS/MS, HPLC, and UPLC-MS/MS

Results




Figure 1. Sum PCBs, DDTs, and PBDEs concentrations (mean + SE) in nine priority Chinook stocks.

- PCBs and DDTs greatest concentrations in 7 out of 9 stocks:
- ECVI, Harrison, Upper Fraser, Mid Fraser, Puget Sound, Columbia, Skeena




Figure 2. Sum alkylphenols, sum chlordanes, and hexachlorobenzene concentrations (mean + SE) in nine priority Chinook stocks.

- Alkylphenols greatest concentrations in South Thompson and WCVI followed by PCBs

Results




Figure 3. $\delta^{13}C$ and $\delta^{15}N$ values (mean \pm SE) for nine priority Chinook stocks consumed by NRKW and SRKW. $\delta^{13}C$ values were lipid corrected based on Larocque et al., 2021.¹²

Discussion and Conclusions

- ECVI and Harrison Chinook stocks are feeding at higher trophic levels and utilizing more coastal food webs than other priority stocks (Fig. 3);
- Seven stocks are feeding at lower trophic levels and are utilizing more open ocean food webs;
- Despite the banning of PCB and DDT 44 and 36 years ago respectively, they are still being found as the dominant contaminants in priority Chinook stocks consumed by SRKW and NRKW;
- This is first study to report on contaminant concentrations for 317 analytes in eight contaminant classes in adult Pacific salmon Chinook stocks;
- PCBs and DDTs likely pose a continued risk to SRKW and NRKW health and may pose a health risk to some resident Chinook stocks;
- These preliminary results will be used in conjunction with fish and marine mammal toxicity thresholds to prioritize/rank COCs for Chinook salmon and resident killer whales;
- Future work will investigate the significance of the varying stable isotope values and their relationship to contaminant concentrations.

Acknowledgements

The authors would like to thank our partners at the University of British Columbia, Albion Test Fishery, DFO, Pacheedaht First Nations, and the anglers who provided samples to the project. We would especially like to thank Helen Jones at Pacheedaht First Nation for partnering with us and coordinating logistics of sample collection. This work was supported by the Government of Canada's *Whales Initiative*, DFO's Species At Risk Program, Ocean Wise Conservation Association, and Mitacs. We authors gratefully acknowledge Brown Lab (DFO) team members for processing the Chinook salmon heads, SGS AXYS Analytical Ltd. for running the contaminant analyses, and the Fisk Lab for running the stable isotope analyses.

References

- Research, C. F. Southern Resident Killer Whale Population (accessed April 7).
- Holbert, S., A. Recovery Strategy for the Southern and Northern Resident Killer Whales (Orcinus orca) in Canada. Fisheries and Oceans Canada, 2008.
- Towers, J. Photo-identification catalogue and status of the southern resident killer whale population in 2011. Fisheries and Oceans Canada, 2015.
- Ford, R. B., Wright, M. J., Ellis, M. J., and Cunniff, J. R. Chinook salmon produced by resident killer whales: assessed and regional productivity, and identity of prey, and consumption rates. Canadian Science Advisory Secretariat: Secretariat scientifique canadien de consultation, 2010.
- Hansen, M. B., Baskett, R. W., Ford, J. K. B., Hightower, B. A., Van Doren, D. M., Conroy, J. E., Emmerson, C. K., Schorr, G. S., Gilmore, B. C., Brown, K. C., Baker, K. C., Baker, B. A., Soren, J. G., Ford, M. J., Species and stock identification of prey consumed by southern resident killer whales in their summer range. *Environ Biol Fish*, 2019.
- Hansen, M. B., Emmerson, C. K., Conroy, J. E., Brown, K. C., Hightower, B. A., Van Doren, D. M., Schorr, G. S., Jacobsen, J. K., Soren, M. S., Soren, J. G., Baskett, R. W., Baskett, L. Endangered predators and endangered prey: Survival identification of prey consumed by southern resident killer whales in their summer range. *Environ Biol Fish*, 2019.
- Department of Environment and Climate Change Canada. *Environmental and Climate Change Canada: Environment and Climate Change Canada: Environment and Climate Change Canada*. Ottawa, ON: Environment and Climate Change Canada, 2020.
- Canada. *Climate Change Canada*. Ottawa, ON: Environment and Climate Change Canada, 2020.
- Brown, T. M., Ross, P. S., Baskett, R. W., Holbert, S. J., Ford, A. T., Hightower, B. A., PCB related effects threshold as derived through prey processing profiles to locally contaminated degraded wild (Pau Hupia). *Environmental science & technology*, 2014, 48 (21), 12952-12961.
- Edwards, P. R., Price, S. Health risk based evaluation of emerging pollutants in killer whales (Orcinus orca) prey: setting a support of recovery. Canada, D. A. D., Ed. 2017.
- Nick, M., Brown, T. M., Contaminants as a Conservation Threat to Marine Mammals. In *Biochemistry Conservation: Addressing Modern Challenges Through Multidisciplinary & Forensic Approaches*, Adams, H. B., Ed. Springer, 2020.
- Larocque, S. M., Ford, M. J., & Johnson, T. B. (2021). Evaluation of trophic level structure and food web for killer whales and their prey: analysis in adult networks. *Applied Conservation in Marine Systems*, 1(1), 1-10.
- Government of Canada. "Management of Toxic Substances." December 11, 2020. <https://www.canada.ca/en/environment-climate-change/services/management-toxic-substances.html>