



Apr 27th, 4:30 PM - 5:00 PM

## Microchemical techniques to evaluate priority contaminant sources along the migration routes of Chinook (*Oncorhynchus tshawytscha*) and Coho Salmon (*Oncorhynchus kisutch*)

Micah Quindazzi  
*Uvic*

Follow this and additional works at: <https://cedar.wwu.edu/ssec>

---

Quindazzi, Micah, "Microchemical techniques to evaluate priority contaminant sources along the migration routes of Chinook (*Oncorhynchus tshawytscha*) and Coho Salmon (*Oncorhynchus kisutch*)" (2022). *Salish Sea Ecosystem Conference*. 164.  
<https://cedar.wwu.edu/ssec/2022ssec/allsessions/164>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).

# Microchemical techniques to evaluate priority contaminant sources along the migration routes of Chinook (*Oncorhynchus tshawytscha*) and Coho Salmon (*Oncorhynchus kisutch*)

Micah Quindazzi<sup>a</sup>, Tanya Brown<sup>b</sup> and Francis Juanes<sup>a</sup>

a. University of Victoria

b. Department of Fisheries and Oceans



## Introduction

Tracking the marine migrations of salmon is important for understanding marine survival and contaminant profiles<sup>1,2,3</sup>. These considerations are important both for the health of salmon populations<sup>4,5</sup>, as well as their predators<sup>6,7</sup>. Current methods of tracking these migrations are constrained by costs, methods of retrieval, and need to individually tag salmon. Intrinsic tags can overcome these problems<sup>8,9</sup>.

## Questions and Hypotheses

1. Is otolith microchemistry an effective way of tracking marine migrations of Chinook and Coho Salmon? I hypothesize that this method will be effective due to overall differences in water chemistry and physiological conditions between regions.
2. Does early marine growth determine the marine migration life history displayed by the individual? I hypothesize that faster growing fish will more likely become residents.

## Methods

- Coho baselines captured from Strait of Georgia (SoG) and West Coast Vancouver Island (WCVI) in 2018
- Coho collected from hatchery facilities across Southern BC (Big Qualicum, Chilliwack, and Quinsam)
- Conducted laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) to collect element and trace element data
  - N=20 per baseline region; N=52 for BQ; N=68 for CH; N=58 for Q
- Scale circuli spacing measured for first 10 circuli post marine check for early marine growth measurement




## Statistical Analysis

- Utilized Random Forest (RF) models to assign salmon to different marine regions
- Created PERMANOVA model to compare marine migration life history type against early marine growth rate, population, origin, and sex

## References

1. Cullon et al., 2009, *Env. Toxicol. Chem.*, 28;
2. Ruff et al., 2017, *Fish. Oceanogr.*, 26;
3. Sobocinski et al., 2021, *Ecol. Indic.*, 124;
4. Debruyne et al., 2004, *Env. Sci. Tech.*, 38;
5. Monod, 1985, *B. Env. Contam. Tox.*, 35;
6. Cullon et al., 2012, *Env. Toxicol. Chem.*, 31;
7. O'Neill et al., 2020, *Sci. Total Env.*, 712;
8. Solomon et al., 2006, *CJFAS*, 63;
9. Hampton et al., 2018, *J. Mar. Sys.*, 188

## Results






## Conclusions

- Otolith microchemistry can differentiate marine migration life history types
- Early marine growth does not influence marine migration life history type overall, seems to be population specific

## Future Directions

- Analyzing thousands of Chinook and Coho otoliths from a broader range of marine regions
- Conducting stable isotope measurements to add in classification success
- Looking at other endogenous effects like estuary entry and estuary residency time
- Collections of Chinook muscle tissue to determine contaminant profiles of different marine regions across the Northeast Pacific Ocean



## Acknowledgements

Funders: Pacific Salmon Foundation, Mitacs, International Year of the Salmon, AFS WA-BC Chapter

Collaborators: Nuu-chah-nulth Tribal Council, Department of Fisheries and Oceans, Alaska Department of Fish and Game, Washington Department of Fish and Wildlife, Will Duguid, Emma Polard, Lydia Walton, Zoe Molder, and Leigh Gaffney