



Western Washington University
Western CEDAR

Salish Sea Ecosystem Conference


2022 Salish Sea Ecosystem Conference
(Online)

Apr 26th, 1:30 PM - 3:00 PM

Forecasting central Salish Sea Dungeness crab populations: A study in patience

Sarah Grossman

Follow this and additional works at: <https://cedar.wvu.edu/ssec>

 Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Grossman, Sarah, "Forecasting central Salish Sea Dungeness crab populations: A study in patience" (2022). *Salish Sea Ecosystem Conference*. 186.
<https://cedar.wvu.edu/ssec/2022ssec/allsessions/186>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.



Forecasting central Salish Sea Dungeness crab populations: A study in patience

Sarah K. Grossman, Claire E. Cook, Julie S. Barber - Swinomish Indian Tribal Community

Salish Sea Ecosystem Conference April 26, 2022

Dungeness crab in Puget Sound

Outdoors | Sports

Unlike elsewhere, crabbing prospects solid in the Puget Sound region

Originally published December 2, 2015 at 7:03 pm | Updated December 2, 2015 at 7:13 pm

While pretty much the entire coast of southern Oregon and northern California wallows in sorrow about Dungeness crab fishing closures, it is quite the opposite around Puget Sound where fishing remains excellent.

By Mark Yuasa
Seattle Times staff reporter

Share story

Puget Sound crabs are plentiful, delicious and easy to catch. Here's how.


Originally published July 20, 2016 at 4:17 pm | Updated July 25, 2016 at 1:06 pm



1 of 2 | Nelson Nakata, of Seattle, inspects a crab. (Mark Yuasa/The Seattle Times)

It could be another record season for the Dungeness fishery in Western Washington waters.

By Mark Yuasa
Seattle Times staff reporter



Commercial Crab Management Regions and Areas. Crab Management Region 3-4 is new for the 2021-2022 season.
WDFW PS Crab Team

Dungeness crab in Puget Sound

Key Peninsula News

The Voice of the Key Peninsula

LOCAL NEWS • FEATURES • OPINION • MILESTONES • COMMUNITY • CALENDAR •


Crabbing Season Closed in South Sound

Posted Sunday, July 1, 2018 4:50 am


Sara Thompson, KP News

In May the Washington state Department of Fish and Wildlife announced that south Puget Sound is closed this season for all crab harvesting, including Dungeness and red rock crabs.

"A combination of trends in crab population over the last five years and some recent scientific papers and presentations about crab survival led to this decision," said Robert Sizemore, Puget Sound shellfish manager for the WDFW. "Harvest reports from south Puget Sound have been dismal."



Courtesy: Washington State Department of Fish and Wildlife




Commercial Crab Management Regions and Areas. Crab Management Region 3-4 is new for the 2021-2022 season.

WDFW PS Crab Team

Factors influencing recruitment to the fishery

Long-term goal: Develop a model to better forecast Dungeness crab harvest in Swinomish management regions

Q: What effect does timing and initial size at settlement have on recruitment to the fishery



Larval Dungeness crab methods

Monitor daily abundance of megalopae (catch/hour) from April to ~September

Measure CW/CH/TH for up to 30 Dungeness crab megalopae per week

*Quantify other crab species

Megalopae = late-stage larvae;

CW = carapace width; CH = carapace height; TH = total height

Early cohort = ~April to mid-June;

Late cohort = mid-June to ~August



Juvenile intertidal methods


Survey beaches monthly

Measure CW/CH for crab >4 mm to ~ 40 mm¹


Plus random molts and larger roaming crab

J1 = Juvenile instar stage 1; J2 = juvenile instar stage 2

¹ Juvenile Dungeness crab occupy intertidal habitats until ~ 40 mm CW.




Larval results: Annual variability 2018 to 2021




Larval abundances varied by up to 12x between years and sites

| Region 2E | | | |
|------------------|------------------|-------------------------|--------------------------|
| Recruitment Year | Megalopae Caught | Commercial Fishing Year | Commercial Landings, lbs |
| 2018 | 20,592 | 2021/2022 | 1,954,000 |
| 2019 | 19,740 | 2022/2023 | ??? |
| 2020 | 8,427 | 2023/2024 | |
| 2021 | 100,480 | 2024/2025 | |

Larval results: Temporal variability 2018 to 2021




Larval results: Temporal variability 2018 to 2021



Megalopae sizes

Why is timing of
delivery important?


1.2 mm decrease in mean
monthly CW April to August



Intertidal J1 sizes

Size at settlement tracks
with larval sizes, over time

1.7 mm
decrease in
monthly mean
CW, May to
August




Intertidal juvenile growth: Early vs. late cohorts

EC instars that settle in May reach 40 mm by the fall of year 1

LC instars overwinter in the intertidal, reach 40 mm in spring of year 2


All instars of the previous settlement year emigrate from the intertidal by May/June



Juvenile growth: continued

EC settlers reach sexual maturity by year 2: ~ 1 to 2 years earlier than LC settlers

LC molt 1-2 times more than EC before sexual maturity




Summary

What effect does timing and initial size at settlement have on recruitment to the fishery?

EC settlers grow to harvest limit in ~3.5-4.5 years vs. LC in ~5-6 years

To date, majority of larvae delivered to central Salish Sea are LC

Orensanz and Gallucci (1988): San Juan Island, WA; MacKay and Weymouth (1935): Boundary Bay, B.C.; Armstrong et al. (1989): Grays Harbor, WA



This work would not be possible without support from

Swinomish Senate and Fish and Game Committee, Tandy Wilbur, and all of the Fisheries Shellfish Team

Pacific Northwest Crab Research Group, Deception Pass State Park, Port of Anacortes, and our wonderful volunteers from the Salish Sea Stewards and the Sound Water Stewards

In memory of Lorraine Loomis, who encouraged and enthusiastically supported our research



U.S. Environmental Protection Agency #PA-01J27601, Bureau of Indian Affairs #A19AP00216 , U.S. Fish and Wildlife Service #F18AP00618

