



Apr 27th, 4:30 PM - 5:00 PM

## 2021 Mapping of Microplastics in Surface Sediments of Elliot Bay near Seattle, WA to Determine Impacts on Benthic Communities

Sade McNight  
*University Of Washington Tacoma*

Olivia Kirby  
*University Of Washington Tacoma*

Follow this and additional works at: <https://cedar.wwu.edu/ssec>

---

McNIGHT, Sade and Kirby, Olivia, "2021 Mapping of Microplastics in Surface Sediments of Elliot Bay near Seattle, WA to Determine Impacts on Benthic Communities" (2022). *Salish Sea Ecosystem Conference*. 244.

<https://cedar.wwu.edu/ssec/2022ssec/allsessions/244>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).

# 2021 MAPPING OF MICROPLASTICS IN SURFACE SEDIMENTS OF ELLIOT BAY NEAR SEATTLE, WA TO DETERMINE IMPACTS ON BENTHIC COMMUNITIES




Olivia Kirby, Julie Masura, and Cheryl Greengrove – University of Washington Tacoma

## MICROPLASTICS PROBLEM



In 2018, 359 million tonnes of plastics were produced worldwide, with 59% being common polymers (i.e., polyethylene, polypropylene, polyvinyl chloride; [PlasticsEurope 2019](#)). The rate of input of ocean plastic is estimated to be approximately 9.5 million tonnes per year. Primary plastics are those manufactured at size for making larger plastics, and secondary plastics are those that have broken down in the environment. Size-categories for plastics are macroplastics (> 5 mm) and microplastics (< 5 mm).

## RESEARCH PARTNERSHIP

This project explores microplastics in sediments collected from a fine resolution sampling project in 2021 of Elliot Bay near Seattle, WA to create baseline observations and determine if plastic pollution in sediments have changed over time. Washington State Department of Ecology's Marine Sediment Monitoring Group has provided sediment samples to analyze for microplastics since 2014. Thirty-seven stations were sampled using a 0.1 m<sup>2</sup> stainless steel Van Veen grab sampler to recover 2-3 cm of the top sediment from the seabed.





2021 microplastics concentration for Puget Sound and Elliot Bay.




## UWT MICROPLASTICS QUICK FACTS

- > Every station contained microplastic
- > 92% of the micro-debris are fibers
- > 56% of the micro-debris are 1-2.5 mm
- > Over 800 micro-debris were clear followed by ~300 blue



## UNDERGRADUATE RESEARCH

Each year the sediment samples are processed by undergraduate researchers for grain-size distribution, total organic content percentage, harmful algae abundance, and microplastic concentration as part of a summer research experience. Microplastics were processed and analyzed through a series of density separations, sieving, and manual extractions. Each piece was characterized by type, color, and size.



## STUDENT TEAM

Abdullah Alammam  
Anna Finch  
Olivia Kirby  
Sade McKnight

For more information, contact [jmasura@uw.edu](mailto:jmasura@uw.edu)

