

7-1973

S-Algebras on Sets in C to the power of n

Donald R. Chalice

Western Washington University, donald.chalice@wwu.edu

Follow this and additional works at: https://cedar.wwu.edu/math_facpubs



Part of the [Mathematics Commons](#)

Recommended Citation

Chalice, Donald R., "S-Algebras on Sets in C to the power of n " (1973). *Mathematics*. 16.
https://cedar.wwu.edu/math_facpubs/16

This Article is brought to you for free and open access by the College of Science and Engineering at Western CEDAR. It has been accepted for inclusion in Mathematics by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

S-ALGEBRAS ON SETS IN C^n

DONALD R. CHALICE

ABSTRACT. We give conditions which are necessary and sufficient for polynomial approximation of any continuous function on a compact subset of C^n .

Let X be a compact set in C^n , complex n -space, $P(X)$ the uniform closure of the polynomials on X , $C(X)$ all continuous functions on X , m_{2n} $2n$ -dimensional Lebesgue measure on C^n , and for any λ in C^n let $E(\lambda) = \{z \in C^n \mid z_i = \lambda_i \text{ for some } i\}$.

A given set is a *strong peak set* if it is an intersection of peak sets and meets the boundary of each of them in a set which contains no nonempty perfect subsets. We say a Banach algebra A is an *S-algebra* if when x is in A and \hat{x} , the Gelfand transform of x , vanishes at some p , then there exist x_n in A such that \hat{x}_n vanish in (perhaps different) neighborhoods of p and $\|x_n - x\| \rightarrow 0$. For example, for any locally compact abelian group G , $L^1(G)$ is an *S-algebra* [6, p. 51]. The main question which motivates us here is: If A is a uniform algebra on a compact space X and A is an *S-algebra*, does $A = C(X)$? Our main result is the following.

THEOREM. *A necessary and sufficient condition that $P(X) = C(X)$ is that (i) $P(X)$ is an S-algebra, (ii) for almost all $\lambda \in C^n$ with respect to m_{2n} , $E(\lambda) \cap X$ is a strong peak set, and (iii) each point of X is a peak point for $P(X)$.*

We begin with some observations about uniform algebras which are *S-algebras*.

LEMMA 1. *Let A be a uniform algebra on a compact space X and suppose that A is an S-algebra. Then: (i) The maximal ideal space of A is X . (ii) A is normal. (iii) If each point of X is a peak point then A satisfies condition D [4, p. 86], i.e. if $f \in A$ and $f(p) = 0$ then there exist $f_n \in A$ vanishing on neighborhoods of p such that $f_n f \rightarrow f$.*

PROOF. (i) Let p be a homomorphism on A and μ_p a representing measure for p with minimal closed support. If μ_p is not a point-mass then

Presented to the Society, January 18, 1972 under the title *A condition for polynomial approximation in C^n* ; received by the editors December 6, 1971 and, in revised form, October 18, 1972.

AMS (MOS) subject classifications (1970). Primary 46J10; Secondary 41A10.

© American Mathematical Society 1973

some $q \neq p$ lies in its closed support. Find f in A such that $f(p)=1$ and $f(q)=0$. Since A is an S -algebra we can assume that f vanishes in a neighborhood of q . Thus $f\mu_p$ is a complex representing measure for p , and since it dominates a (positive) representing measure for p [3, p. 33], we have a contradiction to the minimality of μ_p .

(ii) By part (i), to show normality of A we need only show regularity. But if $p \neq q$ then as above there is an f in A such that f vanishes on a neighborhood of p and $f(q)=1$. If K is compact and $q \notin K$ then by compactness one finds a function f in A such that $f=0$ on K and $f(q)=1$.

(iii) Suppose k peaks at p . Then there exist g_n in A such that g_n vanish on neighborhoods of p such that $\|g_n - (1 - k^n)\| \rightarrow 0$. Hence, $\|f - fg_n\| \leq \|f(1 - k^n - g_n)\| + \|fk^n\| \rightarrow 0$ so that $fg_n \rightarrow f$.

Part (iii) allows us to do spectral synthesis on the maximal ideal space of any uniform S -algebra as follows.

LEMMA 2. *Let A be a uniform algebra which is an S -algebra on X and let I be a closed ideal of A . If each point of X is a peak point for A then I contains every element f in A such that $\partial\{x|f(x)=0\} \cap \text{hull}(I)$ contains no nonempty perfect set.*

PROOF. Since A is normal and satisfies condition D, this is immediate from [4, p. 86].

We shall also need the following lemma which generalizes a result in [7] from one variable. A detailed proof is given in [1].

LEMMA 3. *Let X be a compact set in C^n and let μ be a regular bounded Borel measure on X . Let*

$$\hat{\mu}(z) = \int \frac{d\mu(\lambda)}{(\lambda_1 - z_1) \cdots (\lambda_n - z_n)}$$

and

$$N_\mu(z) = \int \frac{d|\mu|(\lambda)}{|\lambda_1 - z_1| \cdots |\lambda_n - z_n|}.$$

Then $N_\mu(z) < \infty$ a.e. with respect to m_{2n} and if $\hat{\mu}(z) = 0$ a.e. m_{2n} then $\mu = 0$.

PROOF OF THE THEOREM. Let $E_1(X) = \bigcup \{E(z) | z \in X\}$. Let μ be a measure on X such that $\mu \perp P(X)$. We must show that $\mu = 0$. Now clearly if $z \notin E_1(X)$ then $\hat{\mu}(z) = 0$. Now call $E(X)$ the set of z for which $E(z) \cap X$ is a strong peak set and for which $N_\mu(z) < \infty$. Since this only differs from $E_1(X)$ by a set of measure 0, we need only show that $\hat{\mu}$ vanishes on $E(X)$. Now if $\lambda \in E(X)$, we know that $E(\lambda) \cap X = \bigcap_{i=1}^\infty K_i$ with k_i peaking on K_i and $E(\lambda) \cap \partial K_i$ contains no nonempty perfect subset. Note that the hull of the closed ideal generated by $(z_1 - \lambda_1) \cdots (z_n - \lambda_n)$ is $E(\lambda) \cap X$ so that, by Lemma 2, $1 - k_i^{n_i} \in$ the uniform closure of $P(X)(z_1 - \lambda_1) \cdots (z_n - \lambda_n)$

for any positive n_i . Now choose n_i so that $k_i^{n_i} \rightarrow \chi_{E(\lambda)}$ boundedly pointwise on X . Then find g_j in $P(X)$ such that $\|g_j(z_1 - \lambda_1) \cdots (z_n - \lambda_n) + 1 - k_j^{n_j}\| \rightarrow 0$. In other words, $f_j = 1 + g_j(z_1 - \lambda_1) \cdots (z_n - \lambda_n) \rightarrow \chi_{E(\lambda)}$ boundedly pointwise on X . Since $N_\mu(\lambda) < \infty$, $|\mu|$ vanishes on $E(\lambda)$. Also as $j \rightarrow \infty$,

$$\frac{f_j}{(\lambda_1 - z_1) \cdots (\lambda_n - z_n)} \rightarrow 0$$

pointwise on $X - E(\lambda)$, and dominatedly. Hence

$$\hat{\mu}(\lambda) = \int \frac{f_j}{(\lambda_1 - z_1) \cdots (\lambda_n - z_n)} d\mu \rightarrow 0 \text{ as } j \rightarrow \infty,$$

so $\hat{\mu}(\lambda) = 0$. Thus $\hat{\mu} = 0$ a.e. and, by Lemma 3, $\mu = 0$ and the theorem is proved.

For a uniform algebra A and a point x in $M(A)$, the maximal ideal space of A , call the 0-germ at x the set of functions in A which vanish on a neighborhood of x . We close with an example of a uniform algebra A such that for each point x in a dense set in $M(A)$ the 0-germ is dense in the maximal ideal determined by x . In other words the S -algebra condition is satisfied on at least a dense subset. McKissick [5] has proved the following.

LEMMA 4. *Let D be the open unit disk. Then there is a sequence $\{a_k\}$ in D , $0 < |a_k| \leq |a_{k+1}| \rightarrow 1$, such that for any $\epsilon' > 0$ there is a sequence $\{J_k\}$ of open disks in D centered at $\{a_k\}$ respectively such that:*

- (1) $\sum_1^\infty \text{length}(\partial J_k) < \epsilon'$.
- (2) *There exist rational functions r_n with poles at a_1, \dots, a_n such that $r_n \rightarrow f$ uniformly on $(\bigcup_{k=1}^\infty J_k)'$ and $f = 0$ on D' while $f(0) = 1$.*

Using the above lemma we prove the following.

LEMMA 5. *Let $c = |a_1|/2$. There is a constant $M > 0$ such that for any positive ϵ, δ there is a δ' and $\{D_k\}$ a sequence of open disks in $N(0, \delta'/c) - N(0, \delta'c)$ such that:*

- (1) $\sum_1^\infty \text{length}(\partial D_k) < \delta'c$.
- (2) *There exist rational functions $\{r_n\}$ with poles in $D_1 \cup \dots \cup D_n$ such that $r_n \rightarrow g$ uniformly on $(\bigcup_{k=1}^\infty D_k)'$ and*
 - (i) $|g| \leq M$ on $(\bigcup_1^\infty D_k)'$,
 - (ii) $g = 0$ on $N(0, \delta')$,
 - (iii) $|1 - g| < \epsilon$ on $N(0, \delta)'$.

In fact if f is the function obtained by Lemma 1 with ϵ' a fixed constant (to be determined) independent of ϵ and δ , then δ' can be chosen as $\delta\delta(\epsilon)$ where $\delta(\epsilon)$ is a function such that $|z| < \delta(\epsilon)$ implies $|1 - f(z)| < \epsilon$.

PROOF. For disks $\{J_k\}$ which we now choose in D let $\{D_k\}$ be their respective images under the map $1/cz$. Since $|a_k| \geq 2c$, by taking a sufficiently small ϵ' we can choose the open disks J_k guaranteed by Lemma 1

so that $z \in \bigcup J_k$ implies $|z| > c$ and so that $\sum \text{length}(\partial D_k) < 1$. Thus $D_k \subset N(0, 1/c^2) - N(0, 1)$ for all k . Let f denote the limit on $(\bigcup J_k)'$ of the rational functions guaranteed by Lemma 1, and let M be the maximum of f on this set. Now since $f(0) = 1$, $|z| < \delta(\epsilon)$ implies $|1 - f(z)| < \epsilon$. Set $\delta' = \delta\delta(\epsilon)$ and let $g(z) = f(\delta'/zc)$. Then redefining D_k as $\delta' D_k$ we have $D_k \subset N(0, \delta'/c^2) - N(0, \delta')$, $g(z)$ is obviously defined for $z \notin D_k$, and

- (1) $\sum \text{length}(\partial D_k) < \delta'$,
- (2) (i) $|g| < M$ on $(\bigcup D_k)'$,
- (ii) $g(z) = 0$ on $N(0, \delta'/c)$ since $|\delta'/zc| > 1$ there, and
- (iii) $|1 - g(z)| < \epsilon$ on $N(0, \delta'/c)$ since $|\delta'/zc| < \delta(\epsilon)$ there.

The statement of the lemma follows by replacing δ in the above by δc .

COROLLARY. *There is a constant M such that given positive δ', ϵ there exist D_k and g as in the above lemma satisfying (1) and (2) if δ is taken as $\delta'/\delta(\epsilon)$.*

Of course the above lemmas hold with 0 replaced by any point p . Also since the function $f(z) = \sum_{k=1}^{\infty} 1/[\phi'(a_k)(z - a_k)]$ used by McKissick in Lemma 1 has a $\delta(\epsilon) < \beta\epsilon$ for some fixed β and small enough ϵ we see that $\delta(\epsilon)$ in the above statements can be replaced by ϵ . We now construct the example. Pick $m > 1$ such that $2^m c > 1$. Let $X_{m-1} = D$ and $S_{m-1} = \phi$. Define $S_n \subset X_n, \{D_k^{j,n}\}$, for $n \geq m$ inductively as follows. Suppose that $S_{n-1} = \{a_1, \dots, a_k\}$. Choose other points a_{k+1}, \dots, a_i in X_{n-1} so that each point of X_{n-1} is within $1/2^n$ of some a_i , and let $S_n = \{a_1, \dots, a_i\}$. Let d denote the minimum distance between the points of S_n . Letting $\delta = \epsilon = d/(2^{n+j}c^{1/2})$ find $\{D_k^{j,n}\}_{k=1}^{\infty}$ open disks in $N(a_j, \delta\epsilon/c) - N(a_j, \delta\epsilon c)$ such that $\sum_{k=1}^{\infty} \text{length}(\partial D_k^{j,n}) < d^2/4^{n+j} < 1/2^{n+j}$ and (2) holds. Let $X_n = X_{n-1} - \bigcup_{k,j} D_k^{j,n}$. Observe that since $\delta\epsilon/c < d$ we have $S_n \subset X_n$. Note too that $\sum_{k,j=1}^{\infty} \text{length}(\partial D_k^{j,n}) < 1/2^n$ so that if we set $X = \bigcap_{n=m}^{\infty} X_n$, we have excised a countable number of discs whose boundaries have total length < 1 . Thus by Lemma 1 of [5], $R(X) \subsetneq C(X)$. It is now clear that given any $\epsilon > 0$ and any a_j some $N(a_j, d/(2^{n+j}c^{1/2})) \subset N(a_j, \epsilon)$ so there is a g in $R(X)$ so that $\|g\| \leq M$, g vanishes on a neighborhood of a_j and $|1 - g| < \epsilon$ on $N(a_j, \epsilon)'$. Thus the 0-germ at a_j is pointwise boundedly dense in the maximal ideal at a_j and so is dense. Since the $\{a_j\}$ are a dense subset of X the example has the required properties.

Can the example be altered so that it is an S -algebra? One's first inclination is to cover the disk by smaller and smaller δ'_n neighborhoods given by the Corollary, but clearly it is not possible to do this and even retain $\sum \delta'_n < \infty$. However the example is rather simple-minded in that the same function is used over and over. Perhaps a choice of other functions will extend the example. Some questions raised by the above are: (1) If the 0-germ at p is dense in the maximal ideal determined by p , is p a peak

point? (2) Is the example normal? (3) From an example of Cole (see also Basener [2]), it is well known that (iii) alone is not sufficient to imply the conclusion of the theorem. Are any of the hypotheses of the theorem redundant?

Wilken [8] has shown that if a uniform algebra A is an S -algebra on $[0, 1]$ then $A=C[0, 1]$. In closing we also show the following.

THEOREM. *If A is a uniform algebra and A is an S -algebra on the unit circle T , then $A=C(T)$.*

PROOF. Let p, q be peak points for A in T , so $\{p, q\}$ is a peak set. Let f in A peak there. Then there are g_n vanishing on neighborhoods of p and h_n vanishing on neighborhoods of q such that $\|(1-f^n)-g_n\| < 1/n$ and $\|(1-f^n)-h_n\| < 1/n$ with h_n and g_n in A . Then $\|(1-f^n)^2-h_n g_n\| < 5/n$. Let $k_n=0$ on one of the arcs $[p, q]$ joining p to q and let $k_n=h_n g_n$ on the other arc $[q, p]$. Then because A is normal and hence local, k_n are in A . But $k_n \rightarrow \chi_{(a,p)}$ boundedly pointwise. Thus if $\mu \in A^\perp$, $\mu_{(a,p)} = \mu_{[q,p]} \in A^\perp$. Hence $[q, p]$ is a peak set. Since every closed interval is an intersection of such peak sets, it follows that every closed set is a peak set and thus $A=C(T)$.

BIBLIOGRAPHY

1. D. R. Chalice, *A uniqueness property for measures on C^n* , Portugal. Math. (to appear).
2. R. F. Basener, *An example concerning peak points*, Notices Amer. Math. Soc. **18** (1971), 415-416. Abstract #71T-B60.
3. T. W. Gaelin, *Uniform algebras*, Prentice-Hall, Englewood Cliffs, N.J., 1969.
4. L. H. Loomis, *An introduction to abstract harmonic analysis*, Van Nostrand, Princeton, N.J., 1953. MR **14**, 883.
5. Robert McKissick, *A nontrivial normal sup norm algebra*, Bull. Amer. Math. Soc. **69** (1963), 391-395. MR **26** #4166.
6. Walter Rudin, *Fourier analysis on groups*, Interscience Tracts in Pure and Appl. Math., no. 12, Interscience, New York, 1962. MR **27** #2808.
7. J. Wermer, *Banach algebras and analytic functions*, Advances in Math. **1** (1961), fasc. 1, 51-102. MR **26** #629.
8. D. R. Wilken, *A note on strongly regular function algebras*, Canad. J. Math. **21** (1969), 912-914. MR **39** #6084.

DEPARTMENT OF MATHEMATICS, WESTERN WASHINGTON STATE COLLEGE, BELLINGHAM, WASHINGTON 98225