



May 2nd, 8:30 AM - 10:00 AM

A molecular framework to identify novel modes of action of endocrine disrupting compounds in shellfish

Mackenzie Gavery
University of Washington, mgavery@uw.edu

Steven Roberts
University of Washington

Follow this and additional works at: <https://cedar.wvu.edu/ssec>

 Part of the [Terrestrial and Aquatic Ecology Commons](#)


Gavery, Mackenzie and Roberts, Steven, "A molecular framework to identify novel modes of action of endocrine disrupting compounds in shellfish" (2014). *Salish Sea Ecosystem Conference*. 16.
<https://cedar.wvu.edu/ssec/2014ssec/Day3/16>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

The role of DNA methylation in mediating the effects of estrogens in oysters

Mackenzie Gavery & Steven Roberts
School of Aquatic and Fishery Sciences
University of Washington

Outline



17 α ethinylestradiol
(EE2)



Outline


- Background
 - DNA methylation
 - EDCs & bivalves
- Results
- Implications




17 α ethinylestradiol
(EE2)




Background




Background






Background










Background
















Source: Randy Jirtle



Reproduction in oysters

- Pacific oysters are sequential hermaphrodites
- Sex determination has a genetic component, but influenced by environmental factors



Reproduction in oysters

- Pacific oysters are sequential hermaphrodites
- Sex determination has a genetic component, but influenced by environmental factors

Estradiol

- induces sex reversal (Mori 1969)

17 α ethinylestradiol (EE2)

- \uparrow rate of oocyte development (Andrew 2010)

Nonylphenol

- offspring of exposed larvae had \uparrow intersex (Nice et al. 2003)



Hypotheses

- EE2 exposure will result in phenotypes such as skewed sex ratios and increased rate of gonad development
- DNA methylation patterns will be altered in oysters exposed to EE2

Hypotheses

- EE2 exposure will result in phenotypes such as skewed sex ratios and increased rate of gonad development
- DNA methylation patterns will be altered in oysters in upon exposure to EE2

Estrogen Experiment



500 ng/L EE2: 150 oysters (n=50/tank)



Control: 150 oysters (n=50/tank)

Estrogen Experiment



500 ng/L EE2: 150 oysters (n=50/tank)



Control: 150 oysters (n=50/tank)



Day 0



Day 7



Day 60

- Samples: histology, gonad tissue

Estrogen Experiment



500 ng/L EE2: 150 oysters (n=50/tank)



Control: 150 oysters (n=50/tank)



Day 0




Day 7





Day 60

- Samples: histology, gonad tissue

Results: Day 60 sex determination




Results: Day 60 size of females



Results: Day 60

size of males



Results: DNA methylation



EE2 (500ng/L) 150 oysters (n=50/tank)



control 150 oysters (n=50/tank)



Day 0



Day 7



Day 60

Results: DNA methylation



EE2 (500ng/L) 150 oysters (n=50/tank)



control 150 oysters (n=50/tank)



Day 0




Day 7



Day 60

Results: DNA methylation



Results: DNA methylation

- Results:
 - 45 differentially methylated regions (DMR)
 - DMRs were located in 38 different genes

Protein names

5-hydroxytryptamine receptor 1B
ATP-binding cassette sub-family G member 1
Angiotensin-converting enzyme
Neuronal acetylcholine receptor subunit alpha-6
Anaphase-promoting complex subunit 1
Arrestin domain-containing protein 3
Calmodulin
Corticotropin-releasing factor receptor 2
Carnosine synthase 1
E3 ubiquitin-protein ligase DTX3L
Dynein gamma chain, flagellar outer arm
Elongator complex protein 2
Ryncolin-1
Glutamine synthetase
Glutaredoxin 3
Granulins
Translation factor Guf1, mitochondrial
Apoptosis inhibitor IAP
Interferon-induced protein 44
Kelch-like protein 24
Liprin-beta-1
Low-density lipoprotein receptor-related protein 6
Unconventional myosin-Vb
NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial
Nose resistant to fluoxetine protein 6
Peptidase M20 domain-containing protein 2
60 kDa SS-A/Ro ribonucleoprotein
Solute carrier family 28 member 3
Solute carrier family 45 member 3
Protein transport protein Sec16A
Small integral membrane protein 14
Src kinase-associated phosphoprotein 2-B
DNA topoisomerase 1
tRNA pseudouridine synthase A, mitochondrial
Vasorin
Vacuolar protein sorting-associated protein 13C
WASH complex subunit 7

Protein names
 5-hydroxytryptamine receptor 1B
 ATP-binding cassette sub-family G member 1
 Angiotensin-converting enzyme
 Neuronal acetylcholine receptor subunit alpha-6

Gene Ontology (GO Slim)	Count
transport	10
cell organization and biogenesis	8
other metabolic processes	7
signal transduction	6
protein metabolism	5
RNA metabolism	5
developmental processes	4
cell cycle and proliferation	3
death	2
stress response	2
cell-cell signaling	1
DNA metabolism	1

Small integral membrane protein 14
 Src kinase-associated phosphoprotein 2-B
 DNA topoisomerase 1
 tRNA pseudouridine synthase A, mitochondrial
 Vasorin
 Vacuolar protein sorting-associated protein 13C
 WASH complex subunit 7

Protein names
 5-hydroxytryptamine receptor 1B
 ATP-binding cassette sub-family G member 1
 Angiotensin-converting enzyme
 Neuronal acetylcholine receptor subunit alpha-6

Gene Ontology (GO Slim)	Count
transport	10
cell organization and biogenesis	8
other metabolic processes	7
signal transduction	6
protein metabolism	5
RNA metabolism	5
developmental processes	4
cell cycle and proliferation	3
death	2
stress response	2
cell-cell signaling	1
DNA metabolism	1

Small integral membrane protein 14
 Src kinase-associated phosphoprotein 2-B
 DNA topoisomerase 1
 tRNA pseudouridine synthase A, mitochondrial
 Vasorin
 Vacuolar protein sorting-associated protein 13C
 WASH complex subunit 7

- ATP-binding cassette protein
- Serotonin receptor
- Low density lipoprotein receptor
- Granulin

Summary

- EE2 treatment did not affect sex ratios, but exposed females were larger than controls
- DMRs were identified within 1 week of EE2 exposure
- Genes with DMRs are functionally diverse (e.g. growth, immune, reproduction)



Implications

- DNA methylation may play a role in mediating responses to EDCs in bivalves
- Epigenetic marks may provide early indicators of EDC exposure in aquatic species

Acknowledgements

- Roberts Lab:
Steven Roberts
Samuel White
Emma Timmins-Schiffman
Claire Ellis
Brent Vadopalas
Jake Heare
- Taylor Shellfish:
Joth Davis
Molly Jackson
- Irv Shultz (Battelle PNNL)

